

gram. [Research support by FAPESP (grant 2015/20986-3), Tairana Artificial Insemination Station and Botupharma.]

<https://doi.org/10.1016/j.anireprosci.2018.04.056>

## AB50

### Evaluation of Percoll PLUS as a cushion solution during single layer centrifugation of fresh bull semen: Effects on frozen/thawed spermatozoa motility

**Alicio Martins Jr.**\*, Tairini E. da Cruz, Fernanda N. Marqui, Diego G. de Souza, Tatiana I.U. Berton

São Paulo State University, FMVA, Araçatuba, São Paulo State University, FMVZ, Botucatu, Tairana Artificial Insemination Station, Presidente Prudente, Master Fertility Animal Reproduction, Araçatuba, Brazil

E-mail address: [amartins@fmva.unesp.br](mailto:amartins@fmva.unesp.br) (A. Martins Jr.).

Traditional semen centrifugation can pack spermatozoa on bottom of conical centrifuge tube. Thus, this study was carried out to verify the influence of Percoll PLUS (PP) as a cushion solution during single layer centrifugation (SLC) of fresh bull semen on sperm motility parameters. Eighteen ejaculates from 3 Nellore bulls (1 of each bull/trial) were pooled and divided into 3 groups, as follows: no centrifuged sperm (NC), cushioned centrifugation (CS; with PP as commercially available) and non-cushioned centrifugation (NCS) both during SLC. SLC was performed by layered 1 billion of spermatozoa on top of 9-ml column of PP (70%) followed by centrifugation (839 × g) for 13 min at room temperature. Then, the supernatant was discarded and the sperm pellet diluted in freezing extender to a final concentration of  $24 \times 10^6$  spermatozoa/straw. After cooling and freezing sperm samples were thawed in a water bath (37 °C/30 s) and assessed for total motility (TM, %), progressive motility (PM, %), curvilinear velocity (VCL,  $\mu\text{m/s}$ ), straight line velocity (VSL,  $\mu\text{m/s}$ ), average path velocity (VAP,  $\mu\text{m/s}$ ), amplitude of lateral head displacement (ALH,  $\mu\text{m}$ ), beat cross frequency (BCF, Hz), linearity (LIN, %), straightness (STR, %), and wobble (WOB, %) using computer assisted sperm analyzer (CASA). For statistical analysis ANOVA and Tukey test were used (values expressed as mean  $\pm$  SD) with  $P < 0.05$  taken as significant. Higher percentage for PM ( $51 \pm 8.6$  and  $49.8 \pm 7.1$ ), STR ( $78 \pm 0.02$  and  $79 \pm 0.04$ ), and WOB ( $72 \pm 0.03$  and  $72 \pm 0.04$ ) were found, respectively for CS and NCS than for NC ( $26.5 \pm 1.7$ ,  $65 \pm 0.04$ , and  $64 \pm 0.03$ , respectively). No difference was observed among groups for the other sperm motility parameters, except for ALH which was better for CS and NCS. In conclusion, cushioned centrifugation with PP during SLC yielded similar results as compared to non-cushioned centrifugation, but these findings suggest that PP could be a substitute for other cushion solutions commonly used. [Acknowledgements: FAPESP (grant 2015/20986-3), Tairana AI Station, Botupharma Ltda, Master Fertility Animal Reproduction, Brazil.]

<https://doi.org/10.1016/j.anireprosci.2018.04.057>

## AB51

### Addition of iodixanol in bull freezing extender improves the sperm membranes integrity

Fernanda Nunes Marqui<sup>1,2,3,\*</sup>, **Alicio Martins Jr.**<sup>1,2,3</sup>, Tairini Erica da Cruz<sup>1,2,3</sup>, Tatiana Issa Uherara Berton<sup>1,2,3</sup>, Camila de Paula Freitas-Dell'Aqua<sup>1,2,3</sup>, José Antonio Dell'Aqua Júnior<sup>1,2,3</sup>, Eunice Oba<sup>1,2,3</sup>

<sup>1</sup> São Paulo State University, Botucatu, São Paulo, Brazil

<sup>2</sup> São Paulo State University, Araçatuba, São Paulo, Brazil

<sup>3</sup> Tairana Artificial Insemination Station, Presidente Prudente, São Paulo, Brazil

E-mail addresses: [fernanda\\_nunes.37@hotmail.com](mailto:fernanda_nunes.37@hotmail.com)

(F.N. Marqui), [amartins@fmva.unesp.br](mailto:amartins@fmva.unesp.br) (A. Martins Jr.).

This study was carried out to verify the effects of different concentrations of iodixanol added to the freezing extender on plasma and acrossomal membrane integrity of post-thawed sperm bull. Therefore, ejaculates from three Nellore bulls ( $n = 18$ ) were pooled and extended in a commercial freezing medium (egg yolk-sugar-glycerol) supplemented with iodixanol at concentrations of 0% (control group), 2.5% (I-2.5), 5% (I-5), and 10% (I-10). Subsequently, semen samples were cooled for 5 h at 4 °C, filled in 0.50 ml straws and frozen in a programmable freezer machine. After that, the straws were stored in liquid nitrogen until sperm evaluation, which were performed by thawing the samples in a water bath at 37 °C/30 s. Plasma and acrossomal membrane integrity (PAMI) were simultaneously assessed using Propidium Iodide and FITC-PSA probes, respectively, while translocation of phosphatidylserine (TPS) was identified by Annexin V and plasma membrane destabilization (PMD) by YO-PRO-1. All sperm samples were analyzed by flow cytometry. ANOVA and Tukey's test were used for statistical analysis (data of six replicates), with  $P < 0.05$  taken as significant. Higher percentage ( $P < 0.05$ ) of MPAL was observed in I-10 ( $65 \pm 1.4$ ) than for other groups, with control group showing a lower value ( $56.7 \pm 2.1$ ), while I-2.5 ( $59.8 \pm 2$ ) did not differ from control and I-5 ( $62 \pm 1.6$ ) groups. Regarding to TPS, higher percentage of spermatozoa without TPS (considering intact membrane cells) was observed in I-10 ( $61.2 \pm 1.4$ ) with lower percentage found in control group ( $45.6 \pm 2.4$ ). However, I-2.5 ( $56.3 \pm 1.9$ ) and I-5 ( $57.7 \pm 1.5$ ) exhibited similar results but differed significantly from the other groups. For PMD the values of I-10 and I-5 were no different ( $67.8 \pm 1.9$  and  $67 \pm 1.9$ , respectively), but they were higher than in control ( $62.4 \pm 2.7$ ) and I-2.5 ( $61 \pm 1.9$ ) groups, both with similar results. In conclusion, the addition of iodixanol to the bovine freezing extender significantly improves the sperm membrane integrity in a dose-dependent manner. [Acknowledgements: CAPES,