

RESEARCH ARTICLE

Baroreflex regulation affects ventilation in the cururu toad *Rhinella schneideri*

Lucas A. Zena^{1,2,*}, Glauber S. F. da Silva^{1,2}, Luciane H. Gargaglioni^{1,2} and Kênia C. Bícigo^{1,2,*}

ABSTRACT

Anurans regulate short-term oscillations in blood pressure through changes in heart rate (f_H), vascular resistance and lymphatic f_H . Lung ventilation in anurans is linked to blood volume homeostasis by facilitating lymph return to the cardiovascular system. We hypothesized that the arterial baroreflex modulates pulmonary ventilation in the cururu toad *Rhinella schneideri*, and that this relationship is temperature dependent. Pharmacologically induced hypotension (sodium nitroprusside) and hypertension (phenylephrine) increased ventilation (25°C: 248.7 ± 25.7 ml kg⁻¹ min⁻¹; 35°C: 351.5 ± 50.2 ml kg⁻¹ min⁻¹) and decreased ventilation (25°C: 9.0 ± 6.6 ml kg⁻¹ min⁻¹; 35°C: 50.7 ± 15.6 ml kg⁻¹ min⁻¹), respectively, relative to control values from Ringer solution injection (25°C: 78.1 ± 17.0 ml kg⁻¹ min⁻¹; 35°C: 137.7 ± 15.5 ml kg⁻¹ min⁻¹). The sensitivity of the ventilatory response to blood pressure changes was higher during hypotension than during hypertension [25°C: -97.6 ± 17.1 versus -23.6 ± 6.0 breaths min⁻¹ kPa⁻¹; 35°C: -141.0 ± 29.5 versus -28.7 ± 6.4 breaths min⁻¹ kPa⁻¹, respectively; negative values indicate an inverse relationship between blood pressure and ventilation (or breathing frequency), i.e. as blood pressure increases, ventilation decreases, and vice versa], while temperature had no effect on these sensitivities. Hyperoxia (30%; 25°C) diminished ventilation, but did not abolish the ventilatory response to hypotension, indicating a response independent of peripheral chemoreceptors. Although there are previous data showing increased f_H baroreflex sensitivity from 15 to 30°C in this species, further increases in temperature (35°C) diminished f_H baroreflex gain (40.5 ± 5.62 versus $21.6 \pm 4.64\%$ kPa⁻¹). Therefore, besides an involvement of pulmonary ventilation in matching O₂ delivery to demand at higher temperatures in anurans, it also plays a role in blood pressure regulation, independent of temperature, possibly owing to an interaction between baroreflex and respiratory areas in the brain, as previously suggested for mammals.

KEY WORDS: Baroreceptors, Chemoreceptors, Lung ventilation, Hyperoxia, Autonomic blockade, Anuran amphibians

INTRODUCTION

Across vertebrates, the negative feedback mechanisms that drive short-term blood pressure regulation rely on mechanoreceptors (baroreceptors) located on the wall of the arterial system that detect fluctuations in blood pressure. The afferent baroreceptor

information is then relayed to and integrated in the central nervous system (CNS), where baroreflex buffering capacity is achieved by reflexively altering heart rate (f_H) and vascular resistance (Altimiras et al., 1998; Bianchi-da-Silva et al., 2000; Sandblom and Axelsson, 2005; Hedrick et al., 2015; Zena et al., 2015, 2016). Besides cardiovascular responses, the baroreflex also influences respiratory function in mammals (for a review, see McMullan and Pilowsky, 2010). Although there are some suggestive data for baroreflex modulation of the respiratory system in crocodilians and amphibians (Van Vliet and West, 1986; Altimiras et al., 1998; Hedrick et al., 2013), there is still no direct evidence of a respiratory component in the reflex regulation of blood pressure in non-mammalian vertebrates.

Amphibians' cardiorespiratory homeostatic mechanisms for monitoring blood pressure and blood gases involve sensory information from baroreceptors and chemoreceptors, respectively, located in their major arterial vessels: the pulmocutaneous artery, the carotid artery and the aortic arch (Bianchi-da-Silva et al., 2000; Reyes et al., 2014). While the identity of these receptors has already been described, their relative contributions to cardiorespiratory control remain under investigation. Arterial partial pressure of oxygen homeostasis is maintained by peripheral chemoreceptor regulation of breathing (Wang et al., 1994; Branco and Glass, 1995), whereas regulation of blood pressure by arterial baroreceptors is adjusted by changing vascular resistance and cardiac frequency. Furthermore, anurans possess lymph hearts (pulsatile organs), which are under baroreflex control, that function to pump lymph into the venous system (Crossley and Hillman, 1999).

Besides their role in gas exchange, amphibian lungs are associated with some ecophysiological functions including vocalization, buoyancy and defensive behavior (Hillman et al., 2009; Jared et al., 2009). In addition, lung ventilation has been directly linked to vertical lymph mobilization and, hence, blood volume homeostasis (Hedrick et al., 2007; Hillman et al., 2010). By transmitting pressure and volume changes to the surrounding lymph sacs, pulmonary ventilation in anuran amphibians facilitates lymph movement against gravitational forces towards dorsally located lymph hearts, thereby preventing lymph accumulation (Hedrick et al., 2013). As a component of the lymphatic system in anurans, dorsally located anterior and posterior lymphatic hearts participate in blood volume regulation by pumping lymphatic fluid generated by an increased net transcapillary flux back into the venous system (Crossley and Hillman, 1999). This is evident from the cane toad (*Rhinella marina*), which has an extraordinary ability to acutely compensate for high blood losses (78% of their initial blood volume), but destruction of both anterior and posterior lymph hearts critically impairs this ability, leading to hemoconcentration and death in few days (Baustian, 1988).

The cururu toad *Rhinella schneideri* is a widespread bufonid with a broad distribution in natural habitats in South America (Chaco, savanna-like Cerrado, and Atlantic Forest), and is also found in open

¹Department of Animal Morphology and Physiology, College of Agricultural and Veterinary Sciences, São Paulo State University, Jaboticabal, São Paulo 14884-900, Brazil. ²National Institute of Science and Technology in Comparative Physiology (INCT Fisiologia Comparada), Jaboticabal, São Paulo 14884-900, Brazil.

*Authors for correspondence (lucaszena@yahoo.com.br; keniacb@fcav.unesp.br)

© L.A.Z., 0000-0002-4422-4458

List of symbols and abbreviations

CNS	central nervous system
f_H	heart rate
F_{IO_2}	fraction of oxygen in the inspired air
f_R	breathing frequency
Pa_{O_2}	arterial partial pressure of oxygen
PAP	pulsatile arterial pressure
PE	phenylephrine
P_{MA}	mean arterial blood pressure
SNP	sodium nitroprusside
\dot{V}_E	total ventilation
V_T	tidal volume

and urban areas. Despite daily maximum air temperatures reaching over 35°C during the summer season in southeastern Brazil (CIIAGRO: <http://www.ciiagro.sp.gov.br/ciiagroonline/>), experiments have shown a preferred body temperature of around 23–27°C for this species (Bicego-Nahas et al., 2001; Guerra et al., 2008; Noronha-de-Souza et al., 2015). Therefore, in order to avoid potential damage due to extreme ambient temperatures, amphibians would seek places that maintain a more moderate humidity and temperature (Moreira et al., 2009; Noronha-de-Souza et al., 2015).

Based on the roles played by lung ventilation in blood volume homeostasis in anurans (Crossley and Hillman, 1999; Hedrick et al., 2007; Hillman et al., 2010, 2013) and in O_2 delivery to match metabolic demands, we tested the hypothesis that pulmonary ventilation contributes to the baroreflex responses to blood pressure imbalances in *R. schneideri*, and that this effect is modulated by changes in temperature. To this end, baroreflex function was assessed through pharmacological interventions on arterial blood pressure, i.e. the Oxford method, at 25 and 35°C. Heart rate buffering reflex responses were evaluated along with ventilation, breathing frequency and tidal volume. Considering that anuran peripheral chemoreceptors mainly adjust ventilation in response to changes in the partial pressure of arterial oxygen (Branco and Glass, 1995; Wang et al., 1994), toads were exposed to hyperoxia [fraction of oxygen in the inspired air (F_{IO_2})=0.30] to account for possible influences of peripheral chemoreceptors on the ventilatory response during hypotension at 25°C. Furthermore, ventilatory responses to hypotension were also tested after total autonomic blockade.

MATERIALS AND METHODS**Animals**

Cururu toads *Rhinella schneideri* (Werner 1894), of both sexes (159.5±7.7 g body mass), were collected in swampy areas in the Ribeirão Preto region of São Paulo State, Brazil (approximately 21°10'S and 47°48'W), and transported to our laboratory at the Department of Animal Morphology and Physiology, UNESP, Jaboticabal, Brazil (approximately 21°14'S and 48°17'W). All animals were maintained at 25°C on a 12 h:12 h light:dark cycle with free access to water from an artesian well and a basking area. They were housed in containers with coconut fiber as a substrate and tubes for hiding in, and were held under laboratory conditions for at least 3 weeks before experimentation. Animals were fed 2–3 times a week with captive-bred speckled cockroaches (*Nauphoeta cinerea*), which were dusted with calcium and vitamin D₃. All experiments were performed from September to March, which matches the activity season of this species (Bicego-Nahas et al., 2001; Glass et al., 1997). Animal collection was approved by the Brazilian environmental agency (SISBIO-ICMBio/no. 35484-1), and the study was conducted with the approval of our Institutional Animal

Care and Use Committee (CEUA-FCAV-UNESP; protocol no. 017204/12).

Surgical procedures

Procedures were performed as previously described (Zena et al., 2015). Briefly, anesthesia was performed through immersion in an aqueous 0.25% solution of 3-aminobenzoic acid ethyl ester (MS-222; Sigma, St Louis, MO, USA), buffered to pH 7.7 with sodium bicarbonate, for 10 min or more until animals lost the palpebral reflex. A polyethylene cannula (PE-50; Clay Adams, Parsippany, NJ, USA), filled with heparinized Ringer solution (100 i.u. ml⁻¹ heparin) was occlusively inserted into the iliac artery and sutured in place. The same procedure was used to cannulate the femoral vein (100 i.u. ml⁻¹ heparin in Ringer solution) for drug injections. Just after surgery, toads were treated with a prophylactic antibiotic (enrofloxacin, Flotril®, 5.0 mg kg⁻¹ s.c.; Schering-Plough) and an analgesic (flunixin meglumine, Banamine®, 1.0 mg kg⁻¹ s.c.; Schering-Plough) (Gentz, 2007; Smith, 2007). Each animal was then individually placed in a temperature-controlled chamber at 25°C in the experimental room, where it was allowed to recover for a minimum of 48 h without disturbance before the experimental procedure was begun.

Drugs

Phenylephrine hydrochloride (PE, agonist of α_1 -adrenergic receptors), sodium nitroprusside dihydrate (SNP, nitric oxide donor), atropine sulfate (antagonist of muscarinic receptors) and sotalol hydrochloride (antagonist of β -adrenergic receptors) were purchased from Sigma. All drugs were dissolved in amphibian Ringer solution (mmol l⁻¹: 46.9 NaCl, 21.0 KCl, 2.40 CaCl, 1.29 MgCl, 3.14 NaHCO₃).

Ventilation and arterial blood pressure measurements

Pulmonary ventilation (\dot{V}_E), tidal volume (V_T) and breathing frequency (f_R) were calculated from the recordings of breathing using the pneumotachography method described by Glass et al. (1978). A facemask was built for each toad from 1.0 mm-thick silicon sheets (Bio-Art Equipamentos Odontológicos, São Carlos, Brazil) heat-molded on a plaster cast of the toad's head. The pneumotachograph was attached to the mask and subsequently fixed to the animal's snout (see below), allowing airflow to be measured continuously. A direct relationship exists between laminar airflow and the pressure difference across this tube, which was monitored with a differential pressure transducer (MLT141Spirometer, PowerLab System, ADInstruments) connected to a data acquisition system that included specific application software (PowerLab System, ADInstruments/Chart Software, v7.3, Sydney, Australia). Calibrations were performed for each mask by injecting 1, 3 and 5 ml volumes of air through the pneumotachograph using a graduated syringe. The arterial cannulae were connected to a physiological pressure transducer (MLT844, ADInstruments), calibrated against a mercury column. The transducer was connected to a data acquisition system (PowerLab System, ADInstruments/Chart Software, v7.3) via a bridge amplifier (FE221, ADInstruments). f_H and mean arterial blood pressure (P_{MA}) were calculated from the pulsatile arterial pressure (PAP) recorded in real-time using the cyclic measurements tool from the Chart Software.

Experimental protocol

Toads were housed in an acrylic water-jacketed chamber kept at the experimental temperature of 25°C using a constant-temperature

circulating water bath (PolyScience 9112A11B programmable model 9112 refrigerated circulator). A facemask attached to the pneumotachograph was glued to the animal's snout using a polyether impression material (Impregum Soft, 3M ESPE, St Paul, MN, USA) 24 h after surgery. Experiments began 48 h after surgery and 24 h after facemask attachment. For experiments at high temperature, the toads were acclimated to 35°C for at least 6 h, beginning 48 h after surgery. The experimental chamber was continuously flushed with humidified room air, and the temperature inside was continuously measured using a temperature sensor (MLT415/M thermistor temperature sensor, ADInstruments). After 60 min of basal recordings of f_H , blood pressure and ventilation airflow, Ringer solution (0.4 ml kg⁻¹) was injected into the femoral vein in order to verify any influence of volume injection on all variables subsequently evaluated. Serial intravenous injections of increasing doses of PE and SNP were then performed (doses are given in 'Analysis and assessment of the baroreflex', below). All cardiorespiratory parameters were recorded throughout the experimental protocol. Between drug injections, cardiorespiratory parameters were always allowed to return to similar pre-injection values.

Five animals from the main experimental group described above had full autonomic blockade at 25°C, which was performed on the day after the baroreflex experiment. It consisted of injections of the β -adrenergic receptor antagonist sotalol (3.0 μ g kg⁻¹) and of the muscarinic receptor antagonist atropine (3.0 μ g kg⁻¹). Cardiorespiratory variables were recorded continuously in animals injected with SNP (100 μ g kg⁻¹) before and after full autonomic blockade.

Five additional animals were subjected to hyperoxia ($F_{I_{O_2}}=0.30$) to identify any possible involvement of the peripheral chemoreceptors in the respiratory responses to blood pressure

adjustments. For hyperoxia experiments, following the baroreflex protocol, animals were exposed to 30% O₂ for at least 30 min. After this interval, 100 μ g kg⁻¹ SNP injection was repeated in order to identify respiratory responses under hypotension in the hyperoxic condition.

Analysis and assessment of the baroreflex

Reflex responses of f_H were evaluated by altering blood pressure through serial injections of increasing doses of PE (5, 10, 25, 50 and 100 μ g kg⁻¹; eliciting blood pressure increases) and SNP (5, 10, 25, 50 and 100 μ g kg⁻¹; eliciting blood pressure decreases) at both 25 and 35°C. f_H was plotted against P_{MA} (kPa) data obtained for each animal at each temperature and fitted to a four-variable sigmoidal logistic function as described previously by Reid (1996):

$$f_H = \frac{(A - D)}{1 + (P_{MA}/C)^B} + D, \quad (1)$$

where A is maximum f_H and D is minimum f_H (beats min⁻¹) of the curve, B is the slope coefficient (beats min⁻¹ kPa⁻¹) and C is the P_{MA} at the midpoint of the f_H range ($P_{MA,50}$; kPa). The range of the baroreflex was calculated as the difference between A and D . The maximum gain of the curve (G_{50} ; beats min⁻¹ kPa⁻¹), which occurs when P_{MA} equals C , was determined by taking the first derivative of Eqn 1, resulting in the gain equation:

$$G_{50} = \frac{-B(A - D)}{4C}. \quad (2)$$

To allow meaningful comparisons between temperatures, the maximal gain was normalized as a percentage change in f_H per unit




Fig. 1. Ventilatory airflow, pulsatile arterial blood pressure (PAP) and heart rate (f_H) of the cururu toad *Rhinella schneideri* at 25°C. Original traces from one toad. The arrows indicate the time of intravenous injection of phenylephrine (PE; 50.0 μ g kg⁻¹) and sodium nitroprusside (SNP; 50.0 μ g kg⁻¹). The black line in the PAP signal represents mean arterial pressure (P_{MA}).




Fig. 2. Effect of pharmacological manipulations on P_{MA} , f_H , breathing frequency (f_R), tidal volume (V_T) and ventilation (\dot{V}_E) in *R. schneideri* at 25 and 35°C. Data are for intravenous injection of Ringer solution and serially increasing doses of PE (A–E) and SNP (F–J) at 25 and 35°C. Blue asterisks indicate a significant effect of drugs relative to Ringer solution at 25°C; red asterisks indicate a significant effect of drugs relative to Ringer solution at 35°C (two-way repeated measures ANOVA; $P < 0.05$; Holm–Šidák's *post hoc* test). Cardiovascular parameters, $N=9$; ventilatory parameters, $N=8$.

change in P_{MA} (% kPa^{-1}) relative to the minimum heart rate (D), as described by Altamiras et al. (1998):

$$G = \frac{100B}{D}. \quad (3)$$

Data analysis and statistics

Data are shown as means \pm s.e.m. The effect of temperature on baseline cardiorespiratory and baroreflex variables was analyzed through a Student *t*-test and Mann–Whitney *U*-test for parametric

and non-parametric data, respectively. The effects of Ringer solution and serial bolus injection of PE and SNP on P_{MA} , f_H , f_R , V_T and \dot{V}_E at 25 and 35°C were analyzed using a two-way repeated-measures ANOVA (factors: drugs and temperature). Different $F_{I_{O_2}}$ (0.21 and 0.30) before and after SNP-induced hypotension for \dot{V}_E and the effect of full autonomic blockade on hypotension induced by SNP were analyzed through two-way repeated-measures ANOVA. Regression analyses were used to evaluate temperature influences on f_R/P_{MA} and \dot{V}_E/P_{MA} relationships during hypotension and hypertension stimuli and a two-way ANOVA was used to

Table 1. Baseline cardiorespiratory and baroreflex variables derived from sigmoidal baroreflex–heart rate curves generated after serial bolus injections of phenylephrine and sodium nitroprusside (Eqn 1) at 25 and 35°C in the toad *Rhinella schneideri*

	25°C	35°C	P
Baseline variables			
P_{MA} (kPa)	3.95±0.31	3.83±0.15	0.73
f_H (beats min ⁻¹)	32.2±1.83	64.3±3.69	<0.001*
\dot{V}_E (ml kg ⁻¹ min ⁻¹)	62.0±11.5	143.3±15.9	0.001*
V_T (ml kg ⁻¹)	3.80±0.53	5.79±0.68	0.037*
f_R (breaths min ⁻¹)	18.3±3.95	25.2±1.96	0.14
Baroreflex variables			
Minimum f_H (beats min ⁻¹)	27.0±1.61	54.5±3.47	<0.001*
Maximum f_H (beats min ⁻¹)	62.3±1.45	96.1±4.94	<0.001*
f_H range (beats min ⁻¹)	35.3±2.74	41.6±3.48	0.17
R^2	0.93±0.01	0.94±0.02	0.48
Slope coefficient	10.9±1.62	11.3±2.02	0.93
$P_{MA,50}$ (kPa)	3.10±0.21	3.32±0.09	0.38
G_{50} (beats min ⁻¹ kPa ⁻¹)	30.8±4.49	38.3±9.80	1.00
Normalized gain (% kPa ⁻¹)	40.5±5.62	21.6±4.64	0.017*

P_{MA} , mean arterial blood pressure; f_H , heart rate; \dot{V}_E , ventilation; V_T , tidal volume; f_R , breathing frequency; R^2 , coefficient of determination; slope coefficient, slope of the linear portion of the curve; $P_{MA,50}$, mean arterial blood pressure at the midpoint of the curve; G_{50} , maximum gain.

Data are expressed as mean±s.e.m. Cardiovascular and baroreflex parameters, $N=9$; ventilatory parameters, $N=8$.

*Significant difference between temperatures.

compare slopes (pressure and temperature). In addition, the relationship between P_{MA} and ventilatory responses (f_R , V_T and \dot{V}_E) was binned into P_{MA} categories in order to evaluate the influence of temperature in those relationships (two-way ANOVA; pressure and temperature). In all ANOVA analyses, the differences among means were further assessed by Holm–Šidák *post hoc* tests and were considered significant when $P<0.05$. Data were tested for unequal variance and normality, and when necessary appropriate transformations were performed.

RESULTS

Mean baseline cardiorespiratory variables for toads at 25 and 35°C are shown in Table 1. No significant temperature effect on P_{MA} was observed ($P=0.73$), whereas f_H was significantly elevated at 35°C ($P<0.001$). Temperature significantly augmented resting \dot{V}_E ($P=0.001$) in toads, mainly by increases in V_T ($P=0.037$).

An original PAP trace of one toad at 25°C exhibiting a typical tachycardic reflex response to a reduction in P_{MA} after injection of SNP (50 µg kg⁻¹) is shown in Fig. 1. In addition, a parallel increase in ventilatory airflow was observed during hypotension. In contrast, the increase in P_{MA} after a bolus injection of PE (50 µg kg⁻¹) had a very small effect on f_H (small bradycardia) besides the diminished ventilatory airflow.

Effect of temperature on the blood pressure influence on cardiorespiratory variables

Serial injections of increasing doses of PE elevated P_{MA} (effect of treatment: $P<0.001$; $F_{5,102}=90.141$; Fig. 2A) and evoked a small decrease in f_H (effect of treatment: $P<0.001$; $F_{5,102}=20.049$; Fig. 2B) at both temperatures tested, while temperature had significant effects on f_H (effect of temperature: $P<0.001$; $F_{1,106}=59.548$; Fig. 2B), but not on P_{MA} (Fig. 2A). Along with reflexive effects on f_H , PE-induced hypertension caused a significant decrease in f_R independent of temperature (effect of treatment: $P<0.001$; $F_{5,90}=21.137$; Fig. 2C). In contrast, V_T was significantly reduced by injections of 25, 50 and 100 µg kg⁻¹ PE at

25°C only (effect of treatment: $P<0.001$; $F_{5,90}=5.951$; Fig. 2D). When temperature was increased to 35°C, V_T remained unchanged for all doses of PE that induced hypertension, but was elevated relative to values at 25°C (effect of temperature: $P<0.001$; $F_{1,94}=19.565$; Fig. 2D). Overall, increasing P_{MA} evoked parallel reductions in \dot{V}_E independent of temperature (effect of treatment: $P<0.001$; $F_{5,90}=17.783$; Fig. 2E), mainly due to decreases in f_R . Because V_T was unaffected by hypertension at 35°C, \dot{V}_E values were maintained elevated at 35°C relative to those at 25°C (effect of temperature: $P=0.002$; $F_{1,94}=13.858$; Fig. 2E).

In contrast to PE effects, serial injections of increasing doses of SNP decreased P_{MA} (interaction of temperature×treatment: $P=0.028$; $F_{5,102}=2.671$; Fig. 2F), causing a pronounced reflex increase in f_H (interaction of temperature×treatment: $P=0.049$; $F_{5,102}=2.343$; Fig. 2G). In addition, because high temperature significantly shifted baseline f_H to higher values (see Table 1), reflex tachycardia was also maintained elevated across all SNP-induced hypotension doses relative to 25°C (effect of temperature: $P<0.001$; $F_{1,106}=73.105$; Fig. 2G). Reductions of P_{MA} induced increases in f_R (effect of treatment: $P<0.001$; $F_{5,90}=16.858$; Fig. 2H) that were followed by increases in V_T at higher doses of SNP at 25°C, while SNP-induced hypotension evoked an increase in V_T at the lowest SNP dose at 35°C only (effect of treatment: $P=0.002$; $F_{5,90}=4.398$; Fig. 2I). Thus, regardless of temperature, SNP-induced hypotension effects on V_T and f_R in *R. schneideri* resulted in a significant increase in total ventilation (effect of treatment: $P<0.001$; $F_{5,90}=24.656$; Fig. 2J).

Effect of temperature on the relationship between P_{MA} , f_H and \dot{V}_E

Increasing the temperature from 25 to 35°C resulted in an upward shift in the relationship between P_{MA} and f_H (Fig. 3). The operating point (represented by baseline P_{MA} and f_H values) in this relationship was moved up mainly because of temperature effects on baseline f_H ($P<0.001$; Table 1 and Fig. 3). Despite a 10°C increment in temperature, the f_H baroreflex was not enhanced, as can be observed by similarities between absolute gain at 25 and 35°C ($P=1.0$; Table 1). Because of the significant effect of temperature on resting f_H , absolute gain was then




Fig. 3. Mean baroreflex relationship between P_{MA} and f_H in *R. schneideri* at 25 and 35°C. Curves were generated from the mean values of the curve-fitted variables for each animal (see Table 1). Baseline f_H and P_{MA} are represented by the blue triangle at 25°C ($N=9$) and the red triangle at 35°C ($N=9$). For statistical values, see baroreflex variables derived from sigmoidal baroreflex– f_H curves in Table 1.




Fig. 4. The effect of changing P_{MA} on f_R , V_T and \dot{V}_E in *R. schneideri* at 25 and 35°C. (A,B) f_R ; (C,D) V_T ; (E,F) \dot{V}_E . Baseline f_R , V_T , \dot{V}_E and P_{MA} are represented by blue circles at 25°C ($N=8$) and red circles at 35°C ($N=8$). Different symbols represent different animals.

normalized as a percentage change from minimum f_H per unit change in P_{MA} (kPa), which resulted in a significant reduction of the normalized gain ($P=0.017$; Table 1) at the higher temperature. In addition to temperature effects on resting f_H , minimum and maximum f_H were also significantly affected by temperature ($P<0.001$; Table 1 and Fig. 3).

The effect of changing P_{MA} on ventilatory responses (f_R , V_T and \dot{V}_E) is represented by the relationships presented in Figs 4–6. The overall variability dataset for both temperatures, showing the relationship between P_{MA} and f_R , V_T or \dot{V}_E , is depicted in Fig. 4. Breathing data from one representative toad are illustrated in Fig. 5, showing the slopes in f_R/P_{MA} and \dot{V}_E/P_{MA} during SNP-induced hypotension and PE-induced hypertension at 25 and 35°C; slopes were not presented for V_T/P_{MA} because there was a poor relationship for most animals, as can be observed in Fig. 4C,D. The sensitivity of

the response (slope) due to hypotension or hypertension was not affected by thermal changes for either f_R/P_{MA} ($P=0.84$) or \dot{V}_E/P_{MA} ($P=0.18$) but hypotension exhibited higher average slopes for f_R/P_{MA} ($P<0.001$; $F_{1,90}=40.369$) and \dot{V}_E/P_{MA} ($P<0.001$; $F_{1,30}=52.764$) relationships than hypertension at both temperatures (Table 2).

To test the specific responses of ventilatory variables to blood pressure changes under the two thermal conditions, the relationships between P_{MA} and f_R , V_T or \dot{V}_E were examined for P_{MA} data binned into 1.0 kPa intervals (Fig. 6). f_R was significantly elevated at 35°C for pressures higher than 4.0 kPa only (hypertension; effect of temperature: $P<0.001$; $F_{1,175}=20.174$; Fig. 6A), while \dot{V}_E was maintained at higher values at 35°C across all P_{MA} binned ranges from hypotension to hypertension (effect of temperature: $P<0.001$; $F_{1,175}=61.694$; Fig. 6C), mainly because of




Fig. 5. Representative data from one toad for the relationship between P_{MA} and f_R or \dot{V}_E at 25 and 35°C. (A) The effect of changing P_{MA} on f_R in the presence of SNP (continuous blue line, 25°C: $r^2=0.70$; $P=0.04$; continuous red line, 35°C: $r^2=0.60$; $P=0.07$) and PE (dotted blue line, 25°C: $r^2=0.42$; $P=0.16$; dotted red line, 35°C: $r^2=0.72$; $P=0.03$). (B) The effect of changing P_{MA} on \dot{V}_E in the presence of SNP (continuous blue line, 25°C: $r^2=0.77$; $P=0.02$; continuous red line, 35°C: $r^2=0.72$; $P=0.03$) and PE (dotted blue line, 25°C: $r^2=0.47$; $P=0.13$; dotted red line, 35°C: $r^2=0.72$; $P=0.03$).

changes in V_T (effect of temperature: $P<0.001$; $F_{1,175}=49.823$; Fig. 6B).

Effects of cholinergic and adrenergic receptor blockade on cardiorespiratory parameters

The effect of full autonomic blockade with atropine plus sotalol on P_{MA} , f_H and \dot{V}_E before and after SNP-induced hypotension in *R. schneideri* is shown in Fig. 7. Full autonomic blockade blunted the tachycardic reflex induced by hypotension (interaction of full autonomic blockade \times treatment: $P<0.001$; $F_{1,22}=48.265$; Fig. 7B); in contrast, hypotension-induced increases in \dot{V}_E were not affected by full autonomic blockade ($P=0.74$; Fig. 7C).

SNP-induced hypotension effects on ventilation under normoxia and hyperoxia

In toads exposed to hyperoxia ($F_{I_{O_2}}=0.30$), ventilation was significantly reduced ($P=0.032$; $F_{1,28}=3.177$; Fig. 8). Similar to animals under normoxic conditions, animals breathing 30% O_2 also expressed an increased ventilatory response ($P<0.001$; $F_{1,28}=96.581$; Fig. 8) after SNP-induced hypotension that did not differ from those values obtained from the group breathing normoxic gas ($P=0.6$).

DISCUSSION

Our finding that changes in lung ventilation are driven by pressure loading and unloading in the cururu toad *R. schneideri* provides evidence of an interesting interaction among baroreceptor reflexes, pulmonary ventilation and the lymphatic system in anuran amphibians (Burggren et al., 2013; Hedrick et al., 2013). Furthermore, in contrast to previous data showing increases in baroreflex sensitivity for the thermal interval between 15 and 30°C, an elevation in temperature to 35°C diminished f_H baroreflex sensitivity, but ventilatory responses to hypotension and hypertension were temperature independent.

Blood pressure homeostasis in anuran amphibians

As part of cardiovascular homeostasis in anuran amphibians, baroreceptors located in the arterial system buffer short-term blood pressure fluctuations reflexively via effectors that modulate f_H and peripheral vascular resistance. Given their high rate of lymph formation, approximately 10 times that of mammals (Desai, et al., 2010; Hillman et al., 2010), anurans have a highly efficient lymphatic system to maintain cardiovascular homeostasis. The importance of the two pairs of dorsally located lymph hearts in blood pressure homeostasis has been revealed by findings that

arterial baroreceptor loading is able to reflexively decrease lymphatic f_H , thereby decreasing lymph fluid return into the venous system (Yamane, 1990; Crossley and Hillman, 1999). In addition, lymph heart destruction results in a marked increase in red cell volume, interstitial edema and also death (Zwemer and Foglia, 1943; Baustian, 1988). Generated extravascular fluid in anuran amphibians returns to the cardiovascular system through a combination of changes in pressure and volume of the lymph sacs that is driven by effectors such as lung ventilation (inflation and deflation) and skeletal muscle contraction (Drewes et al., 2007; Hedrick et al., 2007). Therefore, from an evolutionary perspective, ventilatory responses that may be modulated by baroreceptor loading and unloading in *R. schneideri* can contribute to our understanding of a number of pieces of evidence for the influence of the cardiovascular system on the respiratory system in mammals (Brunner et al., 1982; Walker and Jennings, 1998; Baekey et al., 2010; McMullan and Pilowsky, 2010; Stewart et al., 2011). More interestingly, our findings add further support to a role for lung ventilation in blood pressure/volume homeostasis in anuran amphibians (Hedrick et al., 2013), although the relationship between pulmonary ventilation, baroreflex and lymphatic system in our toads is still correlational rather than mechanistic. Therefore, further studies are needed to clarify the real contribution of ventilation-driven pressure changes in lymph return.

The nucleus of the solitary tract is the major brainstem region receiving and integrating peripheral cardiovascular and respiratory afferent inputs in mammals (Lowey and Spyer, 1990). Baroreceptor afferents also converge towards the nucleus of the solitary tract in *R. schneideri* (Bianchi-da-Silva et al., 2000), but any other synaptic transmission in the CNS of amphibians related to baroreflex control is still unknown. High intensity stimulation of the recurrent laryngeal nerve, which innervates the pulmocutaneous baroreceptors, elicits cardiovascular adjustments such as bradycardia and hypotension in addition to an increased apnea period in awake *R. marina* toads (Van Vliet and West, 1986). This observation corroborates the hypertension-induced reduction in ventilation we see in our toads. Extensive overlap between respiratory and cardiovascular areas and the dose-dependent increase in pulse pressure gradually extending expiratory time in rats (*in situ* preparation) indicates the presence of a neural interaction between the baroreflex and respiratory patterns (Baekey et al., 2008, 2010). It is possible that anuran amphibian baroreceptor inputs into the nucleus of the solitary tract could cause suppression or release of the central respiratory drive, contributing to hypopnea and hyperpnea, respectively. Thus, relationships between blood pressure, ventilation and the lymphatic system may depend on




Fig. 6. Relationships between binned P_{MA} and f_R , V_T or \dot{V}_E in *R. schneideri* at 25 and 35°C. (A) f_R ; (B) V_T ; (C) \dot{V}_E . P_{MA} was grouped into six categories covering pressures from lower than 2.0 kPa to higher than 6.0 kPa; note that pressures between 2.0 and 6.0 kPa were binned into 1.0 kPa intervals. *Significant effect of temperature within binned P_{MA} interval (two-way ANOVA; $P < 0.05$; Holm–Šidák's *post hoc* test).

integrated information in the CNS; however, whether these circuits exist and/or are interacting with each other remains to be further investigated in amphibians.

One may argue that the respiratory response to hypotension results from peripheral chemoreflex activation, but evidence does not support this hypothesis. In toads, we know that the hypoxic ventilatory response is mediated by reductions in arterial partial pressure of oxygen (P_{aO_2}) (Wang et al., 1994; Branco and Glass, 1995). In this context, the discharge frequency of the carotid chemoreceptors in *R. marina* did not increase until P_{aO_2} fell between 5.3 and 8.0 kPa (40–60 mm Hg), and remained silenced in hyperoxic gas conditions (30% O_2 in inspired air; Van Vliet and West, 1992). In our animals, an increased F_{IO_2} (0.30) in the air decreased baseline \dot{V}_E and did not abolish the ventilatory response to hypotension (Fig. 8), suggesting little to no involvement of these chemoreceptors. In addition, hypoxic ventilatory responses in *R. schneideri*, in order to improve lung gas exchange, are mediated by altering V_T rather than f_R (Bicego-Nahas et al., 2001; Gargaglioni and Branco, 2001, 2003), while increases in ventilation elicited by hypotension in the present study were achieved through increases in f_R rather than V_T (Fig. 2). We also know that, at least in *R. marina*, carotid labyrinth chemoreceptors are not flow sensitive; afferent nerve activity is not affected by carotid labyrinth blood supply cessation, and the hypoxia stimulus is no longer effective in stimulating chemoreceptors in these conditions (Van Vliet and West, 1992). Thus, our data suggest that ventilatory responses during acute hypotension may instead be a consequence of baroreceptors influencing respiratory areas. Further support for this view comes from our results with hypertension, which induced the opposite response, i.e. ventilation decrease.

Temperature effects on cardiorespiratory reflex responses

In the present study, increasing the temperature to 35°C promoted cardiovascular and respiratory adjustments such as increases in baseline f_H and \dot{V}_E . \dot{V}_E was affected by temperature from 25 to 35°C (127% increase, $Q_{10}=2.27$), mainly as a result of increases in V_T ($Q_{10}=1.57$). Thus, increased \dot{V}_E at high temperature matches an elevated metabolic rate up to 105% (thermal interval range from 25 to 35°C) reported in a closely related species, *R. marina* (Overgaard et al., 2012). High temperature exposure (35°C) in *R. schneideri* elicited cardiovascular adjustments to support an increased metabolic demand in which baseline f_H rose 100% relative to that at 25°C, while blood pressure remained unchanged. Sustaining blood pressure relatively constant, for the thermal interval between 25 and 35°C, while resting f_H increased accordingly, likely reflects a general peripheral vasodilation (Zena et al., 2015), as stroke volume remains fairly constant (Hedrick et al., 1999).

Pharmacologically induced changes in blood pressure not only triggered baroreflex responses in f_H but also affected \dot{V}_E in *R. schneideri*, similar to what has been reported for mammals (Brunner et al., 1982; Walker and Jennings, 1998; McMullan and Pilowsky, 2010; Stewart et al., 2011). Increases in blood pressure in the toads were followed by reductions in \dot{V}_E , primarily through modification of f_R and to a lesser extent V_T ; conversely, \dot{V}_E was significantly increased during hypotension. In contrast to decreases in V_T during hypertension at 25°C (higher doses of PE), hypertension at 35°C was unable to decrease V_T , despite diminished f_R . In addition, the relationship between P_{MA} and the ventilatory responses (binned P_{MA} data; Fig. 6) revealed that at high temperature, toads exhibit elevated ventilation throughout the entire P_{MA} range (<2 to >6 kPa), mainly because of increased V_T . This permits the toads to keep an elevated ventilatory airflow in their lungs, ensuring sufficient gas exchange at the organ level during high metabolic demand, while their ability to change ventilation in response to baroreceptor loading and unloading is preserved.

Table 2. Regression analysis slopes for the relationships between P_{MA} and f_R or \dot{V}_E generated after serial bolus injections of phenylephrine (hypertension) and sodium nitroprusside (hypotension) at 25 and 35°C in the toad *R. schneideri*

Temperature (°C)	f_R slope (breaths min ⁻¹ kPa ⁻¹)		<i>P</i>	\dot{V}_E slope (ml kg ⁻¹ min ⁻¹ kPa ⁻¹)		<i>P</i>
	Hypotension	Hypertension		Hypotension	Hypertension	
25	−16.8±4.8	−7.1±2.1	<0.001*	−97.6±17.1	−23.6±6.0	<0.001*
35	−19.8±3.0	−4.8±0.7	<0.001*	−141.0±29.5	−28.7±6.4	<0.001*

*Significant difference between hypotension and hypertension slopes (Student’s *t*-test) within each temperature. f_R slope, slope relationship between f_R and P_{MA} ; \dot{V}_E slope, slope relationship between \dot{V}_E and P_{MA} . *N*=8.

(Table 2 and Fig. 5). In agreement with the ability of anurans to respond to hypotension primarily by increasing f_H (Hedrick et al., 2015; Zena et al., 2015), *R. schneideri* also exhibited a prominent ventilatory response to reductions rather than increases in blood pressure. This provides correlational data for the ability of anurans to use their lungs to regulate blood volume (Hedrick et al., 2013); in addition, it contributes to a prevailing view that amphibians strongly

defend short-term blood pressure imbalances against low pressure events (Hedrick et al., 2015; Zena et al., 2015).

Beyond temperature-independent sensitivity of the ventilatory reflex response, the baroreflex of f_H in our toads in fact diminished at 35°C, as seen by the normalized gain (Table 1). This suggests that the cardiac limb of the baroreflex response is possibly approaching its thermal limit. In a previous study (Zena et al., 2015), we showed that f_H baroreflex sensitivity is enhanced by temperature for a thermal interval between 15 and 30°C. Nevertheless, our data seem to indicate that further increases in temperature (above 30°C), instead of improving f_H baroreflex sensitivity, actually reduce it in our toads. A possible explanation for this may involve a thermal impairment of the tachycardic response to hypotension. This is indicated by a lower Q_{10} effect (1.5) for the maximum f_H response achieved for the highest dose of SNP. Baseline f_H was profoundly affected by temperature, shifting the operating point of the baroreflex upward (100% increase) with a Q_{10} effect of 2.0. Hence, in order to sustain f_H baroreflex sensitivity from 25 to 35°C, we expected that temperature would affect maximum and minimum f_H responses through the same Q_{10} effect as baseline f_H ; however, this was not the case for maximum f_H .

The ability to mount reflex f_H responses mainly to hypotension in anurans accounts for most of the f_H reflex responses due to changes in blood pressure (Hedrick et al., 2015; Zena et al., 2015) and can be strongly affected by the inability to further increase f_H when the animal has an already elevated baseline value. Such a situation can be encountered by the animal at very high temperatures (>35°C; Overgaard et al., 2012) or during enforced activity (Wahlqvist and Campbell, 1988); the latter is already known to reset the baroreflex to higher f_H and P_{MA} values in humans (Norton et al., 1999).

Overall it seems that anurans rely on a combination of mechanisms to maintain cardiovascular homeostasis. The baroreflex in anurans is clearly linked to the lymphatic system because lymph hearts are




Fig. 7. Effect of SNP on P_{MA} , f_H and \dot{V}_E in *R. schneideri* before and after full autonomic blockade with atropine and sotalol. Changes in P_{MA} (A), f_H (B) and \dot{V}_E (C) induced by SNP (100 μ g kg⁻¹) before (*N*=7) and after (*N*=5) full autonomic blockade with atropine (3.0 mg kg⁻¹) plus sotalol (3.0 mg kg⁻¹) at 25°C. Different lowercase letters indicate a significant effect of SNP injections; different uppercase letters indicate a significant effect of full autonomic blockade before or after SNP (two-way repeated measures ANOVA; *P*<0.05; Holm–Šidák’s *post hoc* test).




Fig. 8. Effect of SNP-induced hypotension on \dot{V}_E in *R. schneideri* under normoxia and hyperoxia. Different lowercase letters indicate a significant difference between normoxia and 30% hyperoxia; different uppercase letters indicate a significant difference between control and SNP injection within each gas condition (two-way repeated measures ANOVA; *P*<0.05; Holm–Šidák’s *post hoc* test). Normoxia, *N*=10; hyperoxia, *N*=5.

under feedback control of arterial baroreceptors (Crossley and Hillman, 1999) and pulmonary ventilation works as an efficient effector for lymph mobilization (Hedrick et al., 2007). Based on our data, short-term imbalances in blood pressure are defended by mechanisms that involve reflex f_H adjustments along with changes in pulmonary ventilation (mainly breathing frequency) that may essentially arise from afferent integration of baroreceptors in the CNS, modulating respiratory areas, as proposed for mammals (McMullan and Pilowsky, 2010). Furthermore, high environmental temperatures decrease the f_H baroreflex by impairing maximum f_H response effectiveness (lower Q_{10}) when baseline f_H is already elevated, while the ventilatory component of this response is preserved. Although the loop between ventilation and lymph flow based on baroreflex regulation deserves further experimentation, our present findings in *R. schneideri* clearly provide a correlational link among the baroreflex, lung ventilation and the lymphatic system in anurans, in addition to providing evolutionary insights for better understanding of the links between blood pressure and ventilation in mammals. Among amphibians, however, whether the role of lung ventilation is a conserved feature contributing to blood pressure homeostasis is still unknown, as some lungless amphibians like salamanders (Hutchison, 2008) and the Bornean flat-headed frog, *Barbourula kalmantanensis* (Bickford et al., 2008), may rely on other effectors for vertical lymph movement (i.e. skeletal muscle contraction; Drewes et al., 2007), and thus blood pressure/volume regulation dependent on ventilation certainly does not exist.

Acknowledgements

We thank Dr Lynn Hartzler who kindly reviewed our manuscript for English grammar and Dr Brent Foy for his contributions to our statistical analysis of P_{MA} and ventilatory response relationships. We are grateful to Mr Wagner Bianchini for allowing us to access his aquatic plants farm for toad collection.

Competing interests

The authors declare no competing or financial interests.

Author contributions

L.A.Z. conceived, designed and performed the experiments, and analyzed the data; L.A.Z., G.S.F.d.S., L.H.G. and K.C.B. interpreted the data; L.A.Z., K.C.B., G.S.F.d.S. and L.H.G. wrote the manuscript.

Funding

The present study was supported by the Fundação de Amparo à Pesquisa do Estado de São Paulo–FAPESP [08/57712-4 (INCTFisiologia Comparada); 2012/17379-0; 2nd year PhD fellowship to L.A.Z.] and Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES; 1st year PhD fellowship to L.A.Z.). This study was part of the activities developed by L.A.Z. during his PhD at the Joint Graduate Program in Physiological Sciences (PIPGCF) from UFSCar/UNESP and was presented at the Experimental Biology meeting April 2016 (FAPESP-Brazil 2012/17379-0 and APS-USA).

References

- Altamiras, J., Franklin, C. E. and Axelsson, M. (1998). Relationships between blood pressure and heart rate in the saltwater crocodile *Crocodylus porosus*. *J. Exp. Biol.* **224**, 2235–2242.
- Baekey, D. M., Dick, T. E. and Paton, J. F. R. (2008). Pontomedullary transection attenuates central respiratory modulation of sympathetic discharge, heart rate and the baroreceptor reflex in the in situ rat preparation. *Exp. Physiol.* **93**, 803–816.
- Baekey, D. M., Molkov, Y. I., Paton, J. F. R., Rybak, I. A. and Dick, T. E. (2010). Effect of baroreceptor stimulation on the respiratory pattern: Insights into respiratory-sympathetic interactions. *Respir. Physiol. Neurobiol.* **174**, 135–145.
- Baustian, M. (1988). The contribution of lymphatic pathways during recovery from hemorrhage in the toad *Bufo marinus*. *Physiol. Zool.* **61**, 555–563.
- Bianchi-da-Silva, L. M., Menescal-de-Oliveira, L. and Hoffmann, A. (2000). Baroreceptor control of heart rate in the awake toad: peripheral autonomic effectors and arterial baroreceptor areas. *J. Auton. Nerv. Syst.* **80**, 31–39.
- Bicego-Nahas, K. C., Gargaglioni, L. H. and Branco, L. G. S. (2001). Seasonal changes in the preferred body temperature, cardiovascular, and respiratory responses to hypoxia in the toad, *Bufo paracnemis*. *J. Exp. Zool.* **289**, 359–365.
- Bickford, D., Iskandar, D. and Barlian, A. (2008). A lungless frog discovered on Borneo. *Curr. Biol.* **18**, R374–R375.
- Branco, L. G. and Glass, M. L. (1995). Ventilatory responses to carboxyhaemoglobinaemia and hypoxic hypoxia in *Bufo paracnemis*. *J. Exp. Biol.* **198**, 1417–1421.
- Brunner, M. J., Sussman, M. S., Greene, A. S., Kallman, C. H. and Shoukas, A. A. (1982). Carotid sinus baroreceptor reflex control of respiration. *Circ. Res.* **51**, 624–636.
- Burggren, W. W., Christoffels, V. M., Crossley, D. A., Enok, S., Farrell, A., Hedrick, M. S., Hicks, J. W., Jensen, B., Moorman, A. F. M., Mueller, C. A. et al. (2013). Comparative cardiovascular physiology: future trends, opportunities and challenges. *Acta Physiol.* **197**, 257–276.
- Crossley, D. A., II and Hillman, S. S. (1999). The role of pulmocutaneous baroreceptors in the control of lymphatic heart rate in the toad *Bufo marinus*. *Physiol. Biochem. Zool.* **72**, 109–115.
- Desai, P., Williams, A. G., Jr., Prajapati, P. and Downey, H. F. (2010). Lymph flow in instrumented dogs varies with exercise intensity. *Lymphat. Res. Biol.* **8**, 143–148.
- Drewes, R. C., Hedrick, M. S., Hillman, S. S. and Withers, P. C. (2007). Unique role of skeletal muscle contraction in vertical lymph movement in anurans. *J. Exp. Biol.* **210**, 3931–3939.
- Gargaglioni, L. H. and Branco, L. G. (2001). Effect of nitric oxide in the nucleus isthmi on the hypoxic and hypercarbic drive to breathing of toads. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* **281**, R338–R345.
- Gargaglioni, L. H. and Branco, L. G. S. (2003). Role of glutamate in the nucleus isthmi on the hypoxia- and hypercarbia-induced hyperventilation of toads. *Respir. Physiol. Neurobiol.* **135**, 47–58.
- Gentz, E. J. (2007). Medicine and surgery of amphibians. *ILAR J.* **48**, 255–259.
- Glass, M. L., Wood, S. C. and Johansen, K. (1978). The application of pneumotachography on small unrestrained animals. *Comp. Biochem. Physiol. A Physiol.* **59**, 425–427.
- Glass, M. L., Fernandes, M. S., Soncini, R., Glass, H. and Wasser, J. S. (1997). Effects of dry season dormancy on oxygen uptake, heart rate, and blood pressures in the toad, *Bufo paracnemis*. *J. Exp. Zool.* **279**, 330–336.
- Guerra, A. R. G., Gargaglioni, L. H., Noronha-De-Souza, C. R., Abe, A. S., Branco, L. G. S. and Bicego, K. C. (2008). Role of central nitric oxide in behavioral thermoregulation of toads during hypoxia. *Physiol. Behav.* **95**, 101–107.
- Hedrick, M. S., Palioca, W. B. and Hillman, S. S. (1999). Effects of temperature and physical activity on blood flow shunts and intracardiac mixing in the toad *Bufo marinus*. *Physiol. Biochem. Zool.* **72**, 509–519.
- Hedrick, M. S., Drewes, R. C., Hillman, S. S. and Withers, P. C. (2007). Lung ventilation contributes to vertical lymph movement in anurans. *J. Exp. Biol.* **210**, 3940–3945.
- Hedrick, M. S., Hillman, S. S., Drewes, R. C. and Withers, P. C. (2013). Lymphatic regulation in nonmammalian vertebrates. *J. Appl. Physiol.* **115**, 297–308.
- Hedrick, M. S., McNew, K. A. and Crossley, D. A. (2015). Baroreflex function in anurans from different environments. *Comp. Biochem. Physiol. A Mol. Integr. Physiol.* **179**, 144–148.
- Hillman, S. S., Withers, P. C., Drewes, R. C. and Hillyard, S. D. (2009). *Ecological and Environmental Physiology of Amphibians*. Oxford: Oxford University Press.
- Hillman, S. S., Hedrick, M. S., Drewes, R. C. and Withers, P. C. (2010). Lymph flux rates from various lymph sacs in the cane toad *Rhinella marina*: an experimental evaluation of the roles of compliance, skeletal muscles and the lungs in the movement of lymph. *J. Exp. Biol.* **213**, 3161–3166.
- Hutchison, V. H. (2008). Amphibians: lungs' lift lost. *Curr. Biol.* **18**, R392–R393.
- Jared, C., Antoniazzi, M. M., Jordão, A. E. C., Silva, J. R. M. C., Greven, H. and Rodrigues, M. T. (2009). Parotid macroglands in toad (*Rhinella jimi*): Their structure and functioning in passive defence. *Toxicon* **54**, 197–207.
- Lowey, A. D. and Spyer, K. M. (1990). *Central Regulation of Autonomic Functions*. New York, NY: Oxford University Press.
- McMullan, S. and Pilowsky, P. M. (2010). The effects of baroreceptor stimulation on central respiratory drive: a review. *Respir. Physiol. Neurobiol.* **174**, 37–42.
- Moreira, L. a., Fenolio, D. B., Silva, H. L. R. and Silva, N. J. Jr. (2009). A preliminary list of the herpetofauna from termite mounds of the cerrado in the upper Tocantins river valley. *Pap. Avulsos Zool.* **49**, 183–189.
- Noronha-de-Souza, C. R., Bovo, R. P., Gargaglioni, L. H., Andrade, D. V. and Bicego, K. C. (2015). Thermal biology of the toad *Rhinella schneideri* in a semiarid environment in southeastern Brazil. *Temperature* **2**, 554–562.
- Norton, K. H., Boushel, R., Strange, S., Saltin, B. and Raven, P. B. (1999). Resetting of the carotid arterial baroreflex during dynamic exercise in humans. *J. Appl. Physiol.* **87**, 332–338.
- Overgaard, J., Andersen, J. L., Findsen, A., Pedersen, P. B. M., Hansen, K., Ozolina, K. and Wang, T. (2012). Aerobic scope and cardiovascular oxygen transport is not compromised at high temperatures in the toad *Rhinella marina*. *J. Exp. Biol.* **215**, 3519–3526.
- Reid, I. A. (1996). Angiotensin II and baroreflex control of heart rate. *News Physiol. Sci.* **11**, 270–274.
- Reyes, C., Fong, A. Y., Brink, D. L. and Milsom, W. K. (2014). Distribution and innervation of putative arterial chemoreceptors in the bullfrog (*Rana catesbeiana*). *J. Comp. Neurol.* **522**, 3754–3774.

- Sandblom, E. and Axelsson, M.** (2005). Baroreflex mediated control of heart rate and vascular capacitance in trout. *J. Exp. Biol.* **208**, 821–829.
- Smith, S. A.** (2007). Compendium of drugs and compounds used in amphibians. *ILAR J.* **48**, 297–300.
- Stewart, J. M., Rivera, E., Clarke, D. A., Baugham, I. L., Ocon, A. J., Taneja, I., Terilli, C. and Medow, M. S.** (2011). Ventilatory baroreflex sensitivity in humans is not modulated by chemoreflex activation. *Am. J. Physiol. Heart Circ. Physiol.* **300**, H1492–H1500.
- Van Vliet, B. N. and West, N. H.** (1986). Cardiovascular responses to electrical stimulation of the recurrent laryngeal nerve in conscious toads. *J. Comp. Physiol. B* **156**, 363–375.
- Van Vliet, B. N. and West, N. H.** (1992). Functional characteristics of arterial chemoreceptors in an amphibian (*Bufo marinus*). *Respir. Physiol.* **88**, 113–127.
- Wahlqvist, B. Y. I. and Campbell, G.** (1988). Autonomic influences on heart rate and blood pressure in the toad, *Bufo marinus*, at rest and during exercise. *J. Exp. Biol.* **134**, 377–396.
- Walker, J. K. L. and Jennings, D. B.** (1998). Respiratory effects of pressor and depressor agents in conscious rats. *Can. J. Physiol. Pharmacol.* **76**, 707–714.
- Wang, T., Branco, L. G. and Glass, M. L.** (1994). Ventilatory responses to hypoxia in the toad *Bufo paracnemis* before and after a decrease in haemoglobin oxygen-carrying capacity. *J. Exp. Biol.* **186**, 1–8.
- Yamane, M.** (1990). The arterial baroreceptor-lymph heart inhibitory reflex and its pathway in the bullfrog *Rana catesbeiana*. *Kawasaki Med. J.* **16**, 93–100.
- Zena, L. A., Gargaglioni, L. H. and Bicego, K. C.** (2015). Temperature effects on baroreflex control of heart rate in the toad, *Rhinella schneideri*. *Comp. Biochem. Physiol. A Mol. Integr. Physiol.* **179**, 81–88.
- Zena, L. A., Dantonio, V., Gargaglioni, L. H., Andrade, D. V., Abe, A. S. and Bicego, K. C.** (2016). Winter metabolic depression does not change arterial baroreflex control of heart rate in the tegu lizard *Salvator merianae*. *J. Exp. Biol.* **219**, 725–733.
- Zwemer, R. and Foglia, V.** (1943). Fatal loss of plasma volume after lymph heart destruction in toads. *Exp. Biol. Med.* **53**, 14–17.