



Notes: The authors declare no conflict of interest.

In vitro versus *in vivo* protein digestibility techniques for calculating PDCAAS (protein digestibility-corrected amino acid score) applied to chickpea fractions



Olga Luisa Tavano^{a,*}, Valdir Augusto Neves^b, Sinézio Inácio da Silva Júnior^c

^a Federal University of Alfenas, Nutrition Faculty, 700 Gabriel Monteiro da Silva St, Alfenas, MG 37130-000, Brazil

^b Food and Nutrition Department, School of Pharmaceutical Sciences, São Paulo State University, Araraquara-Jaú Road, Km 01, CEP 14801-902 Araraquara, São Paulo State, Brazil

^c Food and Medicines Department, School of Pharmaceutical Sciences, Federal University of Alfenas, Gabriel Monteiro da Silva Street, 700, CEP: 37130-000, Alfenas, Minas Gerais State, Brazil

ARTICLE INFO

Article history:

Received 22 August 2016

Received in revised form 4 October 2016

Accepted 5 October 2016

Available online 6 October 2016

Keywords:

Protein digestibility

In vitro digestibility

PDCAAS

Chickpea

Protein fractions

ABSTRACT

Seven different *in vitro* methods to determine the protein digestibility for chickpea proteins were considered and also the application of these methodologies for calculating PDCAAS (protein digestibility-corrected amino acid score), seeking their correlations with the *in vivo* methodology. *In vitro* digestibility of raw and heated samples were determined using pepsin-pancreatin hydrolysis, considering soluble nitrogen *via* Kjeldahl (ppKJ) and hydrolysed peptide linkages using trinitrobenzenesulfonic acid and o-phthaldialdehyde. *In vitro* digestibility was also determined using trypsin, chymotrypsin and peptidase (3-Enz) or trypsin, chymotrypsin, peptidase and pronase solution (4-Enz). None of the correlations between *in vitro* and *in vivo* digestibilities were significant (at $p < 0.0500$), but, strong correlations were observed between PDCAAS calculated by *in vitro* and *in vivo* results. PDCAAS-ppKJ, PDCAAS-3-Enz and PDCAAS-4-Enz presented the highest correlations with *in vivo* method, $r = 0.9316, 0.9442$ and 0.9649 ($p < 0.0500$), respectively. The use of *in vitro* methods for calculating PDCAAS may be promising and deserves more discussions.

© 2016 Elsevier Ltd. All rights reserved.

1. Introduction

Dietary protein quality is measured using a variety of methods, many of which are related to animal responses (often rats) after feeding them the test protein. These methods have the advantage of reflecting both the supply of essential amino acids and the digestibility of the protein, which are two important factors in protein evaluation. For dietary proteins to meet indispensable amino acid demands, their potential digestibility must be considered (Boye, Wijesinha-Bettoni, & Burlingame, 2012).

Although growth tests using rats have been used for many years (PER—Protein Efficiency Ratio, NPR—Net Protein Ratio, or NPU—Net Protein Utilization), these tests suffer from the disadvantage that rats may exhibit different amino acid requirements than humans. In this sense, nitrogen balance assays in rats, such as digestibility evaluation, may provide results that are the more directly relevant to humans (Schaafsma, 2012).

In 1989, the FAO/WHO Expert Consultation on Protein Quality Evaluation recommended the use of the Protein Digestibility Corrected

Amino Acid Score (PDCAAS) method, which considers both the indispensable amino acid content of the test protein and its digestibility. In this framework, amino acid score = content of the first limiting amino acid in the test protein/corresponding content of this amino acid in the WHO/FAO/UNU reference pattern (preschool child requirements, mg/g), and PDCAAS = protein digestibility × amino acid score. This is an extremely important measure, despite recent criticism and constant revision (Schaafsma, 2012; Boye et al., 2012; Leser, 2013; WHO/FAO/UNU, 2007).

Ileal digestibility is most appropriate for quantifying digestion, because true fecal nitrogen digestibility does not account for essential amino acids that are lost into the colon *via* intestinal flora activity (Schaafsma, 2012). However, rat assays for true digestibility have instead been extensively used, in part because of their practicality compared to the use of ileal-fistulated pigs. On the other hand, *in vivo* experiments suffer from high costs and long experimental run times (approximately 9 days at minimum) (McDonough et al., 1990), as well as ethical objections to animal use (Schaafsma, 2012). Thus, the employment of *in vitro* alternative methodologies must be considered and further developed to improve correlations with *in vivo* assays.

Although some studies have already PDCAAS data calculated with the use of *in vitro* digestibility values, a large number of *in vitro* methods

* Corresponding Author.

E-mail address: tavanool@yahoo.com.br (O.L. Tavano).

can be found in the literature. The purpose of this study was to compare protein digestibility measurements for chickpea proteins using a variety of *in vitro* methods and to use these methodologies to compute PDCAAS while evaluating them as potential substitutes for *in vivo* procedures, contributing to the choice of a methodology that, once adopted by different authors, facilitate comparison of data and well express what would be obtained by *in vivo* assays.

Chickpeas protein fractions were chosen as the basis for discussions on the comparison between *in vitro* and *in vivo* methodologies because these samples had previously been studied in our laboratory and their nutritional values have been well characterized using different *in vivo* and antinutritional assays (Tavano, Silva Junior, Demonte, & Neves, 2008, 2005). This gives a clearer picture and context for the results. The *in vitro* protein digestibility methods employed in this study are based on the measurement of potential susceptibility of protein peptide bonds to proteolysis by digestive enzymes, which is one of the first characteristics of proteins that interferes with the real process of *in vivo* protein digestion, and therefore in their nutritional value. These assays used enzyme mixtures or different enzymes sequences for proteolysis of protein sample, which try to mimic the effects of the human digestion process. We highlight the methods of Akeson and Stahman (1964) and Hsu, Vavak, Satterlee, and Miller (1977). The first is an ordered enzyme method, where pepsin is the first enzyme applied followed by pancreatin action, which hydrolysis effectiveness calculations were based on the following principles: the quantification of soluble nitrogen in the presence of a precipitation agent compared to the total nitrogen of the sample (e.g., the ppKj), because increased hydrolysis leads to greater solubility of protein fragments; or quantification of cleaved peptide bonds calculated by determining the free amino groups released through the reaction of compounds that react with amino groups using TNBS (trinitrobenzenesulfonic acid) or OPA (o-phthalaldehyde) reactions. The second consists of a multienzyme mix system (trypsin, chymotrypsin, and peptidase) with determination of protein sample pH before and after 10 min of reaction, which monitors the H⁺ ions released by broken peptide bonds and free carboxyl group “generation” (pH-drop assays). This second method is quick and simple to apply. In addition, another titrimetric method was used, including an additional step of hydrolysis by using pronase. In this way, besides the issues related to the procedure itself (which mainly involve enzyme choice), the adopted principle in each hydrolysis measurement method can influence the results of *in vitro* protein digestion assays and should be carefully considered as discussed in this paper.

2. Materials and methods

2.1. Materials

Chickpea seeds (*Cicer arietinum* L.) cv. IAC-Marrocos, were cultivated by and obtained from the Instituto Agronômico de Campinas, São Paulo, Brazil. Chickpea defatted flour and its protein fractions were obtained as previously described (Tavano et al., 2008), using differential protein solubility in different solvents (defatted flour, albumins, total and major globulins, and glutelins presenting 25.01, 75, 91.2, 97.3, and 68.6 g/100 g of protein, respectively). Pepsin (from porcine gastric mucosa), trypsin (porcine, Type IX), chymotrypsin (Type II), peptidase (from porcine intestinal mucosa), pancreatin, pronase (bacterial protease from *Streptomyces griseus*, Type XIV), 2,4,6-trinitrobenzene sulfonic acid (TNBS), o-phthalaldehyde (OPA) and benzoyl-DL-arginine-p-nitroanilide (BapNa) were purchased from Sigma Chemical Co., St. Louis, MO. Others chemicals were reagent grade.

2.2. Heat treatment

A portion of defatted flour and protein fractions were suspended in distilled water (1:6 w/v), autoclaved at 121 °C for 15 min, and immediately cooled and lyophilized again, as described by Tavano et al. (2008).

2.3. Nitrogen determination

Nitrogen was determined using the Kjeldahl method (AOAC, 1990). Crude protein was calculated as N × 6.25 for chickpea proteins and N × 6.38 for casein samples.

2.4. Pepsin/pancreatin method

In vitro protein digestibility of 50 mg protein from raw and heated samples were determined as described by Akeson and Stahman (1964) using a pepsin-pancreatin (pp) incubation sequence (0.75 mg pepsin, 1.641 units, and 2.0 mg pancreatin), at 37 °C, for 3 h and 24 h, respectively. The enzymatic reaction was then interrupted by adding 10% trichloroacetic acid, followed by centrifugation at 7000g for 15 min. The resulting supernatant was used for determining the degree of hydrolysis using the following methods:

2.4.1. ppKj

Determination of soluble nitrogen by Kjeldahl (AOAC, 1990). Percent hydrolysis was calculated as the percent of soluble nitrogen compared to the total nitrogen in the sample and using the following ratio:

% Hydrolysis = $(N_a - N_{ba} - N_{be}) / N_t \times 100$, where: N_a = soluble nitrogen in the supernatant of the hydrolysed sample; N_{ba} = soluble nitrogen in the supernatant of the non-hydrolysed sample (a blank sample without enzymes); N_{be} = soluble nitrogen in the supernatant of solution containing only enzyme (a blank sample without samples); N_t = total nitrogen in the sample.

2.4.2. Determination of the percent of peptide linkages hydrolysed, as determined by the number of free amino groups in the supernatant. This was calculated using the following methods:

2.4.2.1. ppTNBS. Free amino groups in the supernatant were determined using trinitrobenzenesulfonic acid (TNBS), as described by Fields (1972) and modified by Spadaro, Draghetto, Del Lama, Camargo, and Greene (1979). The mixture contained the following: 0 to 0.4 mL of sample, 0.4 mL borate/KOH buffer, pH 9.5 and 0.2 mL of TNBS solution (5 mM). After 50 min at 25 °C, the reaction was interrupted by addition of 0.2 mL of 18 mM sodium sulfite in 2 M sodium phosphate buffer, and the absorbance was measured at 420 nm using the reagent solution as a blank.

2.4.2.2. ppOPA. Free amino groups in the supernatant were measured using o-phthalaldehyde (OPA), as described by Church, Swaisgood, Porter, and Catignani (1983). Aliquots of samples between 0 and 130 µL were combined with 1 mL of OPA reagent that was freshly prepared every day (25 mL of 100 nM sodium tetraborate, 2.5 mL SDS 20%, 40 mg OPA/mL methanol, 100 µL β-mercaptoethanol and adjusted to 50 mL with distilled water). After 2 min of reaction the absorbance was measured at 340 nm using the reagent solution as a blank.

Analytical reference curves for both ppTNBS and ppOPA were constructed using L-leucine as a standard. The total moles of amino acids present in the initial mass of protein were estimated using the average molecular weight of amino acids (MW = 113) and also *via* corrected values with respect to the individual real concentration values in the sample and molecular weights of each amino acid considering the previously determined sample aminograms (Tavano et al., 2008). The equation used to calculate percent hydrolysis (%H) was:

%H = $(AAs - AAba - Aabe) \times 100 / Aatm$ or $Aatc$, where: AAs = amino acid mols in the sample supernatant; AAba = amino acid mols in the blank supernatant of the samples; Aabe = amino acid mols in the blank supernatant of the enzymes; Aatm = total amino acid mols in the sample considering the mean molecular weight of the amino acids (ppOPA and ppTNBS); Aatc = total amino acid mols in the sample considering the individual molecular weight of each amino acid (ppOPAc and ppTNBSc).

2.5. 3-Enzyme method (3-Enz)

In vitro protein digestibility (IVPD) was applied to unheated and heated chickpea protein samples using the multienzyme method as described by Hsu et al. (1977). Protein was dispersed in 10 mL distilled water (6.25 mg mL⁻¹) and adjusted to pH 8.0 at 37 °C in a water bath followed by addition of 1.0 mL multienzyme solution (1.6 mg trypsin, 24,320 units, 3.1 mg chymotrypsin, 155 units and 1.3 mg of peptidase from porcine intestinal mucosa, 0.133 units mL⁻¹). The pH variation was then recorded after 10 min reaction. The equation $Y = 210.46 - 18.10 X$, was used for calculating *in vitro* digestibility, where X is pH change after 10 min (Hsu et al., 1977).

2.6. 4-Enzyme method (4-Enz)

The AOAC-982.30 (Association of Official Analytical Chemists, 2000) method was then applied to the unheated and heated chickpea protein samples. Protein was dispersed in water (6.25 mg nitrogen or 62.5 mg protein in 10 mL distilled water) and adjusted to pH 8.0 at 37 °C in a water bath. Multienzyme solution A (22,704 units of trypsin, 182 units chymotrypsin and 0.052 units of peptidase from porcine intestinal mucosa, in 1.0 mL) was added. After exactly 10 min at 37 °C, 1.0 mL of enzyme solution B (6.5 units of pronase in 1.0 mL distilled water) was added, and the sample was incubated for an additional 20 min at 55 °C. The equation $Y = 234.84 - 22.56 X$, was used for calculating *in vitro* digestibility, where X is the pH value after the final 20 min.

2.7. Protein Digestibility Corrected Amino Acid Scoring

Protein Digestibility Corrected Amino Acid Scoring (PDCAAS) was calculated on the basis of the WHO/FAO/UNU essential amino acid scoring pattern for 1- to 2-year-old children (WHO/FAO/UNU, 2007), as described by Sarwar and McDonough (1990), using the protein digestibility values obtained *in vitro* or *in vivo* methods for correction. PDCAAS = (limiting amino acid content in sample protein/same amino acid content of the reference pattern) × protein digestibility.

When *in vivo* true protein digestibility was used in calculations, the previously obtained results (Tavano et al., 2008) were considered.

2.8. Statistical analyses

All assays were performed in triplicate and are expressed as means ± SD. Analysis of variance (ANOVA) was used to compare results ($p \leq 0.05$) using Student's *t*-test. Statistical analysis was conducted using Statsoft STATISTICA 8.0 (Statsoft, 2007).

3. Results and discussion

Results of pepsin-pancreatin hydrolysis assays varied widely between the different methods of determining the degree of hydrolysis (ranging from 34.73 to 64.49% for unheated flour, for instance), as shown in Table 1. The ppKj method (pepsin-pancreatin hydrolysis followed by Kjeldahl determination of soluble nitrogen) was not able to detect the improvement in digestibility of heated samples, while most other *in vitro* methods could detect such differences. Although ppTNBS (pepsin-pancreatin hydrolysis followed by TNBS amino group determination), ppOPA (pepsin-pancreatin hydrolysis followed by OPA amino group determination) and ppKj are based on the same sample hydrolysate (pepsin-pancreatin hydrolysates stopped *via* TCA precipitation), different interpretations of the products generated may arise because ppKj is not capable of differentiating whether nitrogen is from peptides or free amino acids. In other words, ppKj can confuse one tetrapeptide for four free amino acids. This does not occur using ppOPA and ppTNBS, e.g., a tetrapeptide released by hydrolysis that remains in solution will be recognized as a single α -amino group released (one broken link), while using the ppKjeldahl method this same fragment is interpreted as four nitrogens in solution, even if only one peptide bond has been broken. This explains the high hydrolysis values for most samples using this method compared to data generated by ppOPA and ppTNBS (Table 1).

While this observation may suggest that ppKj overestimates digestibility, it is important to consider that for *in vivo* assays (true digestibility *via* fecal rat assays), protein digestibility is evaluated considering the

Table 1
In vitro and *in vivo* chickpea protein digestibility of chickpea flour and fractions using different methodologies.

Samples ^a	<i>In vitro</i> methods ^b (% hydrolysis degree)							<i>In vivo</i> ^c (TD)
	ppOPA	ppOPAc	ppTNBS	ppTNBSc	ppKj	3-Enz	4-Enz	
Casein	65.99 ^{aA} ± 2.90	88.41 ^B ± 3.88	47.04 ^a ± 1.17	63.02 ^A ± 1.57	87.09 ^B ± 4.09	86.05 ^{aB} ± 0.55	107.75 ± 1.84	96.08 ^a ± 1.61
Flour (U)	42.20 ^{bA} ± 1.19	48.91 ± 1.38	34.73 ^b ± 0.99	40.25 ^{aA} ± 1.15	64.49 ^a ± 3.42	70.99 ^B ± 0.27	80.53 ^C ± 0.60	78.42 ^{bBC} ± 7.88
Flour (H)	53.00 ^{dEFA} ± 4.76	61.42 ^{acdAB} ± 5.52	38.05 ^{bc} ± 2.13	44.10 ^{ab} ± 2.47	63.59 ^{aB} ± 3.17	80.26 ^{aC} ± 0.58	82.03 ^{aD} ± 0.34	78.75 ^{bCD} ± 3.42
Albumin (U)	40.10 ^{cA} ± 0.88	45.64 ^b ± 1.01	27.64 ± 0.43	31.46 ± 0.49	57.28 ^{bc} ± 1.25	68.19 ± 0.36	78.08 ± 0.85	88.76 ^C ± 4.57
Albumin (H)	51.01 ^{dA} ± 0.47	58.06 ^{aB} ± 0.53	39.36 ^C ± 0.65	44.80 ^b ± 0.74	54.38 ^{cAB} ± 3.70	76.37 ± 0.29	81.96 ^a ± 0.34	93.00 ^a ± 5.04
Total globulin (U)	47.51 ^c ± 1.33	55.66 ^C ± 1.56	31.17 ± 0.22	36.52 ± 0.25	75.22 ^{dA} ± 3.26	80.68 ^{aA} ± 2.26	93.84 ± 0.60	87.05 ^C ± 2.46
Total globulin (H)	68.49 ^a ± 2.04	80.23 ^A ± 2.39	47.36 ^a ± 1.33	55.48 ^C ± 1.56	75.26 ^{dA} ± 3.95	88.65 ^b ± 1.01	96.40 ^B ± 0.47	94.61 ^{aB} ± 2.17
Major globulin (H)	61.53 ^{fA} ± 3.91	71.86 ^{dB} ± 4.57	46.81 ^a ± 2.43	54.67 ^{cA} ± 2.84	70.68 ^B ± 0.80	89.31 ^{bc} ± 0.84	89.63 ^{bC} ± 0.13	94.65 ^a ± 2.27
Glutelin (H)	36.56 ^{cA} ± 3.84	42.26 ^{bA} ± 4.43	37.43 ^{bcA} ± 4.53	43.27 ^{abA} ± 5.23	59.45 ^{ab} ± 3.40	83.52 ± 0.21	88.72 ± 1.07	78.19 ^b ± 2.01

Values were expressed as means ± standard deviation (*in vitro* assays n = 3, *in vivo* assays n = 8). Means followed by the same letter (small in the same column and capital in the same line) indicate no significant difference between samples ($p < 0.0500$).

^a U = Unheated sample, H = heated sample. The heated samples were autoclaving at 121 °C/15 min.

^b ppOPA, ppTNBS, ppKj, ppOPAc and ppTNBSc: pepsin/pancreatin hydrolysis method (Akeson & Stahman, 1964) using o-phthalaldehyde α -amino group detection (OPA), trinitrobenzenesulfonic acid α -amino group detection (TNBS) or soluble nitrogen by Kjeldahl (Kj) for hydrolysis degree calculation (OPAc and TNBSc considerate total amino acid μ Mols using each sample aminograms); 3-Enz = Hsu et al. (1977) method using 10 min hydrolysis with trypsin, chymotrypsin and peptidase solution; 4-Enz = AOAC-982.30 (AOAC, 2000) method using 10 min hydrolysis with trypsin, chymotrypsin and peptidase solution, plus additional 10 min using pronase, as described in Materials and methods section.

^c TD = rats fecal true digestibility values, as previously reported in Tavano et al. (2008).

Table 2
Correlation coefficients between chickpea protein digestibility values obtained using different methods.

Correlated methods	Correlation coefficients (r) ^a
Pepsin-pancreatin method	
ppOPA/ppTNBS	0.8172 (<i>p</i> = 0.0132)*
ppOPAc/ppTNBS	0.8289 (<i>p</i> = 0.0109)*
ppOPA/ppKj	0.6066 (<i>p</i> = 0.1108)
ppOPAc/ppKj	0.6380 (<i>p</i> = 0.0887)
ppTNBS/ppKj	0.3805 (<i>p</i> = 0.3524)
ppTNBS/ppKj	0.4260 (<i>p</i> = 0.2926)
3-Enzymes/pepsin-pancreatin method	
3-Enz/ppOPA	0.6920 (<i>p</i> = 0.0572)
3-Enz/ppOPAc	0.7086 (<i>p</i> = 0.0491)*
3-Enz/ppTNBS	0.8233 (<i>p</i> = 0.0120)*
3-Enz/ppTNBS	0.8409 (<i>p</i> = 0.0089)*
3-Enz/ppKj	0.6159 (<i>p</i> = 0.1039)
4-Enzymes/pepsin-pancreatin method	
4-Enz/ppOPA	0.5544 (<i>p</i> = 0.1538)
4-Enz/ppOPAc	0.5804 (<i>p</i> = 0.1314)
4-Enz/ppTNBS	0.5240 (<i>p</i> = 0.1824)
4-Enz/ppTNBS	0.5558 (<i>p</i> = 0.1525)
4-Enz/ppKj	0.8132 (<i>p</i> = 0.0140)*
4-Enzymes/3-enzymes method	
3-Enz/4-Enz	0.8275 (<i>p</i> = 0.0112)*
<i>In vivo/in vitro</i>	
TD/ppOPA	0.6785 (<i>p</i> = 0.0640)
TD/ppOPAc	0.6637 (<i>p</i> = 0.0727)
TD/ppTNBS	0.4553 (<i>p</i> = 0.2570)
TD/ppTNBS	0.4481 (<i>p</i> = 0.2654)
TD/ppKj	0.2683 (<i>p</i> = 0.5205)
TD/3-Enz	0.3534 (<i>p</i> = 0.3904)
TD/4-Enz	0.3675 (<i>p</i> = 0.3704)

^a Only chickpea values were considered.

* Marked correlations (*r*) are significant at *p* < 0.0500.

capacity of the protein to be absorbed in the intestinal tract, which is determined through quantification of the ingested nitrogen that is absorbed by the organism. This process is influenced by the hydrolysis susceptibility of the polypeptide chain by digestive enzymes as much

as the availability to absorb the generated products. However, this absorption process can occur in the free amino acid form or even in the form of peptides containing few amino acid residues, such as di-, tri- or tetra peptides, or even more (Pappenheimer, Dahl, Karnovsky, & Maggio, 1994; Jahan-Mihan, Luhovyy, El Khoury, & Anderson, 2011). As such, it is important remember that a small number of disrupted peptide bonds should not necessarily be interpreted as a low percentage of absorption or low digestibility *in vivo*.

On the other hand, it is necessary to observe that *in vitro* methods that employ soluble nitrogen determination can present additional failures due to limitations in the precipitation process. Yvon, Chabanet, and Pélissier (1989) demonstrated that peptide fragments containing up to fifty residues can stay in solution until 12% trichloroacetic acid. We could interpret this to mean that any large peptides could be potentially absorbed in the intestine, but it is not clear that this occurs. Although there is evidence to suggest absorption of large peptides such as octapeptides (Pappenheimer et al., 1994), the relationship between the sizes of the soluble proteins in precipitant solutions and peptides that actually can be absorbed merits further study.

Hydrolysis methods that use ppOPA and ppTNBS have another usual inconvenience relative to the final hydrolysis calculation. The 100% hydrolysis value (total amino acids that would be liberated if all peptide bonds could be cleaved), which is necessary for calculations, may be estimated using the average molecular weight (PM = 113) to calculate the number of total amino acids in the sample, as employed herein (considering the total number of amino acids in 50 mg of test protein), as a simplified protocol for calculation. But, the amino acid profiles of different samples can be very distinct, and considering the large variation between individual molecular weights of the amino acids, the average molecular weight for each protein sample can vary widely. Some authors suggest performing total hydrolysis of a sample (*via* HCl 6 N hydrolysis) for determining the 100% equivalent but this step adds more work to the procedure. The ppKjeldahl method does not suffer from this problem, because the initial values of total nitrogen in the samples (100%) are obtained by the same Kjeldahl methodology used to determine the soluble nitrogen. To verify the influence of the average molecular weight procedure in the final hydrolysis calculations, we compared it with the use of the individual aminogram of each sample as a

Table 3
In vitro and *in vivo* chickpea protein digestibility expressed as % of casein results (100%).

Samples ^a	<i>In vitro</i> methods ^b (%)							<i>In vivo</i> ^c % TD
	ppOPA	OPAc	ppTNBS	ppTNBS	ppKj	3-Enz	4-Enz	
Casein	100	100	100	100	100	100	100	100
Flour (U)	63.95 ^{aeA}	55.32	73.83 ^{ab}	63.87 ^{aA}	74.05 ^{abC}	82.50 ^D	69.30	83.09 ^{aCD}
	± 1.80	± 1.56	± 2.12	± 1.83	± 3.93	± 0.31	± 0.52	± 8.35
Flour (H)	80.31 ^{bcdA}	69.48 ^{acdB}	80.89 ^{abA}	69.98 ^{bb}	73.02 ^{aAB}	93.27 ^a	70.59 ^{ab}	83.44 ^{aA}
	± 7.22	± 6.25	± 4.53	± 3.92	± 3.64	± 0.67	± 0.29	± 3.63
Albumin (U)	60.77 ^{aA}	51.62 ^{bb}	58.76 ^A	49.93 ^B	65.77 ^{bc}	79.25	67.19 ^C	94.04 ^b
	± 1.34	± 1.14	± 0.91	± 0.78	± 1.43	± 0.42	± 0.74	± 4.84
Albumin (H)	77.31 ^b	65.67 ^{aA}	83.66 ^b	71.08 ^{bb}	62.44 ^{ba}	88.75	70.52 ^{ab}	98.54 ^{bc}
	± 0.71	± 0.60	± 1.38	± 1.17	± 4.25	± 0.33	± 0.29	± 5.34
Total globulin (U)	72.98 ^c	62.96 ^c	66.27	57.95	86.37 ^c	93.76 ^{aA}	80.75	92.23 ^{ba}
	± 2.02	± 1.77	± 0.46	± 0.40	± 3.75	± 2.63	± 0.52	± 2.61
Total globulin (H)	103.79 ^A	90.75 ^B	100.67 ^{cA}	88.03 ^{cb}	86.41 ^{cb}	103.02 ^{ba}	82.95	100.25 ^{ca}
	± 3.10	± 2.71	± 2.83	± 2.48	± 4.54	± 1.17	± 0.40	± 2.30
Major globulin (H)	93.25 ^{dA}	81.29 ^{dBC}	99.50 ^{cAD}	86.74 ^c	81.69 ^B	103.79 ^{bd}	77.12 ^{bc}	100.29 ^{cd}
	± 5.93	± 5.17	± 5.17	± 4.51	± 0.92	± 0.97	± 0.11	± 2.40
Glutelin (H)	55.40 ^{aA}	47.80 ^{ba}	79.56 ^{abBCD}	68.66 ^{ab}	68.26 ^{abB}	97.06	76.35 ^{bc}	82.85 ^{ad}
	± 5.81	± 5.02	± 9.62	± 8.30	± 3.90	± 0.24	± 0.92	± 2.13

Values were expressed as means ± standard deviation (*in vitro* assays *n* = 3, *in vivo* assays *n* = 8). Means followed by the same letter (small in the same column and capital in the same line) indicate no significant difference between samples (*p* < 0.0500).

^a U = Unheated sample, H = heated sample. The heated samples were autoclaving at 121 °C/15 min.

^b ppOPA, ppTNBS, ppKj, ppOPAc and ppTNBS: pepsin/pancreatin hydrolysis method (Akeson & Stahman, 1964) using o-phthalaldehyde α-amino group detection (OPA), trinitrobenzenesulfonic acid α-amino group detection (TNBS) or soluble nitrogen by Kjeldahl (Kj) for hydrolysis degree calculation (OPAc and TNBS considerate total amino acid uMols using each sample aminograms); 3-Enz = Hsu et al. (1977) method using 10 min hydrolysis with trypsin, chymotrypsin and peptidase solution; 4-Enz = AOAC-982.30 (AOAC, 2000) method using 10 min hydrolysis with trypsin, chymotrypsin and peptidase solution, plus additional 10 min using pronase, as described in Materials and methods section.

^c TD = rats fecal true digestibility values, as previously reported in Tavano et al. (2008).

Table 4
Indispensable amino acid compositions^a and scores^b of chickpea proteins.

Amino acids	Pattern ^c	Flour		Albumin		Total globulin		Major globulin		Glutelin		Casein	
		AA ^a	Score ^b	AA ^a	Score ^b	AA ^a	Score ^b	AA ^a	Score ^b	AA ^a	Score ^b	AA ^a	Score ^b
Thr	27	39.3	1.00	52.1	1.00	29.5	1.00	28.8	1.00	43.5	1.00	38.0	1.00
Val	42	40.6	0.97	45.5	1.00	29.5	0.70	41.5	0.98	50.5	1.00	55.0	1.00
Met + Cys	26	19.5	0.75	47.5	1.00	9.9	0.38	8.7	0.33	18.7	0.72	35.0	1.00
Ile	31	38.3	1.00	42.7	1.00	40.5	1.00	40.6	1.00	45.2	1.00	44.0	1.00
Leu	63	77.3	1.00	71.8	1.00	78.1	1.00	76.0	1.00	76.8	1.00	76.0	1.00
Tyr + Phe	46	90.0	1.00	77.1	1.00	96.4	1.00	93.2	1.00	91.8	1.00	70.0	1.00
Trp	7.4	6.2	0.84	7.0	0.94	7.8	1.00	8.0	1.00	10.9	1.00	10.0	1.00
Lys	52	73.1	1.00	93.3	1.00	66.1	1.00	65.2	1.00	78.9	1.00	68.0	1.00
His	18	28.6	1.00	30.2	1.00	30.8	1.00	29.7	1.00	29.9	1.00	25.0	1.00

^a AA = Amino acid composition (mg/100 mg protein) as previously reported in Tavano et al. (2008).

^b Amino Acid Score = amino acid content in test protein/amino acid content of reference pattern. Values underlined represent the first limiting amino acid of the sample.

^c Amino acid scoring patterns for 1–2 years children according to 2007 WHO/FAO/UNU report (2007).

reference to calculate the individual value of 100% of amino acids in the samples. These corrections are not common in the *in vitro* protein digestibility literature, because the necessity of determining the total amino acid makes it difficult to execute and runs counter to one of the justifications for replacement of *in vivo* by *in vitro* methods, which is the simplicity. The total μmol s of amino acids/50 mg proteins was calculated using the individual molecular weight of each amino acid and their individual concentration observed in each sample (Tavano et al., 2008). The following values were achieved: 381.70, 388.68, 377.64, 378.81, 382.74 and 330.21 μmol s of amino acids/50 mg proteins for flour, albumin, total globulin, major globulin, glutelin and casein, respectively. It is possible to observe the differences between the corrected (ppOPAc and ppTNBSc) and non-corrected (ppOPA and ppTNBS) values (Table 1), which can expose the variation enclosed in the standardization of the initial amino acid concentration (which assumes 50 mg protein = 442 μmol s amino acids for all samples).

The adoption of the individual amino acid concentration in each sample produced a relative increase in the percentage of hydrolysis values (ppTNBSc and ppOPAc in comparison of ppTNBS and ppOPA), as shown in Table 1 because all results for the total μmol s of amino

acids were lower than those values calculated by the average molecular weight of the amino acids. However, when these percentages are expressed in relation to casein (%) the proportions are reversed (Table 3), and there is a reduction in ppTNBSc and ppOPAc digestibility values compared to ppTNBS and ppOPA. This is because the values by the total number of amino acids of casein suffer the greatest reduction when the aminogram is used (Tables 1 and 3).

This correction in the calculations by aminograms, although made to obtain the most accurate results and therefore increase similarity to *in vivo* condition, caused no major changes in the correlations of these methods with the results generated by testing with animals (Table 2). In fact, none of the correlations established between the digestibility *in vitro* and *in vivo* were significant (at $p < 0.0500$). The only significant correlations found were between some *in vitro* methods and most of these were moderate correlations (Table 2).

Panasiuk, Amarowicz, Kostyra, and Sijtsma (1998) found a correlation coefficient of 0.964 for the TNBS and OPA methods for raw pea protein samples after hydrolysis. In this study, the correlation between ppOPA and ppTNBS was 0.8172 ($p = 0.0133$), and the correlation between ppOPAc and ppTNBSc was 0.8289 ($p = 0.0109$), as shown in

Table 5
Protein digestibility-corrected amino acid score (PDCAAS) of chickpea proteins and casein calculated using different *in vitro* and *in vivo* protein digestibility assays.

Samples ^a	PDCAAS							AAS \times TD ^c
	First limiting amino acid score (AAS) \times <i>in vitro</i> digestibilities ^b							
	ppOPA	ppOPAc	ppTNBS	ppTNBSc	ppKj	3-Enz	4-Enz	
Casein	65.99 ^A	88.41 ^{aBC}	47.04	63.02 ^A	87.09 ^{aBC}	86.05 ^B	107.75	91.59 ^{aC}
Flour (U)	± 2.90	± 3.88	± 1.17	± 1.57	± 4.09	± 0.55	± 1.84	± 1.67
	31.65 ^A	36.68	26.05 ^a	30.19 ^{abA}	48.37 ^{ab}	53.24	60.40 ^B	58.82 ^{bB}
Flour (H)	± 0.89	± 1.03	± 0.74	± 0.86	± 2.56	± 0.21	± 0.45	± 5.91
	39.75 ^{aA}	46.07 ^{aAB}	28.54 ^a	33.08 ^a	47.69 ^{abB}	60.20 ^{aC}	61.52 ^C	59.06 ^{bC}
Albumin (U)	± 3.57	± 4.14	± 1.60	± 1.85	± 2.38	± 0.43	± 0.25	± 2.56
	37.69 ^a	42.90 ^a	25.98 ^a	29.57 ^b	53.84 ^c	64.10	73.40	83.43 ^c
Albumin (H)	± 0.83	± 0.95	± 0.40	± 0.46	± 1.17	± 0.34	± 0.80	± 4.29
	47.95 ^A	54.58 ^B	37.00	42.11	51.12 ^{bcAB}	71.79	77.04	87.42 ^{aC}
Total globulin (U)	± 0.44	± 0.50	± 0.61	± 0.70	± 3.48	± 0.27	± 0.32	± 4.74
	18.05	21.15 ^b	11.84	13.88	28.58 ^{dA}	30.66 ^{bA}	35.66	33.08 ^d
Total globulin (H)	± 0.50	± 0.59	± 0.09	± 0.02	± 1.24	± 0.84	± 0.23	± 0.93
	26.03 ^{bA}	30.49 ^{cb}	18.00	21.08	28.60 ^{dAB}	33.69	36.63 ^C	35.95 ^C
Major globulin (H)	± 0.77	± 0.91	± 0.50	± 0.59	± 1.50	± 0.38	± 0.18	± 0.82
	20.30 ^A	23.71 ^{bb}	15.44	18.04 ^A	23.32 ^B	29.47 ^{bcD}	29.58 ^C	31.23 ^{dd}
Glutelin (H)	± 1.29	± 1.51	± 0.80	± 0.94	± 0.26	± 0.28	± 0.05	± 0.75
	26.32 ^{bA}	30.43 ^{cA}	26.95 ^{aA}	31.15 ^{abA}	42.80 ^a	60.13 ^a	63.88	56.30 ^b
	± 2.76	± 3.19	± 3.26	± 3.76	± 2.45	± 0.15	± 0.77	± 1.45

Values were expressed as means \pm standard deviation (*in vitro* assays $n = 3$, *in vivo* assays $n = 8$). Means followed by the same letter (small in the same column and capital in the same line) indicate no significant difference between samples ($p < 0.0500$).

^a U = Unheated sample, H = heated sample. The heated samples were autoclaving at 121 °C/15 min.

^b ppOPA, ppTNBS, ppKj, ppOPAc and ppTNBSc: pepsin/pancreatin hydrolysis method (Akeson & Stahman, 1964) using o-phthalaldehyde α -amino group detection (OPA), trinitrobenzenesulfonic acid α -amino group detection (TNBS) or soluble nitrogen by Kjeldahl (Kj) for hydrolysis degree calculation (OPAc and TNBSc considerate total amino acid μmol s using each sample aminograms); 3-Enz = Hsu et al. (1977) method using 10 min hydrolysis with trypsin, chymotrypsin and peptidase solution; 4-Enz = AOAC-982.30 (AOAC, 2000) method using 10 min hydrolysis with trypsin, chymotrypsin and peptidase solution, plus additional 10 min using pronase, as described in Materials and methods section.

^c TD = rats fecal true digestibility values, as previously reported in Tavano et al. (2008).

Table 6

Correlation coefficients between *in vitro/in vivo* chickpea protein digestibility corrected amino acid score values.

Correlated methods	Correlation coefficients (r) ^a
TD/ppOPA	0.9007 ($p = 0.0279$) [*]
TD/ppOPAc	0.8916 ($p = 0.0319$) [*]
TD/ppTNBS	0.8852 ($p = 0.0009$) [*]
TD/ppTNBSc	0.8755 ($p = 0.0013$) [*]
TD/ppKj	0.9316 ($p = 0.0247$) [*]
TD/3-Enz	0.9442 ($p = 0.0135$) [*]
TD/4-Enz	0.9649 ($p = 0.0104$) [*]

^a Only chickpea values were considered.

^{*} Marked correlations (r) are significant at $p < 0.0500$.

Table 2. The ppOPA method showed the greatest ability to differentiate between raw and heated samples, followed by ppTNBS, 3-Enz and 4-Enz (Table 1).

Although the 3-Enz and 4-Enz methods were able to distinguish the crude from heated samples, the percentage increase of digestibility after heat treatment expressed by these two methods is, in most cases, much lower than that demonstrated by pepsin-pancreatin. For albumin samples, for example, ppOPA and ppTNBS showed increases of approximately 27% and 42%, respectively, whereas 3-Enz and 4-Enz found increases of approximately 12 and 5% (Table 1). The same was found for total globulin sample, where ppOPA and ppTNBS showed increases of approximately 44% and 52% and 3-Enz and 4-Enz showed increases of approximately 10 and 3%, respectively.

These two last methods, besides using a different hydrolysis degree detection principle, also use a much larger amount of enzymes, and their ratio is approximately 1:10 enzyme:substrate, while for pepsin-pancreatin this ratio is between 1:50 (pepsin) and 1:25 (pancreatin). That may help explain the higher percentage of hydrolysis observed for these methods. The 4-Enz method in particular makes use of an additional hydrolysis step with pronase. Pronase is a commercial preparation containing several proteases from *Streptomyces griseus* (Trop & Birk, 1970), usually applied when a high degree of hydrolysis is desired.

Values for the 4-Enz method are in fact among the highest results for digestibility, including the extrapolation of casein dates (presenting 107.75%), suggesting a possible overestimation when this method is used. Another important observation regarding these two methods is that the entire hydrolysate is used for computing the amount of hydrolysis, while in the pepsin-pancreatin method only the supernatant after TCA precipitation is used.

According to Wolzak, Bressani, and Brenes (1981), different enzymatic and chemical methods were not equally sensitive to certain physical and chemical characteristics of food samples, such as presence of anti-nutritional and cell wall structures, which contributes to the difficulty of establishing a single protocol that is applicable to different samples and showing a close relationship with *in vivo* phenomena. In the physiological process of digestion, other processes and enzymes could contribute indirectly to protein digestion, as they help with the destruction of food matrix and thus allow improved access of proteases. These variables are generally not computed *in vitro*, as they usually only consider isolated proteases.

With regard to antinutritional compounds, especially protease inhibitors, *in vitro* assays may react to their presence in different ways than seen in for *in vivo* digestion. Although most *in vitro* assays have pointed to an improvement of digestibility of flour, albumin and total globulin samples after heating, the TD data (Tavano et al., 2008) (true digestibility) indicated that this is not always detectable *in vivo* (Table 1). Using this method, flour and albumin sample digestibilities were not significantly increased after heating. It is interesting to note that these fractions have higher values of the trypsin inhibitor unit (TIU), according to previously observed data (38.72, 2.31, 239.20 and 5.47 TIU/mg protein, respectively for unheated and heated flour, and unheated and heated albumin) (Tavano et al., 2008). A significant improvement in their digestibilities would be expected, therefore, after heat treatment, but this was not observed for *in vivo* testing. At first sight, this indicates that these *in vivo* tests were not greatly influenced by the presence of these inhibitory components. This could be due to an adaptive effect, demonstrated by the changes observed in the pancreas of rats fed the same samples of unheated flour and albumin, as shown in previously published studies (Tavano et al., 2005). These data clearly

Table 7

Protein digestibility-corrected amino acid score (PDCAAS) of chickpea proteins calculated using different *in vitro* and *in vivo* protein digestibility assays and expressed as percentage of casein value (100%).

Samples ^a	PDCAAS (% of casein values) ^b							
	ppOPA	ppOPAc	ppTNBS	ppTNBSc	ppKj	3-Enz	4-Enz	TD ^c
Casein	100	100	100	100	100	100	100	100
Flour (U)	47.96 ^A	41.49	55.37 ^{aB}	47.90 ^{abA}	55.54 ^{acB}	61.87 ^C	56.05 ^B	61.21 ^{aBC}
	± 1.35	± 1.16	± 1.26	± 1.36	± 2.98	± 0.24	± 0.41	± 6.15
Flour (H)	60.24 ^{aABC}	52.10 ^{aABC}	60.67 ^{aA}	52.48 ^{ab}	54.76 ^{acAB}	69.95 ^a	57.10 ^A	61.47 ^{aC}
	± 5.40	± 4.98	± 3.40	± 2.93	± 2.73	± 0.50	± 0.23	± 2.67
Albumin (U)	57.12 ^{aA}	48.53 ^{ab}	55.23 ^{aA}	46.93 ^{bb}	61.82 ^b	74.49	68.12	86.84 ^b
	± 1.26	± 1.07	± 0.85	± 0.73	± 1.34	± 0.39	± 0.74	± 4.46
Albumin (H)	72.66 ^A	61.73 ^B	78.65	66.82	58.69 ^{bcB}	83.43	71.50 ^A	90.99 ^B
	± 0.67	± 0.56	± 1.29	± 1.11	± 4.00	± 0.31	± 0.30	± 4.93
Total globulin (U)	27.36	23.92 ^b	25.18	22.02	32.82 ^{dA}	35.63 ^b	33.09 ^A	34.43 ^C
	± 0.76	± 0.67	± 0.19	± 0.01	± 1.42	± 0.97	± 0.21	± 1.04
Total globulin (H)	39.44 ^{ba}	34.48 ^{bcB}	38.26 ^{AB}	33.45 ^{BC}	32.84 ^{dC}	39.15 ^A	34.00 ^{BC}	37.42 ^{AB}
	± 4.13	± 3.61	± 4.62	± 4.06	± 1.88	± 0.09	± 0.41	± 0.85
Major globulin (H)	30.77 ^{AB}	26.82 ^{baC}	32.83 ^B	28.63 ^{AC}	26.78 ^C	34.25 ^b	27.45 ^C	32.51 ^{cB}
	± 1.96	± 1.68	± 1.70	± 1.51	± 0.30	± 0.33	± 0.04	± 0.78
Glutelin (H)	39.89 ^{baB}	34.42 ^{cA}	57.29 ^{aCD}	49.44 ^{abBC}	49.15 ^{ac}	69.88 ^a	59.28 ^D	58.59 ^{aD}
	± 4.18	± 3.61	± 6.91	± 5.99	± 2.81	± 0.17	± 0.71	± 1.51

Values were expressed as means ± standard deviation (*in vitro* assays $n = 3$, *in vivo* assays $n = 8$). Means followed by the same letter (small in the same column and capital in the same line) indicate no significant difference between samples ($p < 0.0500$).

^a U = Unheated sample, H = heated sample. The heated samples were autoclaving at 121 °C/15 min.

^b ppOPA, ppTNBS, ppKj, ppOPAc and ppTNBSc: pepsin/pancreatin hydrolysis method (Akeson & Stahman, 1964) using o-phthalaldehyde α-amino group detection (OPA), trinitrobenzenesulfonic acid α-amino group detection (TNBS) or soluble nitrogen by Kjeldahl (Kj) for hydrolysis degree calculation (OPAc and TNBSc considerate total amino acid uMols using each sample aminograms); 3-Enz = Hsu et al. (1977) method using 10 min hydrolysis with trypsin, chymotrypsin and peptidase solution; 4-Enz = AOAC-982.30 (AOAC, 2000) method using 10 min hydrolysis with trypsin, chymotrypsin and peptidase solution, plus additional 10 min using pronase, as described in Materials and methods section.

^c TD = rats fecal true digestibility values, as previously reported in Tavano et al. (2008).

showed that animals suffered hypertrophy of this organ (42% of pancreatic increase in animals fed with unheated flour and 45.5% for rats fed with unheated albumin after 14 days, while the other groups showed no changes). This suggests that the results found for protein digestibility could reflect a physiological adaptation of these animals, which could generate a compensatory effect enzyme production to overcome differences between heated and unheated samples. In those *in vitro* assays that utilize high concentrations of enzymes (3-Enz and 4-Enz), especially trypsin and chymotrypsin, this type of compensating effect might also be occurring.

On the other hand, digestibility increases caused by heat treatment, as observed for total sample globulin (TIU low levels) (Tavano et al., 2008), may be more reflective of conformational changes than thermolabile antinutritional.

It is very difficult to isolate the main factor responsible for the effect of heating on proteins, as many effects are likely taking place. In general, *in vitro* assays appear to be more sensitive to variations in the levels of protease inhibitors, though at different proportions. Considering that this test aims to mimic what would occur *in vivo*, then, could be considered that a lower sensitivity of *in vitro* methods to these protease inhibitors could be more interesting. However, if the goal is to understand differentiation between samples as a result of heat treatment, then *in vitro* tests that are more sensitive to all factors would be most relevant.

Some authors express *in vitro* digestibility data as direct percentages of a standard protein result, such as casein. This form of expression may reduce differentiation in the data between the different methods employed, allowing comparison of data generated by the various methods. Table 3 shows digestibility data in relation to casein (Casein value = 100%). Although many results were similar, there is still a wide range of results for the same sample. These observations emphasize the need for care when comparisons between published studies are performed; the same sample assessed using different methods can produce different conclusions. Therefore, it is recommended to compare data produced using the same method. Urbano et al. (2005) also found different values for chickpea samples when two *in vitro* (Hsu et al., 1977; and Akeson & Stahman, 1964, methods) and one *in vivo* methods were used.

Table 4 shows aminograms and amino acid scores for each sample, calculated according to WHO/FAO/UNU (2007) patterns for 1- to 2-year old children. Considering the results of *in vitro* assays for the calculation of PDCAAS (Table 5), even with the difficulties in establishing correlations between digestibility methods (Table 2), strong correlations were observed (at $p < 0.0500$) between PDCAAS results produced by *in vitro* and *in vivo* digestibility results (Table 6).

Although we observe variation between the values obtained using each method (Table 5) for a given sample, the use of *in vitro* methods for calculating PDCAAS may be promising. In addition to the strong correlations found (Table 6), when the results were expressed as % casein (Table 7), the percentage values were much closer to those obtained with the use of *in vivo* methods (PDCAAS-TD), and several *in vitro* results were not significantly different from the PDCAAS data calculated using TD (Table 7). For example, among the eight PDCAAS-TNBS results, four of them significantly coincided with PDCAAS-TD. In addition, all trials using *in vitro* data were able to point to the sample of heated albumin as the highest values, and they were also able to point to globulins as the worst quality proteins, which coincides with PDCAAS-TD results. In fact, PDCAAS-ppKJ, PDCAAS-3-Enz and PDCAAS-4-Enz methods presented the highest correlations with *in vivo* method, $r = 0.9316$, 0.9442 and 0.9649 ($p < 0.0500$), respectively (as shown in Table 6).

4. Conclusions

While there have been a great difficulty in establishing a correlation between the results of protein digestibility generated by *in vitro* methods, the data generated in this study suggest that *in vitro* digestibility results can be apply to calculate the PDCAAS with prospects of good

correlations with results generated by *in vivo* tests. The lower results found for correlations between PDCAAS-*in vivo* and PDCAAS-*in vitro* were found when the PDCAAS-TNBS was used ($R = 0.8755$, and $p = 0.0013$). The highest correlation was $r = 0.9649$ ($p = 0.0104$), considering PDCAAS-TD X PDCAAS-4-Enz. Considering the three *in vitro* methods that generated the highest correlation, perhaps 3-Enz method could be considered the highlight, in the conditions of this work, presenting as less labor and less time consuming. Moreover, the application of *in vitro* methods allowed for differentiation between samples of chickpea protein, including showing sensitivity to differentiate samples after heat treatment. In addition, the tests showed other important advantages over *in vivo* assays such as simplicity, convenience, speed, lower cost and specially the reduction in animal usage. Thus, the use of *in vitro* methods for this digestibility analysis deserves further study and discussion, with a special focus on seeking the standardization of one method as the most appropriate. Larger studies will be important as they can consider greater numbers and types of proteins, including different compositions of samples with variations in presence of anti-nutritional and other components that could interfere with protein digestibility. The choice of a method as a reference will reduce the present difficulties in comparing data, as this study demonstrated that the same sample may display different results when different methods are applied.

References

- Akeson, W. R., & Stahman, M. A. A. (1964). Pepsin-pancreatin digest index of protein quality evaluation. *The Journal of Nutrition*, *83*, 257–261.
- Association of Official Analytical Chemists (1990). *Official methods of analysis*. Association of Official Analytical Chemists: Washington D. C.
- Association of Official Analytical Chemists (2000). *Official methods of analysis of Association of Official Analytical Chemists: Gaithersburg, MD. Official method 982*. (pp. 30), 30.
- Boye, J., Wijesinha-Bettoni, R., & Burlingame, B. (2012). Protein quality evaluation twenty years after the introduction of the protein digestibility corrected amino acid score method. *The British Journal of Nutrition*, *108*, S183–S211.
- Church, F. C. C., Swaisgood, H. E., Porter, D. H., & Catignani, G. L. (1983). Spectrophotometric assay using o-phthalaldehyde for determination of proteolysis in milk and isolated milk proteins. *Journal of Dairy Science*, *66*, 1219–1227.
- Fields, R. (1972). The rapid determination of amino groups with TNBS. *Methods in Enzymology*, *25*, 464–468.
- Hsu, H. W., Vavak, D. L., Satterlee, L. D., & Miller, G. A. (1977). A multienzyme technique for estimating protein digestibility. *Journal of Food Science*, *42*, 1269–1273.
- Jahan-Mihan, A., Luhoivy, B. L., El Khoury, D., & Anderson, G. H. (2011). Dietary proteins as determinants of metabolic and physiologic functions of the gastrointestinal tract. *Nutrients*, *3*, 574–603.
- Leser, S. (2013). The 2013 FAO report on dietary protein quality evaluation in human nutrition: Recommendations and implications. *Nutrition Bulletin*, *38*, 421–428.
- McDonough, F. E., Steinke, F. H., Eggum, B. O., Bressani, R., Huth, P. J., Barbeau, W. E., ... Phillips, J. G. (1990). *In vivo* rat assay for true protein digestibility: Collaborative study. *Journal of Association of Analytical Chemistry*, *73*, 801–805.
- Panasiuk, R., Amarowicz, O. R., Kostyra, H., & Sijtsma, L. (1998). Determination of a-amino nitrogen in pea protein hydrolysates: A comparison of three analytical methods. *Food Chemistry*, *62*, 363–367.
- Pappenheimer, J. R., Dahl, C. E., Karnovsky, M. L., & Maggio, J. E. (1994). *Intestinal absorption and excretion of octapeptides composed of D amino acids* Proceedings of the National Academy of Sciences of the United States of America *91*. (pp. 1942–1945).
- Sarwar, G., & McDonough, F. E. (1990). Evaluation of protein digestibility corrected amino acid score method for assessing protein quality of foods. *Journal of the Association of Official Analytical Chemists*, *73*, 347–356.
- Schaafsma, G. (2012). Advantages and limitations of the protein digestibility-corrected amino acid score (PDCAAS) as a method for evaluating protein quality in human diets. *The British Journal of Nutrition*, *108*, S333–S336.
- Spadaro, A. C. C., Draghetto, W., Del Lama, S. N., Camargo, A. C. M., & Greene, L. T. (1979). A convenient manual trinitrobenzenesulfonic acid method for monitoring amino acids and peptides in chromatographic column effluents. *Analytical Biochemistry*, *96*, 317–321.
- Statistica 8.0 (2007). *Statistics computer program for Windows. Vol. 1*, Tulsa: StatSoft Inc. Software.
- Tavano, O. L., Silva Junior, S. I., Demonte, A., & Neves, V. A. (2005). Pancreatic hypertrophy in rats caused by chickpea (*Cicer arietinum* L.) protein intake. *Alimentos e Nutrição*, *16*, 5–10.
- Tavano, O. L., Silva Junior, S. I., Demonte, A., & Neves, V. A. (2008). Nutritional responses of rats to diets based on chickpea (*Cicer arietinum* L.) seed meal or its protein fractions. *Journal of Agricultural and Food Chemistry*, *56*, 11006–11010.
- Trop, M., & Birk, Y. (1970). The specificity of proteinases from *Streptomyces griseus* (Pronase). *Biochemical Journal*, *116*, 19–25.

- Urbano, G., López-Jurado, M., Frejnagel, S., Gómez-Villalva, E., Porres, J. M., Frías, J., ... Aranda, P. (2005). Nutritional assessment of raw and germinated pea (*Pisum sativum* L.) protein and carbohydrate by *in vitro* and *in vivo* techniques. *Nutrition*, *21*, 230–239.
- WHO/FAO/UNU Expert Consultation (2007). Protein and amino acid requirements in human nutrition. *WHO technical reports series no. 935*. Geneva: World Health Organization.
- Wolzak, A., Bressani, R., & Brenes, R. G. (1981). A comparison of *in vivo* and *in vitro* estimates of protein digestibility of native and thermally processed vegetable proteins. *Plant Foods for Human Nutrition*, *31*, 31–43.
- Yvon, M., Chabanet, C., & Pélissier, J. P. (1989). Solubility of peptides in trichloroacetic acid (TCA) solutions. Hypothesis on the precipitation mechanism. *International Journal of Peptide and Protein Research*, *34*, 166–176.