



Contents lists available at ScienceDirect

journal homepage: www.elsevier.com/locate/humimm

Erratum

Erratum to “Interleukin 4 haplotypes of susceptibility to chronic periodontitis are associated with IL-4 protein levels but not with clinical outcomes of periodontal therapy” [Hum. Immunol. 74 (2013) 1688–1695]



Giovana Anovazzi^{a,b}, Livia S. Finoti^{a,b}, Sâmia C.T. Corbi^{a,b}, Yeon J. Kim^b, Andrea M. Marcaccini^c, Raquel F. Gerlach^c, Marisa V. Capela^d, Silvana R.P. Orrico^b, Joni A. Cirelli^b, Raquel M. Scarel-Caminaga^{a,*}

^a Department of Morphology, School of Dentistry at Araraquara, UNESP – São Paulo State University, Araraquara, São Paulo, Brazil

^b Department of Oral Diagnosis and Surgery, School of Dentistry at Araraquara, UNESP – São Paulo State University, Araraquara, São Paulo, Brazil

^c Department of Morphology, Stomatology and Physiology, Dental School of Ribeirão Preto, USP – University of São Paulo, Ribeirão Preto, São Paulo, Brazil

^d Department of Physical Chemistry, Institute of Chemistry at Araraquara, UNESP – São Paulo State University, Araraquara, São Paulo, Brazil

The author has discovered errors in Table 3 of their article. The corrected Table 3 is provided below.

Table 3
Distribution of probing pocket depth (PPD) (mm), bleeding on probing (BOP) (%) and clinical attachment level (CAL) among groups at different time points: the median of the measurements (min–max).

	Groups	Baseline Median (min–max)	45 days Median (min–max)	90 days Median (min–max)	Difference among the periods ^{II}
PPD (mm)	SCP	2.4 (2.0–3.9)	1.9 (1.2–3.8)	1.8 (1.3–3.7)	<0.0001
	SH	1.6 (1.1–2.1) [*]	1.5 (1.2–1.9) [*]	1.2 (1.0–1.9) [*]	NS
	PCP	2.3 (2.1–3.7)	1.6 (1.5–2.3)	1.3 (1.2–1.5)	0.0001
	PH	1.5 (1.1–1.9) [†]	1.2 (1.0–1.6) [†]	1.1 (1.0–1.3) [†]	NS
	Difference between the groups [*]	0.00005 [*] 0.000001 [†]	0.004 [*] 0.0003 [†]	0.002 [*] 0.004 [†]	
BOP (%)	SCP	26.9 (11.3–70.1)	6.8 (2.9–18.7)	3.6 (0.6–10.5)	<0.0001
	SH	0.61 (0–14.8) [*]	0.3 (0–6.0) [*]	0.003 (0–3.0) [*]	0.007
	PCP	16.0 (7.5–54.8)	4.0 (2.4–8.0)	2.5 (0.1–5.0)	0.001
	PH	0.59 (0–13.0) [†]	0.5 (0–3.7) [†]	0.002 (0–1.2) [†]	<0.0001
	Difference between the groups [*]	0.00003 [*] 0.00013 [†]	0.00008 [*] 0.0027 [†]	0.0016 [*] 0.00065 [†]	
CAL (mm)	SCP	2.7 (2.1–4.2)	2.2 (1.5–3.8)	1.8 (1.3–3.7)	0.008
	PCP	2.8 (2.1–5.2)	2.2 (1.5–2.8)	1.6 (1.3–3.1)	0.001
	Difference between the groups [§]	NS	NS	NS	
PPD ≥ 5 mm and CAL ≥ 3 mm (% sites)	SCP	12.0 (3.0–37.0)	4.0 (1.0–13.0)	2.0 (0–7.0)	0.0001
	PCP	13.0 (4.0–29.0)	4.0 (1.0–10.0)	1.0 (0–3.0)	0.0001
	Difference between the groups [§]	NS	NS	NS	

^{*} Significant differences compared with the group SCP the same period.

[†] Differences when compared to group PCP the same period. NS: not significant.

[§] Mann–Whitney test for the evaluations between the groups in the same period.

^{II} Friedman test for evaluation between different periods.

DOI of original article: <http://dx.doi.org/10.1016/j.humimm.2013.08.286>

* Corresponding author. Address: Department of Morphology, School of Dentistry at Araraquara, UNESP – São Paulo State University, CP. 331, CEP 14801-903 Araraquara, São Paulo, Brazil.

E-mail address: raquel@foar.unesp.br (R.M. Scarel-Caminaga).

<http://dx.doi.org/10.1016/j.humimm.2014.10.001>

0198-8859/© 2014 American Society for Histocompatibility and Immunogenetics. Published by Elsevier Inc. All rights reserved.