




Liquid-liquid extraction of protease from cold-adapted yeast *Rhodotorula mucilaginosa* L7 using biocompatible and biodegradable aqueous two-phase systems

Luciana Daniela Lario, Luciana Pellegrini Malpiedi, Jorge Fernando Brandão Pereira, Lara Durães Sette & Adalberto Pessoa-Junior


To cite this article: Luciana Daniela Lario, Luciana Pellegrini Malpiedi, Jorge Fernando Brandão Pereira, Lara Durães Sette & Adalberto Pessoa-Junior (2016) Liquid-liquid extraction of protease from cold-adapted yeast *Rhodotorula mucilaginosa* L7 using biocompatible and biodegradable aqueous two-phase systems, *Separation Science and Technology*, 51:1, 57-67, DOI: [10.1080/01496395.2015.1080276](https://doi.org/10.1080/01496395.2015.1080276)

To link to this article: <https://doi.org/10.1080/01496395.2015.1080276>

 View supplementary material [↗](#)

 Published online: 15 Dec 2015.

 Submit your article to this journal [↗](#)

 Article views: 144

 View related articles [↗](#)

 View Crossmark data [↗](#)

 Citing articles: 6 View citing articles [↗](#)

Liquid-liquid extraction of protease from cold-adapted yeast *Rhodotorula mucilaginosa* L7 using biocompatible and biodegradable aqueous two-phase systems

Luciana Daniela Lario^a, Luciana Pellegrini Malpiedi^{a,b}, Jorge Fernando Brandão Pereira^c, Lara Durães Sette^d, and Adalberto Pessoa-Junior^a

^aDepartment of Biochemical and Pharmaceutical Technology, University of São Paulo, São Paulo, Brazil; ^bDepartamento de Química-Física, Facultad de Ciencias Bioquímicas y Farmacéuticas, Universidad Nacional de Rosario, Rosario, Argentina; ^cDepartment of Bioprocess and Biotechnology, School of Pharmaceutical Sciences, UNESP – Univ Estadual Paulista, Araraquara, SP, Brazil; ^dDepartment of Biochemistry and Microbiology, Institute of Biosciences, University of São Paulo (UNESP), São Paulo, Brazil

ABSTRACT

This work aimed to optimize the extraction of an extracellular protease produced by the cold-adapted yeast *Rhodotorula mucilaginosa* L7 using aqueous two-phase systems (ATPS) comprising polyethylene glycol (PEG) and sodium citrate or sodium tartrate. First, the biocompatibility of the phase forming agents was assessed. The results obtained with PEG-2000, PEG-4000, and PEG-6000 demonstrated that even at large PEG concentrations (32 wt%) the protease maintains its activity after 3 h of reaction, whereas an increase in salt concentration provokes a gradual decrease in protease stability. Subsequently, the partitioning of the protease in both types of ATPS was assessed, evaluating the effect of temperature, molecular weight, and concentration of PEG on protease purification, using two 2³-full factorial designs. The best partitioning conditions were obtained in PEG-6000/sodium tartrate-based ATPS, at 30°C (with a yield of 81.09 ± 0.66% and a purification factor of 2.51 ± 0.03). Thus, considering the biodegradable characteristics of the system, the PEG/sodium tartrate ATPS is a viable and economic low-resolution step in protease purification, with a strong potential for future industrial application.

ARTICLE HISTORY

Received 26 February 2015
Accepted 3 August 2015

KEYWORDS

Protease; cold-adapted microorganisms; liquid-liquid extraction; aqueous two phase systems; purification

Introduction

Currently, the use of enzymes at an industrial level has become a common practice, and the most utilized are the hydrolytic enzymes. Among them, carbohydrases, proteases, and lipases are the dominant enzymes on the market, accounting for more than 70% of all sales. Proteases refer to a group of proteins whose catalytic function is to hydrolyze peptide bonds, and have been used in a large array of applications, mostly in the chemical, food, cosmetic, and pharmaceutical industries. In addition to these main applications, proteolytic enzymes have been applied in leather treatment and in several bioremediation processes (1).

Although proteases are widespread in nature, microorganisms are the preferred source of these enzymes because of their rapid growth, the small space required for cultivation, and the ease with which they can be genetically manipulated to produce enzymes with customized properties (2). Among the several classes of

industrially used microorganisms, the cold-adapted type have been the focus of interest, because the enzymes they produce catalyze reactions at low and moderate temperatures more efficiently than at high temperatures. Thus, the use of these enzymes decreases the energetic and processing costs that are typically associated with high-temperature processes and industrial heating steps (3). Moreover, due to their heat-labile activity, these enzymes can be quickly inactivated by a mild heat treatment, which is useful in many biotechnological applications that require a rapid heat-inactivation rate (3-6).

Traditionally, the purification of extracellular proteases involves an initial fractionation step with ammonium sulfate (a compound of some environmental concern), followed by several steps of chromatography (high cost), which is difficult in their large-scale bio-production (7-9). Thus, as stated by Straathof (10), it is imperative to continue the search for more sustainable, economical, and suitable separation methods.

CONTACT Luciana Daniela Lario ✉ lucianalario@usp.br Department of Biochemical and Pharmaceutical Technology, University of São Paulo, São Paulo, Brazil.

Color versions of one or more of the figures in the article can be found online at www.tandfonline.com/lsst.

Supplemental data for this article can be accessed on the [publisher's website](#).

Since the beginning of the 1970s, aqueous two-phase systems (ATPS) have been suggested as a good alternative for the purification of several biomolecules and other biological products (11). By definition, ATPS are systems that are spontaneously formed by the mixing of two different aqueous solutions above certain concentrations, such as the aqueous solutions of two polymers, one polymer and one salt, or two salts (11-13). Because of the water-rich content of both co-existing phases, ATPS provides a more biocompatible environment for the separation of a broad range of biological materials. Moreover, the majority of components used in the formation of these systems are not expensive, easily recycled, and, similar to other liquid-liquid extraction operations, ATPS can be readily scaled up (14). Although largely studied (15-18), conventional ATPS using polymer and inorganic salts presents several disadvantages, mainly observed when potassium phosphate is employed as the phase forming agent, because the high salt concentration required for ATPS formation leads to environmental concerns, due to the high eutrophication potential of phosphate (19). This phenomenon leads to the depletion of dissolved oxygen in water, resulting in a consequent loss of aquatic life (20-23). Therefore, ATPS composed of biodegradable anions (such as citrate or tartrate) and polyethylene glycol (PEG) has been suggested to be a more biocompatible alternative (14, 24-25). These two salts are particularly interesting because of their low environmental impact and high biodegradability, and because they can be discharged into common biological wastewater treatment plants.

Despite the large number of studies employing traditional polymer/salt ATPS (15-18), these biodegradable ATPS are less studied, mainly to extract proteases from fermentative broths. Thus, in this paper, the extraction and pre-purification of the protease secreted by the cold-adapted yeast *R. mucilaginosa* L7 using biodegradable polymer/salt ATPS is reported. First, the protease stability and activity in different concentrations of each phase forming compound was evaluated. To determine the significance of different parameters (sodium salt anion, PEG molecular weight, PEG concentration, and temperature) on protease purification, a 2³-full factorial design was implemented for each ATPS. In the latter part of the paper the effect of protease re-concentration changing the ATPS volume ratio (V_r) was studied. The goal of this paper was to find the best conditions to separate proteases from the cold-adapted yeast *R. mucilaginosa* L7.

Experimental section

Chemicals

Polyethylene glycols (PEG) 2000, 4000, and 6000 g/mol were purchased from Sigma-Aldrich (St. Louis, MO, USA). Each PEG is identified in the manuscript as the following acronyms: PEG-2000, PEG-4000, and PEG-6000. Both salts used in the study, sodium citrate and sodium tartrate, were also acquired from Sigma-Aldrich (St. Louis, MO, USA). All of the reagents were of analytical grade with a minimum purity of 99% and were used as received. All of the stock solutions were prepared using water purified by filtration through a Millipore Milli-Q ion exchange system (Bedford, MA, USA).

Yeast strain and culture conditions

The yeast studied herein (strain L7) was previously isolated from an Antarctic marine alga and taxonomically identified as *Rhodotorula mucilaginosa* through similarity to the partial 26S rDNA gene, according to the procedure described by Duarte et al. (26). The strain was deposited at the Brazilian Collection of Environmental and Industrial Microorganisms (Paulinia, Brazil) under the access number CBMAI 1528. The microorganism was maintained in potato-dextrose-agar (PDA) solid medium at 15°C in a dark room.

The *R. mucilaginosa* L7 strain was grown in Sabouraud-dextrose liquid medium (compose of 40 g/L dextrose and 10 g/L peptone) at 25°C, pH 5.6 with shaking at 150 rpm until the exponential phase (18 h). One mL aliquots from this culture were pipetted into 250 mL flasks containing 50 mL of fresh Sabouraud-dextrose medium and were grown at 25°C, 150 rpm, and an initial pH of 5.6 for 48 h. Then, the cells were removed by centrifugation at 3500 xg in a centrifuge model Jouan B4i (Thermo, Waltham, MA, USA). The supernatant, which exhibited an enzymatic activity equal to 200 U/mL, was stored in an ultrafreezer at -70°C for use as the protease source in the partitioning studies.

Evaluation of proteolytic activity and stability

Protease stability was evaluated in the presence of the polymers, PEG-2000, PEG-4000, and PEG-6000, or the biodegradable salts sodium citrate and sodium tartrate. For this purpose, stock solutions of each polymer (48 wt%) and salt (22 wt% and 34 wt%, pH 5.0, for the sodium tartrate and sodium citrate,

respectively) were prepared. Then, different volumes of each stock solution were mixed with 75 μL of crude extract, and brought to a final volume of 150 μL with purified water. Each mixture was placed at 20°C and aliquots of the solutions were withdrawn at time intervals of 0, 1, 2, and 3 h to determine the residual proteolytic activity at 50°C, as described below.

The percent of residual proteolytic activity (A_r ,%) of each solution was determined by calculating the ratio of the enzymatic activity after treatment (A , U/mL) to that of the initial extract (A_0 , U/mL), according to the following equation:

$$A_r(\%) = \frac{A}{A_0} \times 100$$

Protease partitioning studies using the biodegradable ATPS

According to our previous studies, the protease is stable at acidic pH, and has an optimum pH of 5.0 (9). Thus, the selected ATPS were systems of PEG/salt whose phase diagrams had been determined at an acidic pH. PEG/sodium tartrate and PEG/sodium citrate-based ATPS previously reported (26, 27), were chosen to develop the respective assays. The ATPS were prepared with sodium citrate or sodium tartrate and various PEG molecular weights (MW) and concentrations. All of the mixture points were prepared according to the previously determined phase diagrams (27-28). All of the components of the ternary systems were added to 15 mL glass tubes by weight, and mixed by turning the tubes up-side down several times. After complete homogenization, 4.6 g of the culture supernatant (free of cells) was added to the tubes and brought to 10.0 g with deionized water. Then, all of the systems were mixed at 8 rpm for 30 min at 25°C in an end-to-end mixer (Branstead/Thermolyne, Dubuque, IA, USA). After mixing, the ATPSs were stood at each temperature of interest (4, 17, and 30°C) in a thermo bath model 521/2DE (New Ethics, Vargem Grande Paulista, SP, Brazil) until they reached two-phase separation (at least 1 h). After the phase settling, the volumes of the upper and lower phases were measured and aliquots from both phases were carefully collected using Pasteur-pipettes for the subsequent quantification of proteolytic activity and total protein concentration. The partitioning behavior of the protease in each ATPS was evaluated according to the protease partition coefficient (K_e), protease yield (Y ,%) and purification factor (PF). All of the

partitioning parameters and their respective equations are detailed below.

The protease partition coefficient (K_e) was defined as the ratio of the enzymatic activity (U/mL) in the top phase (A_{Top}) to that in the bottom phase (A_{Bot}):

$$K_e = \frac{A_{Top}}{A_{Bot}}$$

The protease yield (Y ,%) was determined by the ratio of the total activity (U) in the top phase to that added in the starting extract, and expressed as a percentage:

$$Y(\%) = \left(A_{Top} * V_{Top} / A_0 * V_0 \right) \times 100$$

where V_{Top} and V_0 are the volumes of the top phase and of the crude extract added to the system, respectively.

The protein partition coefficient (K_p) was calculated as the ratio of the equilibrium concentration (mg/mL) of the total protein in the top phase $[P]_{Top}$ to that in the bottom phase $[P]_{Bot}$:

$$K_p = \frac{[P]_{Top}}{[P]_{Bot}}$$

The purification factor (PF) of the protease was calculated as the ratio of the specific activity (U/mg) in the phase where it was preferentially partitioned (top phase) to the specific activity in the culture supernatant before partitioning:

$$PF = \frac{A_{Top}}{C_{Top}} \bigg/ \frac{A_0}{C_0}$$

where C_{Top} and C_0 are the total protein concentrations (expressed as mg/mL) in the top phase and the crude extract, respectively.

Determination of proteolytic activity

Proteolytic activity was measured by digestion of azocasein (Sigma, St. Louis, MO, USA), in accordance with Charney and Tomarelli (29). An appropriate dilution of culture supernatant (150 μL) was incubated with 150 μL of 0.5 wt% azocasein in 50 mM sodium acetate buffer (pH 5.0) during 40 min at 50°C. The reaction was then stopped by adding 150 μL of 10% (w/v) trichloroacetic acid. After centrifugation of the reaction mixture, 100 μL of the supernatant was mixed with 100 μL of 0.5 M KOH, and the absorbance at 430 nm was measured against an appropriate blank. All of the samples were measured in duplicate, and the respective proteolytic activity was expressed in units of enzyme activity (U). One U was defined as the amount of enzyme leading to a

0.001 increase in the absorbance under the assay conditions. The enzyme activity was determined using a microplate spectrophotometer (Spectra Max Plus 384, Molecular Device, Sunnyvale, CA, USA).

Determination of total protein concentration

The total protein content in both phases of all of the systems was determined using the Bicinchoninic Acid (BCA) method according to previously reported procedures (30). To avoid interference from the phase components, samples were diluted at least tenfold with distilled water and 10 μ L of each was pipetted in triplicate into a 96 well microplate. Bovine serum albumin (BSA) standards were made ranging from 1,500 μ g/mL to 25 μ g/mL, and 10 μ L of each standard was pipetted in triplicate into the same plate. For blanks, 10 μ L of distilled water was pipetted in triplicate into the plate. Next, the working BCA reagent was prepared by diluting 1 part of copper (II) sulfate pentahydrate 4% (w/v) solution (Sigma Aldrich, St. Louis, MO, USA) into 50 parts bicinchoninic acid solution (Sigma Aldrich, St. Louis, MO, USA), and 200 μ L of the working reagent was placed in each well. The plate was briefly mixed on a plate shaker and incubated at 37°C for 30 min. The UV/Vis measurements were performed in a microplate spectrophotometer at 562 nm (Spectra Max Plus 384, Molecular Device, Sunnyvale, CA, USA). The protein concentration was determined by comparison with the absorbance values of the BSA standard curve.

Statistical analysis

Taking into account the large number of variables in the study, two 2³-full factorial experimental designs were performed: one for PEG/sodium tartrate ATPS and another for PEG/sodium citrate ATPS. In both cases, the independent variables (X_i) were PEG molecular weight (PEG MW) (X_1), PEG concentration [PEG] (wt%) (X_2), and temperature (T) (X_3), whereas the responses (Y_i) were protease partition coefficient (K_c) (Y_1), purification factor (PF) (Y_2), and protease yield in the top phase (Y_3) (Y_3).

As shown in Table 1, three different levels were adopted for each independent variable, namely PEG-2000, PEG-4000, and PEG-6000 for X_1 ; 13.00, 13.50, and 14.00 wt% for X_2 (in PEG/Sodium Tartrate ATPS); 15.01, 15.24, and 15.48 wt% for X_2 (in PEG/Sodium Citrate ATPS); and 4, 17, and 30°C for X_3 , respectively. It is important to note that the ATPS compositions (27–28) were selected to avoid a significant loss of protease activity in the presence of high salt concentrations (see

Table 1. Factor levels of the 2³ experimental designs used for the study of protease partitioning in PEG/sodium tartrate (a) and PEG/sodium citrate (b) ATPS.

Variables	Levels		
	Low (-1)	Central (0)	High (+1)
PEG MW (g/mol)	2000 (a, b)	4000 (a, b)	6000 (a, b)
[PEG] (wt%)*	13.00 (a)	13.50 (a)	14.00 (a)
	15.01 (b)	15.24 (b)	15.48 (b)
Temperature (°C)	4	17	30

*A maximum concentration of PEG was defined for all of the ATPS studied (16 wt%) because the protease loses activity at high salt concentrations and high polymeric concentrations induces the precipitation of azocasein not allowing the respective protease to be measured.

below in Section titled “Influence of Salts and Polymers on Protease Stability” and Fig. 1).

For statistical elaboration, the actual values of each independent variable were coded to give -1, 0, and +1 coded levels according to the equation:

$$x_i = \frac{X_i - X_0}{\Delta X_i}$$

where x_i represents the corresponding coded values, X_0 the actual values at the central point, and ΔX_i the step change value. Both experimental design approaches were performed with four repetitions at the central point.

Furthermore, due to several experimental limitations related to proteolytic quantification (at high polymeric concentrations the precipitation of azocasein occurs), a maximum PEG concentration was defined for all ATPS (lower than 16 wt%). It is important to note that in both experimental design approaches, the same amount of crude extract (4.6 g) (which correspond to 200 U/mL of total protease activity) was used. Similar concentrations of PEG-2000, PEG-4000, or PEG-6000 were combined with aqueous solutions of sodium tartrate (10 wt %) and sodium citrate (11.5 wt%) to perform the experimental design analysis.

After the experiments at pre-determined conditions, the statistical results were determined with the aid of the software *Statistica* version 10 (StatSoft, Tulsa, OK, USA).

Results

Influence of salts and polymers on protease stability

Before performing the protease partitioning, an analysis of its stability in the presence of each phase forming agent was carried out. The stability of the protease secreted by *R. mucilaginosa* L7 was investigated by incubating the enzyme without the substrate for 3 h, in the presence of PEG-2000, PEG-4000, and PEG-




Figure 1. Residual protease activity (%) of *R. mucilaginosa* L7 protease at different concentrations of: a) sodium tartrate; b) sodium citrate; and c) PEG with different molecular weights (2000, 4000, and 6000 g/mol). The stability of the enzyme was determined at 1, 2 and 3 h at 20°C. The residual enzyme activity was assessed with 0.5 % (w/v) azocasein at pH 5.00 and 50°C.

6000, or with the sodium citrate and sodium tartrate aqueous solutions. Afterwards, the residual proteolytic activity in 0.5% (w/v) azocasein was evaluated. The concentrations of each PEG and salt aqueous solutions were defined according to the previously reported phase diagrams (27-28).

As shown in Figs. 1a and 1b, a gradual decrease in protease stability was observed with an increase in salt concentration; the highest reduction was observed after 3 h of incubation with 18 wt% of added salt (in the presence of sodium tartrate and sodium citrate the residual protease activities were 20% and 25% lower than the initial values, respectively). A detailed analysis of the protease activity of Fig. 1b clearly indicates that after a small addition of sodium citrate (4 wt%) no negative deviations were observed after 1, 2, or 3 h of incubation (a small positive deviation was observed after 3 h of incubation, which can be related to some interferences in the azocasein quantification). However, when a large amount of sodium citrate was added, a significant decrease in protease activity was observed after 2 h of incubation. From the analysis of Fig. 1a,

small deviations in the protease activity were observed with different sodium tartrate aqueous solutions (4, 8, or 18 wt%), but in all cases, the protease only began to lose activity after 2 h of incubation. The mechanisms of proteins-ion interactions are poorly understood. However, previous evidence has indicated that several kosmotropic anions, such as tartrate and citrate, can reduce the hydration shell of macromolecules and consequently induce the process of protein aggregation (31). Accordingly, the decrease of proteolytic activity (observed in Figs. 1a and 1b) can be related to the low protein solubility as a consequence of its aggregation in the presence of a high concentration of sodium tartrate or sodium citrate.

In contrast, the presence of different PEGs did not affect protease stability, as exhibited in Fig. 1c. The results obtained with PEG-2000, PEG-4000, or PEG-6000 clearly demonstrated that even at large PEG concentrations (32 wt%), the protease maintains its activity after 3 h of reaction. These results were expected because PEG is frequently used as a polyhydroxy stabilizer to improve the biocompatibility of proteins for

both *in vivo* and *ex vivo* applications (32). Similar results have been reported for other enzyme-PEG pairs, in which it was demonstrated that in the presence of PEGs with molecular weight above 4000 g/mol, the enzymatic activity was not significantly affected due to an unspecific exclusion of the polymer from the protein surface and a structure-stabilizing effect, as predicted by Timasheff's theory (33).

Protease partitioning with PEG/sodium tartrate and PEG/sodium citrate-based ATPS

In light of the protease stability results, in which a decrease in enzymatic activity with an increase in the concentration of salts was demonstrated, we decided to work with the minimal salt concentration that permits the formation of each proposed ATPS (27-28). Thus, for all assays, the salt concentration was defined: 10.0 wt% for sodium tartrate and 11.5 wt% for sodium citrate-based ATPS, and the PEG MWs (g/mol), PEG concentrations and temperatures varied as presented in Table 1. The results of the protease partitioning in each PEG/sodium tartrate and PEG/sodium citrate-based ATPS are summarized in Tables 2 and 3, respectively.

A quick analysis of Tables 2 and 3 shows that in all of the ATPS studied, the K_e values are higher than 1, indicating a preferential partitioning of the protease to the top phase (PEG-rich phase). This preferential

Table 2. Experimental values of the protease partitioning with PEG/sodium tartrate ATPS. K_e is the protease partition coefficient, PF is the protease purification factor of the top phase and Y (%) is the protease yield of the top phase. The ternary systems were chosen according to the 2^3 factorial design with centred face and four repetitions at the central point.

Ternary system	PEG MW (g/mol)	[PEG] (wt%)	T (°C)	K_e	PF	Y (%)
1	2000	13.00	4	2.46 ± 0.05	1.68 ± 0.04	74.71 ± 0.86
2			30	2.60 ± 0.02	1.56 ± 0.02	69.63 ± 0.53
3	4000	14.00	4	2.94 ± 0.02	1.49 ± 0.01	66.11 ± 0.43
4			30	2.90 ± 0.06	1.53 ± 0.01	62.22 ± 0.36
5	6000	13.00	4	3.71 ± 0.01	1.40 ± 0.04	78.07 ± 0.76
6			30	3.90 ± 0.02	2.51 ± 0.03	81.09 ± 0.66
7	4000	14.00	4	4.06 ± 0.07	2.42 ± 0.05	74.93 ± 0.76
8			30	3.77 ± 0.03	2.12 ± 0.07	71.85 ± 0.63
9	4000	13.50	17	3.23 ± 0.06	1.93 ± 0.03	72.32 ± 0.84
10			17	3.15 ± 0.02	2.01 ± 0.02	71.10 ± 0.46
11	4000	13.50	17	3.17 ± 0.06	2.00 ± 0.01	71.24 ± 0.35
12			17	3.06 ± 0.02	2.02 ± 0.08	70.30 ± 0.90

Table 3. Experimental values of the protease partitioning with PEG/sodium citrate ATPS. K_e is the protease partition coefficient, PF is the protease purification factor of the top phase and Y (%) is the protease yield of the top phase. The ternary systems were chosen according to the 2^3 factorial design with centred face and four repetitions at the central point.

Ternary System	PEG MW (g/mol)	[PEG] (wt%)	T (°C)	K_e	PF	Y (%)
1	2000	15.01	4	4.58 ± 0.01	1.53 ± 0.01	55.74 ± 0.14
			30	6.06 ± 0.18	1.51 ± 0.04	61.24 ± 0.82
3	4000	15.48	4	7.63 ± 0.10	1.62 ± 0.02	57.95 ± 0.76
			30	13.64 ± 0.06	1.68 ± 0.01	65.01 ± 0.30
5	6000	15.01	4	5.38 ± 0.11	2.13 ± 0.16	58.50 ± 0.46
			30	7.31 ± 0.05	1.80 ± 0.01	58.61 ± 0.42
7	4000	15.48	4	8.74 ± 0.11	2.01 ± 0.03	58.58 ± 0.71
			30	11.97 ± 0.03	2.18 ± 0.06	66.28 ± 0.68
9	4000	15.25	17	5.19 ± 0.05	2.00 ± 0.07	62.72 ± 0.45
			17	5.47 ± 0.06	1.85 ± 0.02	62.81 ± 0.65
11	4000	15.25	17	5.56 ± 0.03	2.00 ± 0.08	62.45 ± 0.41
			17	5.42 ± 0.02	2.07 ± 0.09	63.58 ± 0.83

migration of protease to the PEG-rich phase is extremely interesting when the aim is to extract enzymes from fermentation broths, because cellular debris and other particulate matter typically remain in the bottom phase (salt-rich phase) (34).

In Table 2, there is a comparison of the different protease partitioning parameters for all of the PEG/sodium tartrate ATPS. The highest value observed for Y (%) ($81.09 ± 0.66\%$) and for PF ($2.51 ± 0.03$) occurred in assay 6, which was carried out at 30°C, 10 wt% of sodium tartrate, and 13 wt% of PEG-6000. Contrarily, the worst purification values were obtained from assays 1 through 4; all of these experiments using PEG-2000 based ATPS.

All of the PEG/sodium citrate-based ATPS are depicted in Table 3. The highest values of Y (%) (66.28%) and PF ($2.18 ± 0.03$) were obtained in assay 8, which corresponds to a system composed of 11.5 wt % of sodium citrate and 15.48 wt% of PEG-6000 and prepared at 30°C.

However, taking into account the several variables under study, statistical analysis was used to evaluate the responses of Y (%) to the PEG/sodium tartrate and PEG/sodium citrate ATPS. The results are presented in Figs. 2 and 3, respectively. Analyzing the Pareto chart, the length of each bar is proportional to the standardized effect of the related variable or interaction, and the bars extending beyond the vertical




Figure 2. Pareto chart of ANOVA applied for the evaluation of the effects of different PEG molecular weights (PEG MWs, g/mol) (X_1), PEG concentrations ([PEG], wt%) (X_2), and Temperature (T, °C) (X_3) in protease partitioning using the PEG/sodium tartrate ATPS obtained for Y (%) as response.




Figure 3. Pareto chart of ANOVA applied for the evaluation of the effects of different PEG molecular weights (PEG MWs, g/mol) (X_1), PEG concentrations ([PEG], wt%) (X_2), and temperature (T, °C) (X_3) in protease partitioning using the PEG/sodium citrate ATPS obtained for Y (%) as response.

line correspond to the statistically significant effects at a confidence level of 95%. Thus, according to Fig. 2, it was observed that in the PEG/sodium tartrate system the strongest effects on Y (%) were those exerted by PEG MW (X_1) (positive sign) and PEG concentration (X_2) (negative sign). This statistical analysis means that such parameters would be improved by a simultaneous decrease in concentration and increase in the MW of PEG. Conversely, when the Pareto chart was analyzed for the PEG/sodium citrate ATPS (Fig. 3), the significant variables influencing Y (%) were PEG concentration (X_2) and temperature (X_3), both with positive signs. According to this analysis, an increase in the temperature and/or the PEG concentration causes an increase in Y (%).

In summary, according to the statistical analyses, it is clear that the protease partitioning in the PEG/sodium tartrate or PEG/sodium citrate is controlled by different factors for each set of ATPS. The MW and PEG concentration were the main limiting factors of the first set of ATPS, while the temperature and PEG concentration were the driving factors for the second set. A general comparison between both systems clearly indicates that the Y (%) and PF are significantly higher in the PEG/sodium tartrate ATPS. However, when it is compared to the K_e obtained in both series of ATPS, the K_e values are usually superior in PEG/sodium citrate ATPS. This opposite behavior may be a result of the protein aggregation capacity of citrate salts, as explained in the section titled “Influence of Salts and Polymers on Protease Stability” (35). It is important to note that these results were also confirmed by the total mass balance results which were always lower than 100% in the PEG/sodium citrate ATPS (data not shown).

Taking into account all of the results presented above, the protease concentration and partitioning optimization assays were performed using the ATPS composed of sodium tartrate as the salt and PEG-6000 (high MW leads to high Y (%) and PF) at the lowest possible concentration (a decrease in concentration increases the Y (%) and PF) at 30°C.

Protease concentration and partitioning optimization

A complementary reduction of volume and an increase in the purification factor are fundamental when the aim of the liquid-liquid extraction is an industrial separation process (36). Thus, in the last part of our work we studied the effect of volumetric ratio (V_r) in protease partitioning using PEG/sodium tartrate ATPS, which exhibited the best purification performances (defined with the statistical analysis of the responses in terms of Y (%) and the FP). Five mixture points of ATPS were prepared by combining different compositions of PEG-6000 (with the lowest possible concentration defined according to the critical point of the phase diagram), sodium tartrate, and crude extract (because a small amount of each phase forming component was used, the amount of crude extract added was increased to 5.18 g). The protease concentration studies were prepared to obtain a V_r between 0.3 and 12 (as depicted in Figs. 4a and 4b). After the preparation of the ATPS, the partitioning tests were performed at 30°C, as previously described in the section titled “Protease Partitioning Studies using the Biodegradable ATPS.” A detailed composition and V_r of each PEG/sodium tartrate




Figure 4. Effect of different volume ratios (V_r is the volumetric ratio between top and bottom phase) in protease partitioning using the system PEG-6000/sodium artrate-based ATPS, in which: a) corresponds to the experimental ternary phase diagram with the mixture points studied (numbering 1 to 5), tie-line (grey line) and binodal curve (black line); b) shows the experimental values of volume ratio (V_r); and c) shows the experimental values of protease yield Y (%) and purification factor (PF).

ATPS is presented in Supplementary Information (Table S1).

As observed in Fig. 4c, a decrease in V_r resulted in a decrease of Y (%), and a proportional increase in PF . The increase observed in the PF was expected, because at lower V_r , the volumetric activity (U/mL) was higher in the top phase (high values of K_e), while the concentration of total protein between the co-existing phases remained almost constant ($K_p \approx 0.8$) (Table S1). These results are in accord with those of previous studies, in which a strong influence of V_r in the K_e was demonstrated (37-39). The reduction observed in the Y (%) parameter with a decrease in V_r can be attributed to the reduction of the top phase volume, and the subsequent reduction of the total protease activity (U) in this phase.

For a comparison, a summary of the results obtained in previous work in protease purification

using different types of PEG/salt ATPS is presented in Table 4. With respect to the type of salt chosen for protease recovery, Bacha et al. (2012) evaluated the influence of potassium phosphate, sodium sulfate, ammonium sulfate, and sodium citrate solutions (50% w/v) in combination with PEG-4000 (50% w/v) on the purification of an extracellular protease from *Serratia marcescens* P3 (40). These authors achieved maximum yields (approximately 68%) and purification factors (2.7-fold) with the systems containing sodium citrate. Similarly, for the PEG/citrate ATPS (composed of 11.5 wt% of sodium citrate and 15.48 wt% of PEG-6000), we obtained maximum yields of 66.28% and PF of 2.18 ± 0.03 . It is important to highlight that we used lower concentrations of sodium citrate to diminish the effect of the higher concentrations of salts on protein stability and to achieve a more

Table 4. Comparison of proteases purification (*PF*) and yields (*Y*, %) obtained from different biological sources using aqueous two-phase systems (ATPSs).

ATPS	Biological source	<i>Y</i> (%)	<i>PF</i>	Ref
PEG-6000/Sodium Tartrate	<i>Rhodotorula mucilaginosa</i>	81	2.5	current work
PEG-600/Sodium Tartrate	Bovine pancreas	87	2.8	Pellegrini et al. 2011 (38)
PEG) 4000-sodium citrate	<i>Serratia marcescens</i>	68.7	2.77	Bacha et al. 2012 (40)
PEG-1000/Sodium Citrate	<i>Clostridium perfringens</i>	131	4.2	Porto et al. 2008 (41)
PEG-4000/Potassium Phosphate	<i>Penicillium roqueforti</i>	56	3.1	Pericin et al. 2008 (42)
PEG-8000/Potassium Phosphate	<i>Azadirachta indica</i> leaves	85	1.1	Subattha et al. 2012 (43)
PEG-1000/Potassium Phosphate	<i>Bacillus subtilis</i>	62	6.1	Chouyok et al. 2005 (44)
PEG-6000/Potassium Phosphate	<i>Lentinus citrinus</i>	151	1.1	Kirsh et al. 2012 (45)
PEG-6000/Potassium Phosphate	<i>Escherichia coli</i>	95	1.7	Loc et al. 2013 (46)

economic and ecological system. Porto et al. (2008) (41) also used an ATPS composed of PEG-10000 and citrate-salts to remove proteases from a *Clostridium perfringens* fermentation broth. Similarly to our results, these authors found an increase in protease recovery in the top phase with a decrease in citrate concentration (*Y* of 131% and *PF* of 4.2).

In the case of PEG/tartrate ATPS used for the purification of proteases, Pellegrini et al. (38) evaluated the capability of PEG/sodium tartrate-based ATPS to purify α -chymotrypsinogen from bovine pancreas at a pH of 5.0. The authors found that the systems based on PEG-600 (20.62 wt%) and sodium tartrate (12.85 wt%) with a V_r equal to one have the highest enzyme purification performance (*PF* = 2.8; *Y* = 87%). However, in contrast to our results, which demonstrated that it is possible to concentrate and increase the purity of the protease in a PEG-rich phase using PEG-6000/sodium tartrate-based ATPS by decreasing the V_r , these authors suggested that volume ratio changes were not suitable variables to improve the protease purifying performance in PEG-600 and PEG-4000 ATPSs.

A large number of studies that analyzed the recovery of proteases in PEG/phosphate ATPS have been performed (42-46), in which the *PF* and *Y* values for the partitioning of the proteases were similar to the values reported by our group. However, the concentration of phosphate used in these systems was between 15-50 wt %, which leads to environmental concerns, avoiding a large industrial application. Phosphate pollution is recognized as a major pollution problem throughout the world and some countries have introduced control acts to limit the use of phosphate-based compounds. In fact Switzerland banned the use of phosphate in detergents (47). A report prepared by the Swedish Chemical

Agency (2006) (48) indicated a series of substances that should be used instead of phosphates. According to that report, citrate salts, at concentrations between 2.5 wt% and 15 wt% were found to be easily biodegradable, not bio-accumulative and non-toxic to aquatic organisms. Likewise, tartrate was found to have a low toxicity and was not bio-accumulative. Thus, it is evident that the ATPS studied in this work (with citrate and tartrate sodium salts) are particularly interesting because of their low environmental impact and high biodegradability and, because they can be discharged into common biological wastewater treatment plants.

Conclusions

The potential applicability of biodegradable ATPS to purify proteases directly from *Rhodotorula mucilaginosa* L7 culture was studied. First, it was demonstrated that the protease can remain fully stable in different aqueous solutions of PEG (different MWs and concentrations) and in the presence of low concentrations of sodium tartrate or sodium citrate aqueous solutions. The protease purification results revealed that the best conditions to purify the protease were obtained in the system composed of PEG-6000 and sodium tartrate at 30°C. Additionally, with an experimental design and subsequent statistical analysis, it was shown that the PEG MW or PEG concentration had a strong influence on protease purification parameters.

As discussed, our protease recovery results are similar to or even higher than those reported by other authors, where PEG/salt ATPS were also used to purify different enzymes (38, 40-46). Furthermore, although the *PF* obtained in our study were generally lower than other systems reported in the literature, it is important to note that this parameter is directly related with to the initial protease/total protein ratio. In our case, because *R. mucilaginosa* L7 secretes a small amount of proteins into the extracellular medium (49), the protease sample purified with our biodegradable ATPS exhibited a high degree of purity.

Finally, in order to optimize the protease extraction and its concentration in small volume size, the influence of volume ratio on the enzyme recovery and purification was also evaluated. Our results demonstrated that it is possible to concentrate and to increase the purity of the protease in a PEG-rich phase using PEG-6000/Sodium Tartrate-based ATPS, since a decrease of V_r resulted in the increment on the protease *PF*. Taking into account all of the interesting purification results and the biocompatible (high protease stability) and biodegradable characteristics of PEG and sodium tartrate, in addition to their low costs, it is

clear that PEG/sodium tartrate-based ATPS may be a potential and viable liquid-liquid technique to purify proteases for industrial purposes.

Funding

This work was funded by the Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP PhD fellowship process n° 2012/23726-4 awarded to Luciana Lario, FAPESP PhD fellowship process n° 2011/20521-0 awarded to Luciana Pelegrini and FAPESP grant n° 2013/19486-0 awarded to Lara Sette).

References

- [1] Li, S.; Yang, X.; Yang, S.; Zhu, M.; Wang, X. (2012) Technology prospecting on enzymes: Application, marketing and engineering. *Comput. Struct. Biotechnol. J.*, 2:1–11.
- [2] Rao, M. B.; Tanksale, A. M.; Ghatge, M. S.; Deshpande, V. V. (1998) Molecular and biotechnological aspects of microbial proteases. *Microbiol. Mol. Biol. Rev.*, 62(3): 597–635.
- [3] Cavicchioli, R. (2006) Cold-adapted archaea. *Nat. Rev. Microbiol.*, 4(5): 331–343.
- [4] Margesin, R. F.; Feller, G.; Gerday, C.; Russell, N. J. (2002) Cold-adapted microorganisms. Adaptation strategies and biotechnological potential. In *The Encyclopedia of Environmental Microbiology*; Bitton, G., Ed.; Wiley: New York, pp. 871–885.
- [5] Kasana, R. C. (2010) Proteases from psychrotrophs: An overview. *Crit. Rev. Microbiol.*, 36(2):134–145.
- [6] Feller G. (2013) Psychrophilic enzymes: From folding to function and biotechnology. *Scientifica*, 2013:1–28.
- [7] Ma, C. L.; Ni, X. M.; Chi, Z. M.; Ma, L. Y.; Gao, L. M. (2007) Purification and characterization of an alkaline protease from the marine yeast *Aureobasidium pullulans* for bioactive peptide production from different sources. *Mar. Biotechnol.*, 9: 343–351.
- [8] Nouani, A.; Belhamiche, N.; Slamani, R.; Belbraquet, S.; Fazuouane, F.; Bellal, M. M. (2009) Extracellular protease from *Mucor pusillus*: Purification and characterization. *Int. J. Dairy Technol.*, 62(1): 112–117.
- [9] Abirami, V. M.; Meenakshi, S. A.; Kanyamathy, K.; Bharathidasan, R.; Mahalingam, R.; Panneerselvam, A. (2011) Partial purification and characterization of an extracellular protease from *Penicillium janthinellum* and *Neurospora crassa*. *European J. Exp. Biol.*, 1(3): 114–123.
- [10] Straathof, A. J. J. (2011) The proportion of downstream costs in fermentative production processes In *Comprehensive Biotechnology*; M. Y. Murray, ed.; 2nd ed.; vol. 2, pp. 811–814, Academic Press: Burlington, Mass.
- [11] Albertsson, P. A. (1971) *Partition of Cell Particles and Macromolecules*; 2nd ed., Wiley: New York.
- [12] Zaslavsky, B. Y. (1994) *Aqueous Two-Phase Partitioning: Physical Chemistry and Bioanalytical Applications*; Dekker: New York.
- [13] Li, M. P.; Peeples, T. L. (2004) Purification of hyperthermophilic archaeal amyolytic enzyme (MJA1) using thermoseparating aqueous two-phase systems. *J. Chromatogr. B Analyt. Technol. Biomed. Life Sci.*, 807(1): 69–74.
- [14] Zafarani-Moattar M. T.; Hamidi, A. A. (2003) Liquid-liquid equilibria of aqueous two-phase (polyethyleneglycol)-potassium citrate system. *J. Chem. Eng.*, 48:262–265.
- [15] Asenjo, J. A.; Andrews, B. A. (2012) Aqueous two-phase systems for protein separation: phase separation and applications. *J. Chromatogr. A*, 1238: 1–10.
- [16] Goja, A. M.; Yang, H.; Cui, M.; Li, C. (2013) Aqueous two-phase extraction advances for bioseparation. *J. Bioprocess. Biotechnol.*, 4:1–8.
- [17] Dutra-Molino, J. V.; Viana Marques, D. A.; Pessoa Júnior, A.; Mazzola, G. P.; Gatt M. S. (2013) Different types of aqueous two-phase systems for biomolecule and bioparticle extraction and purification. *Biotechnol. Progr.*, 29(6): 1343–1353.
- [18] Dutra-Molino, J. V.; Araujo Feitosa, V.; de Lencastre-Novaes, L. C.; de Carvalho, V.; Moreni Lopes, A.; Faustino Jozala, A.; de Araújo, D.; Pellegrini Malpiedi, L.; Pessoa Júnior, A. (2014) Biomolecules extracted by ATPS: Practical examples. *Rev. Mex. Ing. Quim.* 13(2): 359–377.
- [19] Perumalsamy, T.; Murugesan, T. (2006) Prediction of liquid-liquid equilibria for PEG 2000–sodium citrate based aqueous two-phase systems. *Fluid Phase Eq.*, 244: 52–61.
- [20] Ames, L. L.; Robert, B. (1970) Phosphorous removal from effluents in alumina columns. *J. Water Pollut. Control Federation*, 42(5): R161–R172.
- [21] Batchelor, B.; Dennis, R. (1987) A surface complex model for adsorption of trace components from wastewater. *J. Water Pollut. Control Federation*, 59(12): 1059–1068
- [22] Zhao, D.; Sengupta, A. K. (1998) Ultimate removal of phosphate from waste water from using a new class of polymeric ion exchanges. *Water Res.*, 32(5): 1613–1625.
- [23] Kofinas, P.; Kioussis, D. R. (2003) Reactive phosphorous removal from aquaculture and poultry production systems using polymeric hydro gels. *Environ. Sci. Technol.*, 37(2): 423.
- [24] Vernau J, Kula, M. R. (1990) Extraction of proteins from biological raw materials using aqueous polyethyleneglycol-citrate phase systems. *Biotechnol. Appl. Biochem.* 12: 397–404.
- [25] Pellegrini Malpiedi, L. P.; Pico, G. A.; Loh, W.; Nerli, B. B. (2010) Role of polymer–protein interaction on partitioning pattern of bovine pancreatic trypsinogen and alphachymotrypsinogen in polyethyleneglycol/sodium tartrate aqueous two-phase systems. *J. Chromatogr. B*, 878: 1831–1836.
- [26] Duarte, A. W.; Dayo-Owoyemi, I.; Nobre, F. S.; Pagnocca, F. C.; Chaud, L. C; Pessoa, A.; Felipe, M. G.; Sette, L. D. (2013) Taxonomic assessment and enzymes production by yeasts isolated from marine and terrestrial Antarctic samples. *Extremophiles* 17(6): 1023–1035.
- [27] Pellegrini Malpiedi, L.; Fernandez, C; Picó, G; Nerli, B. (2008) Liquid–Liquid equilibrium phase diagrams of

- polyethyleneglycol + sodium tartrate + water two-phase systems. *J. Chem. Eng. Data*; 53: 1175–1178.
- [28] Tubio, G. P; Pellegrini, L.; Nerli, B. B.; Picó, G. A. (2006) Liquid-liquid equilibria of aqueous two-phase systems containing poly(ethyleneglycols) of different molecular weight and sodium citrate. *J. Chem. Eng. Data*, 51: 209–212.
- [29] Charney, J.; Tomarelli, R. M. (1974) Azocasein method. *J. Biol. Chem.*, 171: 501–505.
- [30] Smith, P. K.; Krohn, R. I.; Hermanson, G. T.; Mallia, A. K.; Gartner, F. H; Provenzano, M. D.; Fujimoto, E. K.; Goeke, N. M.; Olson, B. J.; Klenk, D. C. (1985) Measurement of protein using bicinchoninic acid. *Anal. Biochem.*, 150(1): 76–85.
- [31] Zhang, J. (2012) Protein-protein interactions in salt solutions. In *Protein-Protein Interactions: Computational and Experimental Tools*; Dr. Weibo Cai, ed., InTech: Croatia.
- [32] Annunziata, O.; Asherie, N; Lomakin, A.; Pande, J.; Ogun, O; Benedek, G. B. (2002) Effect of polyethylene glycol on the liquid-liquid phase transition in aqueous protein solutions. *Proc. Natl. Acad. Sci. USA*, 99 (22): 14165–14170.
- [33] Rocha, M. V.; Nerli, B. B. (2013) Molecular features determining different partitioning patterns of papain and bromelain in aqueous two-phase systems. *Int. J. Biol. Macromol.*, 61: 204–211.
- [34] Pellegrini Malpiedi, L.; Romanini, D.; Picó, G. A.; Nerli, B. B. (2009) Purification of trypsinogen from bovine pancreas by combining aqueous two-phase partitioning and precipitation with charged flexible chain polymers. *Sep. Purif. Technol.*, 65(1): 40–45.
- [35] Kendrick, B. S.; Li, T.; Chang, B. S. (2009) *Physical Stabilization of Proteins in Aqueous Solution*; Plenum Press: New York, pp. 61–84.
- [36] Evans, D. R.; Romero, J. K.; Westoby, M. (2009) Concentration of proteins and removal of solutes. *Methods Enzymol.*, 463: 97–120.
- [37] Marcos, J. C. F.; Ramalho, M. T.; Cabral, J. M. (1998) Variation of penicillin acylase partition coefficient with phase volume ratio in poly(ethylene glycol)-sodium citrate aqueous two-phase systems. *J. Chromatogr. B Biomed. Sci. Appl.* 711(1–2): 295–299.
- [38] Pellegrini Malpiedi, L. P.; Picó, G. A.; Nerli, B. B. (2011) Studies of protein partition in non conventional aqueous two-phase systems as method to purify trypsinogen and alpha-chymotrypsinogen from bovine pancreas. *Sep. Purif. Technol.*, 78: 91–96.
- [39] Huddleston, J. G. W.; Flanagan, J. A.; O'Brien, S.; Lyddiatt, A. (1994) Variation of protein partition coefficients with volume ratio in poly(ethylene glycol)-salt aqueous two-phase systems. *J. Chromatogr. A.*, 668: 3–11.39.
- [40] Bacha, E.; Sant'Anna, V.; Daroib, D. J.; Folmer Correa, A. P; Segalinc, J.; Brandellia, A. (2012) Production, one-step purification, and characterization of a keratinolytic protease from *Serratiamarcescens* P3. *Process Biochem.* 47: 2455–2462.
- [41] Porto, T. S.; Medeiros, G. M.; Porto, C. S.; Cavalcanti, M. T. H.; Neto, B. B.; Lima-Filho, J. L.; Converti, A.; Porto, A. L. D.; Pessoa, Jr, A. (2008) Liquid-liquid extraction of proteases from fermented broth by PEG/citrate aqueous two-phase system. *Chem. Eng. Process*, 47: 716–721.
- [42] Pericin, D. M.; Mađarev-Popović, S. Z.; Vaštag, Z. Q. (2008) Partitioning of acid protease from *Penicillium roqueforti* in aqueous two-phase system polyethylene glycol/phosphate. *Acta Periodica Technologica*, 39: 171–180.
- [43] Subathra, K.; Jeevitha, G. C.; Deepa, R. (2012) Aqueous two phase extraction of protease from neem leaves [*Azadirachta indica*]. *Int. J. Adv. Chem. Sci. Appl.*, 3: 346–351.
- [44] Chouyyok, W.; Wongmongkol, N.; Siwarungson, N.; Prichanont, S. (2005) Extraction of alkaline protease using an aqueous two-phase system from cell free *Bacillus subtilis* TISTR 25 fermentation broth. *Process Biochem.* 40: 3514–3518.
- [45] Kirsch, L.; dos Santos, A. C; Teixeira, M. F. S. (2012) Partition of proteases from *Lentinus citrinus* dpua 1535 by the peg/phosphate aqueous two-phase system. *Quim. Nova* 35(10): 1912–1915.
- [46] Loc, N. H.; Lien, H. T. T.; Giap, D. V.; Quang, H. T. (2013) Purification of recombinant neutral protease (NPRC10) by partitioning in aqueous two-phase systems. *Eur. J. Exp. Biol.*, 3(2): 252–257.
- [47] DivyaJyothi, M.; RohiniKiran, K.; Ravindhranath, K. (2012) Phosphate pollution control in waste waters using new bio-sorbents. *Int. J. Water Res. Environ. Eng.*, 4(4): 73–85.
- [48] Non-surfactant organic ingredients and zeolite based detergents, Final report prepared for the European Commission, RPA June 2006. Available from: http://ec.europa.eu/enterprise/sectors/chemicals/files/studies/rpa_non_surf_organ_zeolites_en.pdf
- [49] Lario, LD.; Chaud, L.; Almeida, MG.; Converti, A.; Durães Sette, L.; Pessoa-Junior, A. (2015) Production, purification and characterization of extracellular acid protease from the marine Antarctic yeast *Rhodotorula mucilaginosa* L7. *Fungal Biol.* (In press).