

# Palaeoenvironmental characterization of a crocodilian nesting site from the Late Cretaceous of Brazil and the evolution of crocodyliform nesting strategies



Júlio C. de A. Marsola<sup>a,\*</sup>, Alessandro Batezelli<sup>b</sup>, Felipe C. Montefeltro<sup>c</sup>, Gerald Grellet-Tinner<sup>d,e</sup>, Max C. Langer<sup>a</sup>

<sup>a</sup> Laboratório de Paleontologia de Ribeirão Preto, FFCLRP, Universidade de São Paulo, Avenida Bandeirantes 3900, Ribeirão Preto, São Paulo, 14040-901, Brazil

<sup>b</sup> Departamento de Geologia e Recursos Naturais, Instituto de Geociências, Universidade Estadual de Campinas, Rua João Pandiá Calógeras, 51, Campinas, São Paulo, 13083-870, Brazil

<sup>c</sup> Departamento de Biologia e Zootecnia, FEIS-UNESP, Rua Monção 226, 15385-000, Ilha Solteira, Brazil

<sup>d</sup> Orcas Island Museum, PO Box 134, 181 North Beach Road, Eastsound, WA 98245, United States

<sup>e</sup> Investigador Correspondiente at Departamento de Geociencias, CRILAR, CONICET, Argentina

## ARTICLE INFO

### Article history:

Received 19 February 2016

Received in revised form 9 June 2016

Accepted 10 June 2016

Available online 11 June 2016

### Keywords:

*Pissarrachampsia sera*

Crocodylian fossil eggs

Palaeoenvironment

Adamantina Formation

Late Cretaceous

Bauru Basin

## ABSTRACT

Despite the vast crocodyliform fossil record, little is known about the reproductive biology and nesting strategies of the extinct members of the group. Here we report a large accumulation of crocodilian fossil eggs from the type-locality of the baurusuchid *Pissarrachampsia sera*. Sedimentary facies and architectural elements of the site support a palaeoenvironmental model with a shallow lacustrine, playa-lake system interacting to ephemeral braided fluvial channels, with aeolian influence and development of sandy soils. The presence of pedogenic calcretes in the palaeosols indicates arid to semi-arid conditions. The crocodilian affinity of the eggs is supported by the thin eggshell which bears wedge-shaped shell units with tabular microstructures. Furthermore, taphonomic data support an autochthonous assemblage of eggs and skeletal remains, hinting at a monotypical stratigraphic horizon and suggesting *P. sera* as the egg-laying taxon. The repeated pattern of four (eventually five) eggs per clutch at the site demonstrates that *P. sera* laid fewer eggs compared to modern crocodilians, indicating that k-selected reproductive strategy pattern is pervasive in the fossil record of Notosuchia. In the crocodyliform phylogenetic framework, the k-strategy and the “egg clutch sizes” optimization of Notosuchia is opposite to the strategy with larger clutches consistently occurring in modern Crocodylia and Neosuchia, the sister clade to Notosuchia. Yet, the lack of data on more early-branching taxa renders unclear which pattern is plesiomorphic for Crocodyliformes as a whole.

© 2016 Elsevier B.V. All rights reserved.

## 1. Introduction

Crocodyliformes achieved high taxonomic diversity through geologic time (Bronzati et al., 2015) and were notably represented in Gondwanan landmasses by the Notosuchia (Pol et al., 2014; Fiorelli et al., 2015), a successful and diverse clade of terrestrial forms (Martinelli and Teixeira, 2015). The Late Cretaceous Brazilian fossil record, particularly that of the Bauru Group, depicts an important episode in notosuchian evolutionary history, when they played major ecological roles, occupying different niches, from first to second level consumers to top predators with complex social behaviour (Avilla et al., 2004; Godoy et al., 2014).

Although crocodilian fossil eggs have been reported worldwide in the past thirty years (Moreno-Azanza et al., 2014, Table 1), their associations with skeletal remains are scarce. The best documented fossil assemblage is reported from the Late Cretaceous notosuchid *Yacarerani boliviensis* (Novas et al., 2009) in Bolivia, and similar associations are known from the Late Jurassic of Portugal (Russo et al., 2014) and the Middle Eocene of Germany (Hastings and Hellmund, 2015), providing insights on their reproductive biology. Moreover, several sites of the Bauru Group are known for the association of fossil eggs and notosuchid skeletal remains (Magalhães Ribeiro et al., 2006; Vasconcellos and Carvalho, 2010; Marsola et al., 2011; Oliveira et al., 2011). However, in contrast to Mesozoic non-avian and avian dinosaurs (e.g. Chiappe et al., 2005; Grellet-Tinner et al., 2006; Grellet-Tinner and Fiorelli, 2010; Fernández et al., 2013; Hechenleitner et al., 2015), combined taxonomic and stratigraphic studies of crocodyliforms nesting sites and strategies are lacking.

Here we report a large accumulation of crocodile fossil eggs, including clustered and isolated elements, from the type-locality of

\* Corresponding author.

E-mail addresses: [juliomarsola@gmail.com](mailto:juliomarsola@gmail.com) (J.C.A. Marsola), [abatezelli@ige.unicamp.br](mailto:abatezelli@ige.unicamp.br) (A. Batezelli), [fcmontefeltro@bio.feis.unesp.br](mailto:fcmontefeltro@bio.feis.unesp.br) (F.C. Montefeltro), [locarnolugano@gmail.com](mailto:locarnolugano@gmail.com) (G. Grellet-Tinner), [mclanger@ffclrp.usp.br](mailto:mclanger@ffclrp.usp.br) (M.C. Langer).

the baurusuchid *Pissarrachampsia sera* (Montefeltro et al., 2011), in the Late Cretaceous Adamantina Formation, Bauru Basin, SE Brazil. This study focuses on (1) describe the depositional and pedological context of the fossil-bearing strata, in an attempt to understand the palaeoenvironment; (2) comparatively describe the oological remains; (3) infer the egg-laying taxon; and (4) discuss the possible implications of the gathered data for understanding the evolution Crocodyliformes nesting strategies.

## 2. Material and methods

### 2.1. Study area and sampling

This study mainly relies on facies and pedological analyses of outcrops, along with laboratory analyses. The descriptive and photographic documentation of the outcrop where the fossils were found (Fig. 1) – Inhaúmas-Arantes farm, Campina Verde, Minas Gerais, Brazil (19°20' 41.8" S; 49°55' 12.9" W) – and the elaboration of columnar sections based on field data provided the basis for the identification of the sedimentary processes and pedogenic features. Facies characterization combining field work, petrography, scanning electronic microscopy (SEM), and X-Ray diffractometric analyses were performed, focusing on the texture, colour, sedimentary structures, and mineralogy, to interpret depositional and diagenetic processes. Following Retallack (2001),

pedological interpretations are based on macro and micromorphology analyses of the palaeosols.

### 2.2. Material

The studied specimens are housed at Laboratório de Paleontologia, Universidade de São Paulo, Ribeirão Preto campus, and include three clutches with four eggs each (LPRP-USP 0046, 0047, 0048) (Fig. 7), one clutch with at least four compressed eggs (LPRP-USP 0648) (Fig. 7), isolated eggs (LPRP-USP 0638, 0640, 0647) (Fig. 8), and eggshells (LPRP-USP 0641, 0639) (Fig. 8). The eggshell samples were thin sectioned and submitted to LEICA DM4000B optical microscopy (OM) for microcharacterization of the eggshell units, including their general morphology and geometry. In order to generate tridimensional images of the shell ultrastructure, SEM analyses were performed in JEOL JSM 6460 and FEI Quanta 650 FEG.

### 3. Geological settings and palaeoenvironmental characterization

The above 300,000 km<sup>2</sup> Bauru Basin covers parts of the states of Minas Gerais, Goiás, Mato Grosso do Sul, São Paulo, and Paraná (Fig. 1). This basin was filled with 300 m of continental deposits, divided into the Caiuá and Bauru groups (Fig. 2). The origins of the Bauru Basin is related to subsidence by isostatic adjustment occurred South of the Alto




Fig. 1. Surface distribution map of the Bauru Basin lithostratigraphic units. (Edited from Batezelli, 2015.)

Paraná Uplift (SAP; Fig. 1) during the Late Cretaceous in South-eastern Brazil (Batezelli, 2015).

The fossils focus of this research were discovered on the top of the Adamantina Formation (Soares et al., 1980) near its contact with the Marília Formation, in western Minas Gerais. This unit crops-out in the central and western parts of the Triângulo Mineiro region, and is composed of reddish, fine- to medium-grained and moderately-sorted sandstones. These are either massive or exhibit small to medium scale cross-stratifications (up to 3 m-sized foresets), planar stratifications characterized by ripple marks and desiccation cracks (Batezelli, 2010). In the Gurinhata and Campina Verde areas, the Adamantina Formation also includes well-sorted and fine-grained sandstones, forming large scale (more than 3 m) and high angle (about 25°) cross-stratification foresets. Reddish mudstones, massive or with ripple marks and planar stratifications, often intercalated by centimetric to decimetric layers of very fine-grained sandstones, cemented or not by calcium carbonate, may also occur (Fig. 3). The sandstone layers are lenticular or tabular, sometimes intercalated with centimetric mudstone layers. Trough cross-stratifications and carbonatic cement are common, and bioturbations (krotovinas and root marks) are also frequent, tunnels are mostly filled by well-sorted, very fine- to fine-grained sandstone, with low angle cross-bedding (Fig. 3). The geometry of the bodies, the fine texture, and the sedimentary structures of the Adamantina Formation, indicate fluvial flows deposition, with the development of muddy plains. Small aeolian dunes of the sand-sheet type, adjacent to the fluvial deposits, interacted with them during dry periods, partially filling abandoned channels.


A typical trait of the Adamantina Formation is the presence of palaeosols, which profiles occur intercalated in the deposits and are characterized by calcretes. These paleosols allow a general comparison to the present soil orders Alfisol, Aridisol, Entisol, and Vertisol,

indicating arid and semi-arid climatic conditions (Dal' Bó et al., 2010; Pereira et al., 2015).

### 3.1. Depositional environment of the Adamantina Formation


The about 2 m thick and 70 m long fossil-bearing outcrop (Fig. 2) is composed of massive muddy sandstones, fine to medium grained, moderately sorted, and with carbonatic cement. At the base, reddish mudstone and sandstone layers (about 7 cm thick) are intercalated on planar-parallel bedding (Fig. 3). These are characterized by fining-upward rhythmic cycles of up to 40 cm thick, with an erosive-abrupt base. Upwards in the section, the sandstones are mostly massive, but planar stratification and cross beddings are also common. The sandstones are poorly to moderately sorted and the quartz and feldspar grains are rounded to subangular. Rhythmites disappear in the top of the section, sandstones and mudstones are mixed up, and dish structures and convoluted folds are present. From the middle to the upper portions of the section, the muddy sandstone turns into massive sandstones (palaeosols), with carbonatic nodules, root marks, and ped-like structures (Fig. 4). The Marília Formation appears locally above the described lithology, where it is composed by reddish to yellowish, fine to medium grained, and poorly to moderately sorted sandstones. These sandstones are be muddy, massive, cemented by calcium carbonate, and occurs associated with centimetric to decimetric layers of fining-upward conglomerates, composed by rounded to sub-angular pebbles of quartz and quartzite.

In the fossil-bearing site, the rhythmic deposits of sand- and mudstones of the Adamantina Formation indicate episodic sedimentary processes, with slow decrease in energy, as typical of those formed by low viscosity turbidity flows generated during flooding periods on depressed areas ("inundites" sensu Della Favera, 1984). The massive,



**Fig. 2.** Chronostratigraphic chart of the Cretaceous sequences of the Paraná/Bauru basins, based on Amaral et al., 1967 (sample CSN); Hasui and Cordani, 1968 (samples AX, C-3, S-10, S-31, A-2-4c, OB-SN, SB, S-1, P, T-2, B-1); Sonoki and Garda, 1988 (samples CT, CS, CCI); Machado Júnior, 1992 (sample CCII); Guimarães et al., 2012 and Fragozo et al., 2013 (Pterosaurs); Gobbo-Rodrigues, 2001 and Dias-Brito et al., 2001 (Ostracods); Santucci and Bertini, 2001 and Martinelli et al., 2011 (Aelosaurus). (Edited from Batezelli, 2015.)






**Fig. 3.** Adamantina Formation. A, Sand bars deposits with trough cross bed, interlayered with mudstone (Prata-Campina Verde road, 20 km from Prata-MG). B, Very fine sandstone with low-angle trough cross bed. C, Channel deposits (BR 364 road, 5 km from Gurinhatã-MG). D, Tabular sand bars interlayered with mudstones (BR 364 road, 5 km from Gurinhatã-MG).

muddy sandstones that appear above the rhythmic facies show root marks and carbonatic nodules, both of which indicate pedogenic processes. The superimposed Marília Formation is formed by high-energy fluvial deposits. This includes channel-geometry conglomerates and sandstones, as well as downstream and lateral accretion macroforms. The depositional geometry and prevalence of bottom sediments suggests a braided river system, whereas the occurrence of thick palaeosol horizons indicate that the sediment-transporting flows were ephemeral. Indeed, the thick massive conglomerates are typical of

density-detrital flows associated to rapid, high-energy events (Ramos and Sopena, 1983; Miall, 1985, 1988a, 1988b, 1996).

Palaeocurrent data (cross stratification front measurements) from the Campina Verde area (Batezelli, 2003) indicate a prevalent east/northeast to west/southwest sedimentary transport, with base level positioned at the central/south part of the Basin. The high variability of the paleocurrents, together with the identified facies and architectural elements, shows an alluvial-fan system, dominated by low-sinuosity braided rivers, as discussed by Stanistreet and McCarthy



**Fig. 4.** Adamantina Formation. A, Calcrete overlying channel deposits. B, Carbonate nodules in palaeosol profile. C, Root mark. D, Prismatic ped structure in palaeosol.






Fig. 5. Depositional model to the study area.

(1993), or a distributive fluvial system according to Weissmann et al. (2010). In general, architectural elements of the Bauru Group resemble those described by Bromley (1991) from the Kayenta Formation, Colorado, and by Jones et al. (2001) from the Rio Vero Formation, Spanish Pyrenees, both related to intermediate to distal alluvia-fan systems (Batezelli, 2003). The proposed depositional model also resembles that of the Huesca alluvial system, Ebro Basin (Spain), where the braided fluvial channels are distributaries, evolving from alluvial-fans that enter the basin from the Pyrenees (Hirst, 1991). A general palaeoenvironment model for the area would include a shallow lacustrine environment, such as a playa-lake, acting as the base level and interacting with ephemeral braided fluvial system, with aeolian influence and the development of sandy soils (Fig. 5). Accordingly, Batezelli (2015) states that the sedimentary characteristics of the Adamantina Formation indicate a playa-lake setting with high sedimentation rate, dominated by suspension settlings in the more internal portions. Finally, density flows generated during floods may be responsible for the upward fining rhythmic deposits, i.e., Tb-d Bouma's sequence turbidity (Bouma, 1962) or "inundites" (Della Favera, 1984). Low-density mudflows were responsible for the muddy deposits. Carbonate cementation and vegetation development would be responsible for the obliteration of sedimentary structures. Fauna and flora (palynomorphs, ostracods, conchostraca, and charophyta) show that the environment had long-lasting waters (Gobbo-Rodrigues, 2001; Dias-Brito et al., 2001).

### 3.2. Paleosols and paleoenvironments consideration

The most striking feature of the fossil-bearing outcrop is the occurrence of paleosols. The profiles have undeveloped "B" (commonly lacking) and "C" horizons, high carbonate content, and colours ranging from 7.5 R to 10 R (Fig. 4). The horizons display light colours, carbonate nodules, medium to large prismatic structures, and have sandy-clay textures, with millimetre- to centimetre-sized clasts and abundant root marks (Fig. 4). The root marks range from 0.3 to 90 cm in length, are tubular in shape and filled with  $\text{CaCO}_3$  or clay (Fig. 4B). Quartz occurs at 50% levels and corresponds to the main detrital phase in the rock. Both poli- and monocrystalline forms are present, with undulated extinction. Calcite substitution is common, generating massive calcrete




Fig. 6. A, X-ray diffractogram showing minerals and clay minerals. B, Photomicrography: micritic cement, authigenic palygorskite (Pa) and silex (Si) (from Batezelli, 2015). C, Detail of the fibrous aggregates of the palygorskite (image obtained by SEM). D, Dolomite and hematite cement.

layers (Fig. 4 and 6A). Illite, montmorillonite, and palygorskite are the most common clay minerals. Palygorskite occurs in intertwining fibrous aggregates forming nodular structures (Fig. 6B).

### 3.3. Pedogenic considerations

The presence of palaeosols in the Bauru Groups is ubiquitous (Suguio et al., 1975; Suguio and Barcelos, 1983; Dal' Bó et al., 2009; Pereira et al., 2015). The described section displays root marks, blocks structures, and is heavily cemented by  $\text{CaCO}_3$ , indicating pedogenic processes with calcrete formation (Watson, 1992; Klappa, 1980; Wright, 2007). Pedogenic calcretes are formed in shallow soils, above the phreatic level, and are variably developed (Alonso-Zarza, 2003).

This allows for the thick calcrete profiles of the site, which may also result from the superposition and repetition of such environments over time. The presence of calcretes is a clear indication of arid to semi-arid conditions, because they are formed by soil accumulation of  $\text{CaCO}_3$  due to intense evapotranspiration and changes in the  $\text{CO}_2$  partial pressure.


Nowadays, pedogenic calcretes occur as  $\text{CaCO}_3$  accumulations near or at the surface; are uniform and continuous over several kilometres; are typical of semi-arid and temperate climates; form on low declivity (below  $25^\circ$ ) areas; are substrate independent and may occur in any rock type; are impoverished in  $\text{CaCO}_3$  towards the base of the horizons; and rarely overcome six metres thick (Klappa, 1983). Soil water may have the  $\text{Mg}^{2+}/\text{Ca}^{2+}$  relation increased under severe evapotranspiration, allowing for authigenic precipitation of dolomite, as described in the Coorong lagoon, near Adelaide, Australia (von Der Borch, 1965). During hot summers, the  $\text{Mg}^{2+}/\text{Ca}^{2+}$  ratio may surpass 10.0, leading to dolomite precipitation. The studied samples bear pedogenic dolomites related to calcrete accumulation, with magnesium source related to palagonitization processes and/or alteration of mafic rock fragments from the northeast border of Bauru Basin (Fig. 1) (Batezelli, 2003). The presence of palygorskite has been associated with arid conditions, because its formation occurs under high Mg and Si concentrations in high pH waters. According to Batezelli (2015), the palaeosol profiles of the Bauru Basin indicate of arid to semi-arid conditions, with a maximum rainfall interval between 500 and 1000 mm (Jenny, 1941; Wright and Tucker, 1991; Retallack, 2005).

### 4. Egg description and comparisons

The three clutches LPRP/USP 0046–0048 are composed by four eggs each, horizontally distributed in the same plane with no overlapping of elements (Fig. 7). Specimens in the clutches are overall well preserved, but LPRP-USP 0048 was heavily weathered due to water percolation, precluding the analysis of its morphology beyond general shape and measurements. A fourth clutch, LPRP/USP 0648, preserves at least four vertically collapsed eggs also arranged in the same plane, with the possible preservation of a fifth egg (Fig. 7). The larger eggs are clustered in LPRP-USP 0047, measuring in average 7.2 by 4.2 cm. Smaller eggs are also present, as seen in the LPRP-USP 0638, a nearly complete isolated egg which is 5.9 by 3.17 cm in size (Table 1, Fig. 8). The LPRP-USP 0047 eggs are much larger than other crocodyliform eggs described for the Bauru Group, including *Bauruoolithus fragilis* (Oliveira et al., 2011) and those possibly associated to *Mariliasuchus amarali* (Magalhães Ribeiro et al., 2006). Indeed, they are one of the largest


**Table 1**  
Preservation status and measurements of *Pissarrachampsia sera* eggs.

Specimen		Preservation status	Egg length (cm)	Egg width (cm)
LPRP-USP 0046	1	Nearly complete eggs	5.94	3.92
	2		6.46	2.94
	3	Weathered eggs	6.78	3.90
	4		6.26	3.59
LPRP-USP 0047	1	Nearly complete eggs	7.24	4.21
	2		7.26	4.20
	3		7.02	4.16
	4	Partial egg	4.90	3.00
LPRP-USP 0048	1	Weathered eggs	5.65	4.81
	2		5.80	4.82
	3		6.88	4.74
	4		6.34	4.61
LPRP-USP 0638	Single egg	Nearly complete egg	5.90	3.17
LPRP-USP 0640	Single egg	Partial, compressed egg	6.26	3.92
LPRP-USP 0647	Single egg	Egg with damaged pole	4.66	3.40
LPRP-USP 0648	1	Partial, compressed eggs	5.92	3.90
	2		5.62	3.64
	3		6.90	4.07
	4		5.57	4.21



**Fig. 7.** *Pissarrachampsia sera* egg clutches LPRP/USP 0046 (A), 0047 (B), 0048 (C) and 0648 (D). In D, dashed lines indicate the possible limits of the collapsed eggs.





**Fig. 8.** *Pissarrachampsia sera* isolated eggs LPRP-USP 0638 (A), LPRP-USP 0640 (B), LPRP-USP 0647 (C) and eggshells LPRP-USP 0639 (D) and LPRP-USP 0641 (E).

known crocodile eggs, both fossil and extant (Marzola et al., 2015, Table 3). In general, the sampled eggs are elongated with blunt ends (Fig. 7), presenting the common elliptical shape to crocodyliform eggs (Hirsch, 1985; Kohring and Hirsch, 1996; Marzola et al., 2015). Likewise, the outer surface appears smooth in macroscopic observation, as seen in other crocodyliform eggs (Ferguson, 1982, 1985; Hirsch, 1985; Marzola et al., 2015). However, stereoscopy shows a complex slightly rugose/undulating surface, as in *B. fragilis* (Oliveira et al., 2011) and in *M. amarali* eggs (Magalhães Ribeiro et al., 2006), also resembling the rugosocavate condition observed in *Crocodylus mindorensis* and *Paleosuchus palpebrosus* eggs (Marzola et al., 2015). No extrinsic degradation pits are seen in the samples, but eggshell thickness is much reduced, although some size-related variation occurs; i.e. LPRP-USP 0047 eggshells are about 150  $\mu\text{m}$  thick, whereas smaller eggs (LPRP-USP 0046, 0638, 0640 and 0647) have thinner eggshells, ranging from 130 to 150  $\mu\text{m}$  (Fig. 9). Putative *M. amarali* eggs, which are much smaller, have eggshells about twice thicker (Magalhães Ribeiro et al., 2006), and are closer in thickness to *B. fragilis*, which range from 150 to 250  $\mu\text{m}$  thick (Oliveira et al., 2011). Crocodyliform eggshells about five times thicker than those dealt with here have been reported, reaching up to 750  $\mu\text{m}$  thick (Panadès I Blas and Patnaik, 2009; Moreno-Azanza et al., 2014). The here newly described eggs are among the thinnest known for crocodyliforms, considering both fossil and extant records (Marzola et al., 2015).


The ultrastructure of the described eggshells is not easily accessed due to diagenetic alterations (weathering and dissolution during fossilization). Fully interlocked shell units are recognized, occasionally intercalated by circular pore openings, similar to those of *Paleosuchus palpebrosus* eggs (Marzola et al., 2015). Pore openings are approximate 40  $\mu\text{m}$  in diameter, visible only in LPRP-USP 0046 and 0047, with average density difficult to establish due to their sparseness (Fig. 9). *Bauruoolithus fragilis* shows pore openings with diameter from 30 to 80  $\mu\text{m}$  (Oliveira et al., 2011), which depart from the pattern described here for their elliptical to subtriangular shape and more abundant distribution. In general, pore openings are evidently more abundant in eggs of living crocodyliforms, such as in *Crocodylus mindorensis* and *Paleosuchus palpebrosus* (Marzola et al., 2015), than in fossil eggs, as those described here.

The shell units exhibit an irregular morphology, with a trapezoid or wedge-shaped arrangement (Fig. 9), usually regarded as a typical crocodyliform feature (Hirsch, 1985; Kohring and Hirsch, 1996; Rogers, 2001). The shell unit height/width ratio ranges from 1.2 to 1.5. However, shell units twice higher than wide or even wider than high are also observed. This height/width ratio resembles that observed for shell units of *B. fragilis* (Oliveira et al., 2011). In addition, wedges are observed in the structure of the shell units in cross section (Fig. 9). The shell inner layer is composed of circular shell units, about 40 to 50  $\mu\text{m}$  in diameter. Only LPRP-USP 0047 preserves basal knobs and surrounding “petal-like” structures (Fig. 9), which resemble those of *Krokolithes helleri* (Kohring and Hirsch, 1996). There is no clear boundary between the middle layer and the outer layer, hampering the precise measurement of their thickness. The typical crocodyliform tabular structures (Ferguson, 1985; Hirsch, 1985; Kohring and Hirsch, 1996; Moreno-Azanza et al., 2014; Russo et al., 2014) are not frequently found in the eggshell ultrastructure, most probably due to the preservation biases. When present (Fig. 9D), however, they are equally spaced, both in the middle and in outer layers, differently of the observed by Moreno-Azanza et al. (2014) in *Krokolithidae* eggshells from the Late Cretaceous of Spain.

## 5. Discussion

### 5.1. Egg-layer, taphonomic and environmental considerations

Fossil eggs and nests give unique clues to the reproductive biology of extinct organisms and are usually related to exceptional preservation



**Fig. 9.** A, OM of LPRP/USP 0047 eggshell cross section showing the general irregular and trapezoid aspect of the shell units and the presence of wedges (white arrow). B, schematic drawing of the shell units seen in A, depicting the height/width variation of the shell units with the presence of both wider (dark grey) and higher (lighter grey) units in the same eggshell. C and D, MEV of the LPRP/USP 0046 and 0047, respectively. C, inner view of the shell units showing their rounded aspect and the presence of “petal-like” structures at the base (black frame). D, eggshell in cross section detailing the occurrence of tabular structures in its arrangement. E, rounded pore opening indicated by the black arrow.

contexts (i.e., Norell et al., 1995; Chiappe et al., 1998; Mateus et al., 1998; Schweitzer et al., 2002; Araújo et al., 2013; Ribeiro et al., 2014). Indeed, eggs are usually fragile and prompt to fragmentation during transport (Hayward et al., 2011). In the type-locality of *Pissarrachampsia sera*, all eggs were preserved without biogenic patterns, such as hatching, predation, or trampling signs. LPRP-USP 0639 and 0641 are more fragmentary (Fig. 8), but they represent leftovers of a single collapsed egg, as in the case of the clutch LPRP-USP 0648. Because of the fact that the eggshell fragments still overlap one another, we exclude any transportation of the specimens during sedimentation. As pointed out by Hayward et al. (2000, 2011), if the eggs were subject to any sort of directional transport prior to sedimentation, eggshell fragmentation would be more conspicuous and the fragments may result organized with an orientation pattern. Additionally, Oser and Jackson (2014) show that the edges of eggshell fragments acquire abrasion signs when interact with the sediment during transport, and such features are absent from LPRP-USP 0637, 0639, and 0648. Absence of transport can also be inferred for the three better preserved clutches, LPRP-USP 0046–0048.

Although the described eggs vary in their overall dimension (Table 1), this size disparity cannot be considered a proxy for taxonomic diversity, considering that the average egg size in the extant *Alligator mississippiensis* is directly proportional to the age of the females (with a variation up to 2 cm) (Ferguson (1985). The eggs of the *Pissarrachampsia sera* type-locality are collected from the same stratigraphic horizon preserving several well-preserved skeletal remains of this crocodyliform, including skulls and articulated postcranial elements of different ontogenetic stages (Godoy et al., 2016). This suggests an autochthonous assembly, as corroborated by the evidences disfavouring

a long range transport of the eggs and the rhythmic deposits that indicate episodic sedimentary processes with slow decrease in energy and low viscosity turbiditic flows. Though an egg-taxon association is only certain in cases of extraordinary preservation and conditions, such as the presence of embryos and pregnant females (Chiappe et al., 1998; Mateus et al., 1998; Zhou and Zhang, 2004; Sato et al., 2005; Grellet-Tinner et al., 2006; Araújo et al., 2013) or in clear cases of parental care (Norell et al., 1995), the studied locality appears to include a monotypic egg-taxon assemblage that allows inferring *P. sera* as the best potential egg-layer, alike the association egg-taxon between *Bauruolithus fragilis* and *Baurusuchus pachecoi* described in Oliveira et al. (2011).

The eggs attributed to *Pissarrachampsia sera* are found in deposits that suggest a sedimentary dynamic characterized by short flood periods, intercalated by long periods of subaerial exposure. During these latter periods deprived of sedimentation, deposits were exposed to weathering factors responsible for the pedogenesis. Although long-lasting water was present in the system, the general climatic pattern was surely arid to semi-arid. Together with the fact that *P. sera* eggs were found in sandy paleosols, the geological data suggests that this crocodilian nested in dryer environments compared to extant forms, that nest in areas near water bodies and take advantage of surrounding humidity to incubate their eggs (Ferguson, 1981; Ferguson, 1985). A proposed relation between total pore area, eggshell thickness, and water vapour conductance, indicates that, in birds, thicker and more porous eggs lose more water than thinner and less porous ones (Ar et al., 1974). Plus, archosaur hardly mineralized eggshells minimize the loss of water, strongly unlinking the embryo from the hydric potential of the surrounding environment (Packard et al., 1982). Tanaka et al.






Fig. 10. Simplified phylogenetic framework mapping egg-clutch sizes across Mesoeucrocodylia evolution.

(2015) found covered nesters, such as crocodilians and megapods, to possess more porous eggs than open nesters birds. Accordingly, Seymour (1979) observed that the high porosity of the eggs of some dinosaurs have a clear association with their incubation in nests with quite humid conditions. Accordingly, the scarcity of pores and the thin shells of the eggs described here could also be interpreted as an adaptation for dryer environments, reducing water loss during the embryo development. Therefore, modifications towards a more terrestrial way of life may not be restricted to the skeletal anatomy of baurusuchids (Riff and Kellner, 2011; Godoy et al., 2016), but also include aspects of their reproductive behaviour.

### 5.2. Evolutionary significance of *Pissarrachampsia sera* egg

The above mentioned attribution of the egg clutches to *Pissarrachampsia sera*, along with the previous records of such fossils, provides a unique opportunity to discuss the reproductive biology of baurusuchids and its implications for understanding the origin of the reproductive biology of living crocodyliiforms. The small number of eggs (four to five maximum) per clutch in *P. sera* differs remarkably from the pattern seen in living crocodilians, ranging from a lower

limit of 10 eggs per clutch in *Alligator sinensis*, *Osteolaemus tetraspis*, *Paleosuchus palpebrosus*, and *Crocodylus mindorensis*, to up to 80 eggs in clutches of *Crocodylus niloticus* (Greer, 1975; Platt et al., 2004; see Brazaitis and Watanabe, 2011 for a review). Offspring number is a key factor in determining the life-history strategy of species, but the broadly used k- and r-selected dichotomy is hard to fit in modern crocodilians (Magnusson, 1986; Tucker, 2001; Abercrombie et al., 2001; Grigg and Kirshner, 2015). In one aspect, the group has a greater number of general k-strategists traits (e.g. large size, long-living individuals, slow grow, and long reproductive life), but also presents some typical characteristics of r-selected species, like the numerous eggs per clutch resulting in large offspring (Abercrombie et al., 2001; Tucker, 2001; Grigg and Kirshner, 2015). As for *P. sera*, although much information is lacking to define its life-history strategy, the small number of eggs per clutch pushes the taxon to the k-selected end of the spectrum represented by modern forms (Grigg and Kirshner, 2015).

Considering the entire documentation of crocodyliiform egg clutches, the four-egg clutch of *Pissarrachampsia sera* does not depart so much from the pattern seen in the fossil record. All but two Mesozoic crocodyliiform egg clutches present a lower number of eggs per clutch than the lowest limit observed in modern crocodilians (Greer, 1975; Brazaitis and Watanabe, 2011); moreover, those with the lowest number of eggs are all from the Cretaceous of South America: two-egg clutches were putatively referred to *Mariliasuchus amarali*, also from the Bauru Group (Magalhães Ribeiro et al., 2006), and to *Yacarerani boliviensis*, from the Late Cretaceous Cajones Formation of Bolivia (Novas et al., 2009). The nesting site reported by Oliveira et al. (2011) from the Adamantina Formation includes one clutch of four eggs (see also Oliveira, 2008). The ootaxon *Bauruoolithus fragilis* was named to encompass these eggs, possibly associated to *Baurusuchus pacheco* (Oliveira et al., 2011). Two other Adamantina Formation crocodyliiform clutches, with four and six eggs, were preliminarily described by Magalhães Ribeiro and Arruda Campos (2009), with no taxonomic or parataxonomic assignment. Finally, Carvalho et al. (2013) reported a presumable adult specimen of *Mariliasuchus amarali* still from the Bauru Group and found in close association to four egg clutches formed by 3–4 eggs each. Thus, the only exceptions to the rule of reduced egg clutches in fossil crocodyliiforms are so far from the Late Jurassic of Portugal (Russo et al., 2014; Araújo et al., 2015) and from the Late Cretaceous of India (Srivastava et al., 2015), respectively assigned to Atoposauridae and Dyrosauridae. With 13 eggs each, they represent the largest crocodyliiform egg clutches known in the fossil record.

It is possible to identify, therefore, a clear discrepancy between the number of eggs in fossil and living crocodyliiform clutches; a pattern that may influence our understanding of the life-history strategies of the group through geological time. In order to further explore this subject, crocodyliiform egg clutch sizes were analysed in a phylogenetic framework (Fig. 10). Baurusuchidae (sensu Montefeltro et al., 2011) is a monophyletic group within Notosuchia (Fig. 10), including both *Pissarrachampsia sera* and *Baurusuchus pacheco* (associated to the oospecies *Bauruoolithus fragilis*). Likewise, *Mariliasuchus amarali* and *Yacarerani boliviensis* are part of the advanced notosuchians clade of Pol et al. (2014). On the other side, Atoposauridae and Dyrosauridae are members of Neosuchia, a broader clade that also includes modern crocodiles (Montefeltro et al., 2013; Pol et al., 2012, 2014; Rabi and Sebök, 2015; Leardi et al., 2015). Notosuchia and Neosuchia are consistently found as sister clades, forming one of the better established dichotomies in crocodyliiform phylogeny (Montefeltro et al., 2013; Pol et al., 2012, 2014; Rabi and Sebök, 2015; Leardi et al., 2015). Accordingly, the optimization of egg clutch sizes in such phylogenetic framework suggests a reduced number of eggs as the plesiomorphic condition at Notosuchia node, and a greater number of eggs as plesiomorphic condition at the Neosuchia node (Fig. 10).

The absence of data on the clutch size for earlier crocodyliiform splits precludes identifying confidently the ancestral condition for the Notosuchia/Neosuchia dichotomy. At this point, it is unknown if the

great number of egg per clutch of the modern crocodilians is a modification from a condition of smaller egg clutches present in notosuchians. Alternatively, the smaller egg clutches in notosuchians might in fact be a modification from a plesiomorphic condition of larger clutches present in neosuchians. The larger number of eggs per clutch of other archosaur, such as dinosaurs (i.e., Horner and Makela, 1979; Norell et al., 1995; Chiappe et al., 1998; Grellet-Tinner et al., 2012), might support the second scenario. However, given the gap in the knowledge about the egg-clutch size of more basal pseudosuchians, we prefer to take a more conservative approach and do not make this leap. New finds, such as 'protosuchian', basal notosuchian and basal pseudosuchian egg clutches, might help defining the ancestral Mesoeucrocodylia offspring size, shedding light onto origin of the unique reproductive biology of modern crocodilians.

## 6. Conclusions

Combined stratigraphic and taxonomic studies provided critical information for understanding nesting strategies of archosaurs as a whole. A previously unreported scenario is presented here, where articulated and isolated remains of the baurusuchid *Pissarrachampsa sera* are found in association to several egg clutches. The main conclusions drawn from the palaeoenvironmental characterization of the site and comparative study of the oological remains on a phylogenetic framework are summarized below:

- The rhythmic deposits of sand- and mudstones of the Adamantina Formation in the site indicate episodic sedimentary processes, with slow energy decrease, as typical of low viscosity turbidity flows generated during flooding periods on depressed areas. The muddy sandstones that appear above the rhythmic facies show root marks and carbonatic nodules, both of which indicate pedogenic processes.
- The pedogenic calcretes in the palaeosols are a clear indication of arid to semi-arid climatic conditions, further corroborated by the presence of palygorskite. As a whole, the paleoenvironment is interpreted as a shallow lacustrine system, such as a playa-lake, interacting with an ephemeral braided fluvial system, with aeolian influence and development of sandy soils.
- The combined presence of thin eggshell and wedge-shaped shell units with tabular structures supports the crocodilian affinity of the described eggs. The general egg morphology resembles that of *Bauruoolithus fragilis*, an oospecies assigned to *Baurusuchus*. However, eggs associated to that oospecies are smaller and probably more porous, which along with the elliptical to subtriangular shape of the pore openings, compose their main differences to *Pissarrachampsa sera* eggs.
- Along with the sedimentary patterns discussed above, the scarcity of scattered eggshells and the lack of abrasion signs in their edges, the good preservation of the eggs in clusters, and the presence of articulated remains of *Pissarrachampsa sera*, preclude a long range transport of the remains, indicating an autochthonous assembly. Also, such monotypic deposit allows inferring *P. sera* as the egg-laying taxon.
- The recurring pattern of four (possibly five) eggs per clutch suggest that *Pissarrachampsa sera* laid fewer eggs if compared to modern crocodilians, placing the taxon in the far end of the k-selected spectrum represented in Crocodyliformes.
- The apparent pattern of small egg clutches in notosuchians suggests a peculiar reproductive strategy for this group among crocodyliforms. The optimization of "egg clutch size" in the current phylogenetic framework, however, defines ambiguously if this pattern was already present or not in earlier splits of Crocodyliformes.

## Acknowledgments

The authors thank to Juliana M. Sayão, João A. Souza Neto and Edvaldo J. P. Santos for their help with SEM analyses at CTG – UFPE,

and also Mírian L. A. F. Pacheco and the Laboratório Nacional de Luz Síncrotron – LNNano/CNPq (Project SEM-1667 to FCM) for additional SEM analyses. The authors also acknowledge Claudia Inês da Silva for her help with the OM analyses, and Henrique Trad and Thiago Giacometti for the CT imaging at Central de Diagnósticos Ribeirão Preto. The authors thank the editor David J. Bottjer, and an anonymous reviewer who greatly improved the paper with its revision. This research is a contribution of the Laboratório de Paleontologia (FFCLRP-USP). JCAM is grateful to the Programa de Pós-Graduação em Biologia Comparada, FFCLRP-USP. This research was supported by the São Paulo Research Foundation (FAPESP – Fundação de Amparo à Pesquisa do Estado de São Paulo), grants 2010/14797-0 and 2013/23114-1 to JCAM, 2010/19787-2 to AB, 2013/11358-3 to FCM, and 2014/03825-3 to MCL.

## References

- Abercrombie, C.L., Rice, K.G., Hope, C.A., 2001. The great alligator-caiman debate: meditations on crocodilian life-history strategies. In: Grigg, G.C., Seebacher, F., Franklin, C.E. (Eds.), *Crocodylian Biology and Evolution*. Surrey Beatty & Sons, Sydney, pp. 409–418.
- Alonso-Zarza, A.M., 2003. Paleoenvironmental significance of palustrine carbonates and calcretes in the geological record. *Earth Sci. Rev.* 60, 261–298.
- Amaral, G., Bushee, J., Cordani, U.G., Kawashita, K., Reynolds, J.H., 1967. Potassium-argon ages of alkaline rocks from southern Brazil. *Geochim. Cosmochim. Acta* 31, 117–142.
- Ar, A., Paganelli, C.V., Reeves, R.B., Greene, D.G., Rahn, H., 1974. Avian egg: water vapor conductance, shell thickness, and functional pore area. *Condor* 76, 153–158.
- Araújo, R., Castanhinha, R., Martins, R.M., Mateus, O., Hendrickx, C., Beckmann, F., Schell, N., Alves, L.C., 2013. Filling the gaps of dinosaur eggshell phylogeny: Late Jurassic Theropod clutch with embryos from Portugal. *Sci. Report.* 3.
- Araújo, R., Castanhinha, R., Martins, G.G., Nhamutole, N., Du, T.Y., Fernandez, V., Tafforeau, P., Larsson, H., Martins, R.M., Sucena And Léon, É., 2015. Deep time conservative developmental patterns revealed by Crocodylomorpha embryos from the Late Jurassic of Portugal. *Society of Vertebrate Paleontology, 75th Annual Meeting*, 80. TX, Dallas.
- Avilla, L.S., Fernandes, R., Ramos, D.F.B., 2004. Bite marks on a crocodylomorph from the Upper Cretaceous of Brazil: evidence of social behaviour? *J. Vertebr. Paleontol.* 24 (4), 971–973.
- Batezelli, A., 2003. *Análise da Sedimentação Cretácea no Triângulo Mineiro e sua Correlação com Áreas Adjacentes* (PhD Thesis) Instituto de Geociências e Ciências Exatas, Universidade Estadual Paulista, Rio Claro, SP (183 pp).
- Batezelli, A., 2010. Arcabouço tectono-estratigráfico e evolução das bacias Caiuá e Bauru no Sudeste brasileiro. *Revista Brasileira de Geologia* 40 (2), 265–285.
- Batezelli, A., 2015. Continental systems tracts of the Brazilian Cretaceous Bauru Basin and their relationship with the tectonic and climatic evolution of South America. *Basin Res.* 2015, 1–25.
- Bouma, A.H., 1962. *Sedimentology of Some Flysch Deposits, a Graphic Approach to Facies Interpretation*. Elsevier, Amsterdam, p. 168.
- Brazaitis, P., Watanabe, M.E., 2011. Crocodilian behaviour: a window to dinosaur behaviour? *Hist. Biol.* 23, 73–90.
- Bromley, M.H., 1991. Variation in fluvial style as revealed by architecture elements, Kayenta Formation, Mesa Creek, Colorado, USA: evidence for both ephemeral and perennial fluvial processes. In: Miall, A.D., Tyler, N. (Eds.), *The Three-Dimensional Facies Architecture of Terrigenous Clastics Sediments and its Implications for Hydrocarbon Discovery and Recovery vol. 3*. Society for Sedimentary Geology, Tulsa, pp. 94–102.
- Bronzati, M., Montefeltro, F.C., Langer, M.C., 2015. Diversification events and the effects of mass extinctions on Crocodyliformes evolutionary history. *Royal Society Open Science* 2, 140385.
- Carvalho, L.B., Azevedo, S.A.K., Grillo, O.N., Nascimento, P.P., Cabral, U.G., Kellner, A.W.A., Kuroki, I.M.H.C., Kuroki, I.R., 2013. A trull caring mom. In: Lopes, J., Brancaglion Jr., A., Azevedo, S.A.K., Werner Jr., H. (Eds.), *3D Technologies: Unveiling the Past, Shaping the Future*. Lexikon, Rio de Janeiro, pp. 82–87.
- Chiappe, L.M., Coria, R.A., Dingus, L., Jackson, F., Chinsamy, A., Fox, M., 1998. Sauropod dinosaur embryos from the Late Cretaceous of Patagonia. *Nature* 396 (6708), 258–261.
- Chiappe, L.M., Jackson, F., Coria, R.A., Dingus, L., 2005. Nesting titanosaurs from Auka Mahuevo and adjacent sites. In: Curry, R., Kristina, A., Wilson, J.A. (Eds.), *The Sauropods: Evolution and Paleobiology*. University of California Press, Los Angeles & London, pp. 285–302.
- Dal' Bó, P.F.F., Basílico, G., Angélica, R.S., Ladeira, F.S.B., 2009. Paleoclimatic interpretations from pedogenic calcretes in a Maastrihtian semi-arid eolian sand-sheet palaeoenvironment: Marília Formation (Bauru Basin, southeastern Brazil). *Cretac. Res.* 30, 659–675.
- Dal' Bó, P.F.F., Basílico, G., Angélica, R.S., 2010. Factors of paleosol formation in a Late Cretaceous eolian sand sheet palaeoenvironment, Marília Formation, Southeastern Brazil. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 292 (1), 349–365.
- Della Favera, J.C., 1984. Eventos de Sedimentação Episódica nas Bacias Brasileiras. Uma Contribuição para Atestar o Caráter Pontual do Registro Sedimentar. *Anais do Congresso Brasileiro de Geologia* 33. Janeiro, Rio de Janeiro, pp. 489–501.
- Dias-Brito, D., Musacchio, E.A., Castro, J.C., Maranhão, M.S.A.S., Suárez, J.M., Rodrigues, R., 2001. Grupo Bauru: uma unidade continental Cretácea no Brasil – concepções baseadas em dados micropaleontológicos, isotópicos e estratigráficos. *Rev. Paléobiol.* 20, 245–304.



- Ferguson, M.W.J., 1981. Extrinsic microbial degradation of the Alligator eggshell. *Science* 214 (4), 1135–1137.
- Ferguson, M.W.J., 1982. The structure and composition of the eggshell and embryonic membranes of *Alligator mississippiensis*. *Transactions of the Zoological Society of London* 36 (2), 99–152.
- Ferguson, M.W.J., 1985. Reproductive biology and embryology of the crocodilians. In: Gans, C., Billet, F., Maderon, P.F.A. (Eds.), *Biology of Reptilia: Development* (A) 14. John Wiley & Sons Inc., New York.
- Fernández, M.S., García, R.A., Fiorelli, L., Scolaro, A., Salvador, R.B., Cotaro, C.N., Kaiser, G.W., Dyke, G.J., 2013. Large accumulation of avian eggs from the Late Cretaceous of Patagonia (Argentina) reveals a novel nesting strategy in mesozoic birds. *PLoS One* 8 (4), e61030.
- Fiorelli, L.E., Leardi, J.M., Hechenleitner, E.M., Pol, D., Basilici, G., Grellet-Tinner, G., 2015. A new Late Cretaceous crocodyliform from the western margin of Gondwana (La Rioja Province, Argentina). *Cretac. Res.* 60, 194–209.
- Fragoso, C.E., Weinschütz, L.C., Vega, C.S., Guimarães, G.B., Manzig, P.C., Kellner, A.W.A., 2013. Short note on the pterosaurs from the Caiuá Group (Upper Cretaceous, Bauru Basin), Paraná State, Brazil. *Rio Pterio* 2013 – International Symposium on Pterosaurs, pp. 71–72 (Rio de Janeiro, RJ).
- Gobbo-Rodrigues, S.R., 2001. Carófitas e ostrácodos do Grupo Bauru. Masters Dissertation. Instituto de Geociências e Ciências Exatas, Universidade Estadual Paulista, Rio Claro, SP (137 pp).
- Godoy, P.L., Montefeltro, F.C., Norell, M.A., Langer, M.C., 2014. An additional baurusuchid from the Cretaceous of Brazil with evidence of interspecific predation among Crocodyliformes. *PLoS One* 9 (5), e97138.
- Godoy, P.L., Bronzati, M., Eltink, Marsola, J.C.A., Cidade, G.M., Langer, M.C., Montefeltro, F.C., 2016. Postcranial anatomy of *Pissarrachampsia sera* (Crocodyliformes, Baurusuchidae) from the Late Cretaceous of Brazil: insights on lifestyle and phylogenetic significance. *PeerJ* 4, e2075. <http://dx.doi.org/10.7717/peerj.2075>.
- Greer, A.E., 1975. Clutch size in crocodilians. *J. Herpetol.* 9 (3), 319–322.
- Grellet-Tinner, G., Fiorelli, L.E., 2010. A new Argentinean nesting site showing neosauropod dinosaur reproduction in Cretaceous hydrothermal environment. *Nat. Commun.* 1, 32.
- Grellet-Tinner, G., Chiappe, L., Norell, M., Bottjer, D., 2006. Dinosaur eggs and nesting behaviors: a paleobiological investigation. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 232, 294–321.
- Grellet-Tinner, G., Codrea, V., Folie, A., Higa, A., Smith, T., 2012. First evidence of reproductive adaptation to “island effect” of a dwarf cretaceous Romanian titanosaur, with embryonic integument *in ovo*. *PLoS One* 7 (3), e32051. <http://dx.doi.org/10.1371/journal.pone.0032051>.
- Grigg, G., Kirshner, D., 2015. *Biology and Evolution of Crocodylians*. Csiro Publishing, Ithaca and London (671 pp).
- Guimarães, G.B., Liccardo, A., Godoy, L.C., Weinschütz, L.C., Manzig, P.C., Vega, C.S., Pilatti, F., 2012. Ocorrência de Pterossauros no Cretáceo da Bacia do Paraná/Bauru: Implicações para a geoconservação, paleontologia e estratigrafia. 46th Geology Brazilian Congress. 48. Santos, SP.
- Hastings, A.K., Hellmund, M., 2015. Rare *in situ* preservation of adult crocodilian with eggs from the Middle Eocene of Geiseltal, Germany. *PALAIOS* 30 (6), 446–461.
- Hasui, Y., Cordani, U.G., 1968. Idades potássio-argônio de rochas eruptivas mesozóicas do oeste mineiro e sul de Goiás. Congresso Brasileiro de Geologia, Belo Horizonte.
- Hayward, J.L., Zelenitsky, D.K., Smith, D.L., Zaft, D.M., Clayburn, J.K., 2000. Eggshell taphonomy at modern gull colonies and dinosaur clutch site. *PALAIOS* 15, 343–355.
- Hayward, J.L., Dickson, K.M., Gamble, S.R., Owen, A.W., Owen, K.C., 2011. Eggshell taphonomy: environmental effects on fragment orientation. *Hist. Biol.* 23 (1), 5–13.
- Hechenleitner, E.M., Grellet-Tinner, G., Fiorelli, L.E., 2015. What do giant titanosaur dinosaurs and modern Australasian megapodes have in common? *PeerJ* 3, e1341.
- Hirsch, K.F., 1985. Fossil crocodilian egg from the Eocene of Colorado. *J. Paleontol.* 59 (3), 531–542.
- Hirst, J.P.P., 1991. Variations in alluvial architecture across the Oligo-Miocene Huesca Fluvial System Ebro Basin, Spain. In: Miall, A.D., Tyler, N. (Eds.), *The Three-Dimensional Facies Architecture of Terrigenous Clastics Sediments and Its Implications for Hydrocarbon Discovery and Recovery* vol. 3. Society for Sedimentary Geology, Tulsa, pp. 111–121.
- Horner, J.R., Makela, R., 1979. Nest of juveniles provides evidence of family structure among dinosaurs. *Nature* 282, 296–298.
- Jenny, H.J., 1941. *Factors of Soil Formation*. McGraw-Hill, New York (281 pp).
- Jones, S.J., Frostick, L.E., Astin, T.R., 2001. Braided stream and flood plain architecture: the Rio Vero Formation, Spanish Pyrenees. *Sediment. Geol.* 139, 229–260.
- Klappa, C.P., 1980. Rhizolites in terrestrial carbonates: classification, recognition, genesis and significance. *Sedimentology* 27, 613–629.
- Klappa, C.P., 1983. A process-response model for the formation of pedogenic calcretes. In: Wilson, R.C.L. (Ed.), *Residual Deposits: Surface Related Weathering Processes and Materials*Oxford: Geological Society, Special Publication. Blackwell Scientific Publications, pp. 211–220.
- Kohring, R., Hirsch, K.F., 1996. Crocodilian and avian eggshells from the Middle Eocene of the Geiseltal, Eastern Germany. *J. Vertebr. Paleontol.* 16 (1), 67–80.
- Leardi, J.M., Pol, D., Novas, F.E., Suarez Riglos, M., 2015. The postcranial anatomy of *Yacarerani boliviensis* and the phylogenetic significance of the notosuchian postcranial skeleton. *J. Vertebr. Paleontol.*, e995187.
- Machado Júnior, D.L., 1992. Idades Rb/Sr do complexo alcalino-carbonatítico de Catalão II (GO). 29th Brazilian Geology Congress Bulletin, pp. 91–93 (São Paulo, SP).
- Magalhães Ribeiro, C.M., Arruda Campos, A.C., 2009. Aspectos tafonômicos de posturas de ovos de crocodyliformes da Formação Adamantina, Bacia Bauru. Livro de Resumos XXI Congresso Brasileiro de Paleontologia, Belém-PA.
- Magalhães Ribeiro, C.M., Carvalho, I.S., Nava, W.R., 2006. Ovos de crocodyliformes da Formação Araçatuba (Bacia Bauru, Cretáceo Superior), Brasil. In: Gallo, V.B., Silva, H.M.A., Figueiredo, F.J. (Eds.), *Paleontologia de Vertebrados, Grandes Temas e Contribuições Científicas*. Interciência, Rio de Janeiro.
- Magnusson, W.E., 1986. The peculiarities of crocodilian population dynamics and their possible importance for management strategies. *Proceedings of the 7th Working Meeting of the IUCN–SSC Crocodile Specialist Group*. IUCN–World Conservation Union, Gland, Caracas, Venezuela, pp. 434–442.
- Marsola, J.C.A., Montefeltro, F.C., Langer, M.C., 2011. Baurusuchid eggs from the Vale do Rio do Peixe Formation (Bauru Group), Late Cretaceous of Minas Gerais, Brazil. *Resúmenes del Congreso Latinoamericano de Paleontología de Vertebrados*, San Juan, Argentina.
- Martinelli, A.G., Teixeira, V.P.A., 2015. The Late Cretaceous vertebrate record from the Bauru Group in the Triângulo Mineiro, southeastern Brazil. *Bol. Geol. Min.* 126.
- Martinelli, A.G., Riff, D., Lopes, R.P., 2011. Discussion about the occurrence of the genus *Aeolosaurus* Powell 1987 (Dinosauria, Titanosauria) in the Upper Cretaceous of Brazil. *J. Geosci.* 7, 34–40.
- Marzola, M., Russo, J., Mateus, O., 2015. Identification and comparison of modern and fossil crocodilian eggs and eggshell structure. *Hist. Biol.* 27 (1), 115–133.
- Mateus, I., Mateus, H., Antunes, M.T., Mateus, O., Taquet, P., Ribeiro, V., Manuppella, G., 1998. Upper Jurassic theropod dinosaur embryos from Lourinhã (Portugal). *Memórias da Academia de Ciências de Lisboa* 37.
- Miall, A.D., 1985. Architectural-element analysis: a new method of facies analysis applied to fluvial deposits. *Earth-Sci. Rev.* 22 (4), 261–300.
- Miall, A.D., 1988a. Reservoir heterogeneities in fluvial sandstones: lessons from outcrop studies. *Am. Assoc. Pet. Geol. Bull.* 72 (6), 682–697.
- Miall, A.D., 1988b. Architectural elements and bounding surfaces in fluvial deposits: anatomy of the Kayenta Formation (Lower Jurassic) Southwest Colorado. *Sediment. Geol.* 55 (2), 233–262.
- Miall, A.D., 1996. *The Geology of Fluvial Deposits*. Springer-Verlag, Berlin (582 pp).
- Montefeltro, F.C., Larsson, H.C.E., Langer, M.C., 2011. A new baurusuchid (Crocodyliformes, Mesoeucrocodylia) from the Late Cretaceous of Brazil and the phylogeny of baurusuchidae. *PLoS One* 6 (7), 1–26.
- Montefeltro, F.C., Larsson, H.C.E., França, M.A.G., Langer, M.C., 2013. A new neosuchian with Asian affinities from the Jurassic of northeastern Brazil. *Naturwissenschaften* 100, 835–841.
- Moreno-Azanza, M., Bauluz, B., Canudo, J.I., Puértolas-Pascual, E., Sellés, A.G., 2014. A re-evaluation of aff. *Megaloolithidae* eggshell fragments from the uppermost Cretaceous of the Pyrenees and implications for crocodylomorph eggshell structure. *Hist. Biol.* 26 (2), 195–205.
- Norell, M.A., Clark, J.M., Chiappe, L.M., Dashzeveg, D., 1995. A nesting dinosaur. *Nature* 378, 774–776.
- Novas, F., Pais, D.F., Pol, D., Carvalho, A.B., Scanfaleria, A., Mones, A., Riglos, M.S., 2009. Bizarre Notosuchian crocodyliform with associated eggs from the Upper Cretaceous of Bolivia. *J. Vertebr. Paleontol.* 29, 1316–1320.
- Oliveira, C.E.M., 2008. *Associações de ovos de crocodylomorfos da Formação Adamantina, Grupo Bauru, Cretáceo Superior, na região de Jales-SP* (PhD Thesis) Instituto de Geociências e Ciências Exatas, Universidade Estadual Paulista, Rio Claro, SP (117 pp).
- Oliveira, C.E.M., Santucci, R.M., Andrade, M.B., Fulfaro, V.J., Basilio, J.A.F., Benton, M.J., 2011. Crocodylomorph eggs and eggshells from the Adamantina Formation (Bauru Group), Upper Cretaceous of Brazil. *Paleontology* 54, 309–321.
- Oser, S.E., Jackson, F.D., 2014. Sediment and eggshell interactions: using abrasion to assess transport in fossil eggshell accumulations. *Hist. Biol.* 26 (2), 165–172.
- Packard, M.J., Packard, G.C., Boardman, T.J., 1982. Structure of eggshells and water relations of reptilian eggs. *Herpetologica* 136–155.
- Panadés I Blas, X., Patnaik, R., 2009. A complete crocodylian egg from the Upper Miocene (Chinji Beds) of Pakistan and its palaeobiographical implications. *J. Vertebr. Paleontol.* 6 (1).
- Pereira, C.T., Batezelli, A., Ladeira, F.S.B., 2015. Paleoprecipitation changes based on paleosols profiles of the Marília Formation (Upper Cretaceous) in the eastern portion of the Bauru Basin in Southeastern Brazil. *Geociências* 34 (2), 238–257.
- Platt, S.G., Resbat, A., Stuart, B.L., 2004. Maximum clutch size of the American Alligator. *Florida Field Naturalist* 32 (31), 102–106.
- Pol, D., Leardi, J.M., Lecuona, A., Krause, M., 2012. Postcranial anatomy of *Sebecus icaeorhinus* (Crocodyliformes, Sebecidae) from the Eocene of Patagonia. *J. Vertebr. Paleontol.* 32, 328–354.
- Pol, D., Nascimento, P.M., Carvalho, A.B., Riccomini, C., Pires-Domingues, R.A., Zaher, H., 2014. A new notosuchian from the Late Cretaceous of Brazil and the phylogeny of advanced notosuchians. *PLoS One* 9, e93105.
- Rabi, M., Sebök, N., 2015. A revised Eurogondwana model: notosuchian crocodyliforms and other vertebrate taxa suggest the retention of episodic faunal links during the most of the Cretaceous. *Gondwana Res.* 28, 1197–1211.
- Ramos, A., Sopena, A., 1983. Gravel bars in low-sinuosity streams (Permian and Triassic, Central Spain). In: Collinson, J.D., Lewis, J. (Eds.), *Modern and Ancient Fluvial Systems* Collinson The International Association of Sedimentologists, Special Publication 6. John Wiley & Sons Inc., New York, pp. 301–312.
- Retallack, G.J., 2001. *Soils of the Past, an Introduction to Pedology*. John Wiley & Sons Inc., New York, p. 404.
- Retallack, G.J., 2005. Pedogenic carbonate proxies for amount and seasonality of precipitation in paleosols. *Geology* 33 (4), 333–336.
- Ribeiro, V., Mateus, O., Holwerda, F., Araújo, R., Castanheira, R., 2014. Two new theropod egg sites from the late Jurassic Lourinhã Formation, Portugal. *Hist. Biol.* 26 (2), 206–217.
- Riff, D., Kellner, A.W.A., 2011. Baurusuchid crocodyliforms as theropod mimics: clues from the skull and appendicular morphology of *Stratiotosuchus maxhechti* (Upper Cretaceous of Brazil). *Zool. J. Linnean Soc.* 163 (s1), S37–S56.
- Rogers, J.V., 2001. A complete crocodyloid egg from the Lower Cretaceous (Albian) Glen Rose Formation, central Texas. *J. Vertebr. Paleontol.* 20 (4), 780–783.

- Russo, J., Mateus, O., Balbino, A., Marzola, M., 2014. Crocodylomorph eggs and eggshells from the Lourinhã Fm. (Upper Jurassic), Portugal. *Comunicações Geológicas* 101 (Especial I), 563–566.
- Santucci, R.M., Bertini, R.J., 2001. Distribuição paleogeográfica e biocronológica dos Titanossauros (Saurishia, Sauropoda) do Grupo Bauru, Cretáceo Superior do sudeste Brasileiro. *Revista Brasileira de Geociências* 31, 307–315.
- Sato, T., Cheng, Y., Wu, X., Zelenitsky, D.K., Hsiao, Y., 2005. A pair of shelled eggs inside a female dinosaur. *Science* 308, 375.
- Schweitzer, M.H., Jackson, F.D., Chiappe, L.M., Schmitt, J.G., Calvo, J.O., Rubilar, D.E., 2002. Late Cretaceous avian eggs with embryos from Argentina. *J. Vertebr. Paleontol.* 22 (1), 191–195.
- Seymour, R.S., 1979. Dinosaur eggs: gas conductance through the shell, water loss during incubation and clutch size. *Paleobiology* 5 (01), 1–11.
- Soares, P.C., Landim, P.M.B., Fulfaro, V.J., Sobreiro Neto, A.F., 1980. Ensaio de caracterização estratigráfica do Cretáceo no Estado de São Paulo: Grupo Bauru. *Revista Brasileira de Geociências* 10 (3), 177–185.
- Sonoki, I.K., Garda, G.M., 1988. Idades K-Ar de rochas alcalinas do Brasil Meridional e Paraguai Oriental: compilação e adaptação às novas constantes de decaimento. *Boletim IG-USP. Série Científica* 19, 63–85.
- Srivastava, R., Patnaik, R., Shukla, U.K., Sahni, A., 2015. Crocodilian nest in a Late Cretaceous sauropod hatchery from the Type Lameta Ghat locality, Jabalpur, India. *PLoS One* 10 (12), e0144369.
- Stanistreet, I.G., McCarthy, T.S., 1993. The Okavango Fan and the classification of sub aerial fan systems. *Sediment. Geol.* 85, 115–133.
- Suguio, K., Barcelos, J.H., 1983. Calcretes of the Bauru Group (Cretaceous) Brazil: petrology and geological significance. *Boletim do Instituto de Geociências* 14, 31–47.
- Suguio, K., Berenholc, M., Salati, E., 1975. Composição química e isotópica dos calcários e ambiente de sedimentação da Formação Bauru. *Boletim do Instituto de Geociências* 6, 55–75.
- Tanaka, K., Zelenitsky, D.K., Therrien, F., 2015. Eggshell porosity provides insight on evolution of nesting in Dinosaurs. *PLoS One* 10 (11), e0142829.
- Tucker, A.D., 2001. Sensitivity analysis of stagestructured demographic models for freshwater crocodiles. In: Grigg, G.C., Seebacher, F., Franklin, C.E. (Eds.), *Crocodylian Biology and Evolution*. Surrey Beatty & Sons, Sydney, pp. 349–363.
- Vasconcellos, F.M., Carvalho, I.S., 2010. Paleochronological assemblage associated with *Baurusuchus salgadoensis* remains, a Baurusuchidae Mesoeucrocodylia from the Bauru Basin, Brazil (Late Cretaceous). *Bulletin of the New Mexico Museum of Natural History and Science* 51, 227–237.
- von Der Borch, C., 1965. The distribution and preliminary geochemistry of modern carbonate sediments of the Coorong area, South Australia. *Geochim. Cosmochim. Acta* 29 (7), 781–799.
- Watson, A., 1992. Desert soils. In: Martini, I.P., Chesworth, W. (Eds.), *Weathering, Soli & Paleosols*. Elsevier, Amsterdam, pp. 225–260.
- Weissmann, G.S., Hartley, A.J., Nichols, G.J., Scuderi, L.A., Olson, M., Buehler, H., Banteah, R., 2010. Fluvial form in modern continental sedimentary basins: distributive fluvial systems. *Geology* 38 (1), 39–42.
- Wright, V.P., 2007. Calcrete. In: Nash, D.J., McLaren, S.J. (Eds.), *Geochemical Sediments and Landscapes*. Blackwell, Oxford, pp. 10–45.
- Wright, P., Tucker, M.E., 1991. Calcretes: an introduction. In: Wright, P., Tucker, M.E. (Eds.), *Calcretes*. Blackwell Scientific Publications, Oxford, pp. 1–22.
- Zhou, Z., Zhang, F., 2004. A precocial avian embryo from the Lower Cretaceous of China. *Science* 306 (5696), 653.