

## SHORT COMMUNICATION

# Macrophyte rafts as dispersal vectors for fishes and amphibians in the Lower Solimões River, Central Amazon

Luis Schiesari<sup>\*1</sup>, Jansen Zuanon<sup>†</sup>, Claudia Azevedo-Ramos<sup>‡</sup>, Marcelo Garcia<sup>§</sup>, Marcelo Gordo<sup>§</sup>, Mariluce Messias<sup>¶</sup> and Emerson Monteiro Vieira<sup>#</sup>

<sup>\*</sup>Departamento de Zoologia, Instituto de Biociências, Universidade de São Paulo, CP 20520, 01452-990, São Paulo-SP, Brazil

<sup>†</sup>Coordenação de Pesquisas em Biologia Aquática, Caixa Postal 478, Instituto Nacional de Pesquisas da Amazônia, 69083–970, Manaus-AM, Brazil

<sup>‡</sup>Núcleo de Altos Estudos Amazônicos-NAEA, Universidade Federal do Pará 67070-100, Belém-PA, Brazil

<sup>§</sup>Departamento de Biologia, Universidade do Amazonas, 69970-000, Manaus-AM, Brazil

<sup>¶</sup>UNESP-Rio Claro, PG Zoologia, Avenida 24-A, 1515, C.P. 199, 13506-900, Rio Claro-SP, Brazil

<sup>#</sup>Laboratório de Ecologia de Mamíferos, Centro de Ciências da Saúde, UNISINOS, CP 275, 93022-000, São Leopoldo-RS, Brazil

(Accepted 9 June 2002)

**Key Words:** Amazon, amphibians, biodiversity, biogeography, dispersal, fishes, macrophytes, várzea floodplains

Large rivers have played a prominent role in biogeographic theory for their potential to act as barriers for the dispersal of terrestrial organisms, and therefore be involved in the generation of species diversity (Brown & Lomolino 1998). In this paper, we document the potential role of macrophyte rafts as a mechanism by which Amazonian rivers could act as dispersal agents rather than barriers, transferring organisms across banks and possibly across very large distances. These vectors could therefore act against speciation and towards homogenization of the local biota.

These rafts originate from extensive macrophyte stands that grow along the margins and banks of the nutrient-rich, white-water rivers and lakes in the Amazonian várzea floodplains (Junk 1970, 1973; Junk & Piedade 1993, 1997; Junk *et al.* 1989). Macrophyte stands support an extremely rich community of over 380 species of herbs of floating and rooted habits, but are commonly dominated by the grasses *Paspalum repens* Berg. and *Echinochloa polystachya* (H.B.K.) Hitch. and floating vegetation such as the water hyacinth (*Eichhornia crassipes* (Mart.) Solms.) and the pteridophyte *Salvinia auriculata* Aubl. (Junk & Piedade 1993). Macrophyte stands constitute a peculiar habitat encompassing aquatic and terrestrial biotopes. The rapid growth and production of detritus, and shelter availability in the root zone of macrophyte stands

provide conditions for the development of an abundant aquatic fauna of invertebrates (especially microcrustaceans, molluscs and insects; Junk & Robertson 1997) and vertebrates (mostly fish; Henderson & Hamilton 1995, Junk *et al.* 1997). The aerial portion is habitat for invertebrates (mostly insects and arachnids), anuran amphibians, which use the stands of floating macrophytes as calling and breeding sites (Hödl 1977), and for other vertebrates such as birds (Petermann 1997).

After explosive growth during the period of rising water levels, stalks of floating meadows often get weak and break, forming drifting rafts that may flush into the main river channel and be carried away by wind and water currents (Junk & Piedade 1997).

Field work was conducted on 20 and 21 August 1994 in the Ilha da Marchantaria (03°14'S, 59°57'W), near Manaus, Central Amazon, Brazil. This island is located in the lower Solimões river, 15 km above the confluence with the Rio Negro. Following the yearly peak in water level in June (Irion *et al.* 1997), around August the strongest water currents occur and many macrophyte rafts drift down the river. We approached the rafts by boat, and measured their length and width. Rafts were encircled with a 7-m × 3.5-m (mesh 0.5 cm) seine net. Rafts were not compact and therefore could be entirely encircled even if their perimeter exceeded net length (7 m). The net was subsequently brought to the boat, where every plant was carefully screened for vertebrates. All vertebrates were collected, identified whenever possible to species,

<sup>1</sup> Corresponding author. Present address: Department of Ecology and Evolutionary Biology, University of Michigan, 48109-1048, Ann Arbor-MI, USA. Email: lschiesia@umich.edu

and counted. Fish were identified as adults or juveniles according to Géry (1977), Kullander (1986), Burgess (1989) and Botero & Araújo-Lima (2001). Voucher specimens were deposited in the collections of the Instituto Nacional de Pesquisas da Amazônia, Manaus, Brazil.

We sampled eight floating macrophyte rafts of 3.02–24.7 m<sup>2</sup> (mean  $\pm$  SD: 9.11  $\pm$  6.30 m<sup>2</sup>) which were found drifting down the river. Two rafts were constituted predominantly of *Eichhornia crassipes*, and six of *Paspalum repens*. Associated vertebrates were represented by fishes and amphibians (Table 1).

Fishes were present in the root zone of all rafts. We found 286 individuals of 39 species and 19 families (Table 1). Characidae were the most abundant (32.9% of all individuals,  $n = 94$ ) and species-rich (35.9% of all species,  $n = 14$ ) family, followed closely by Cichlidae in abundance (29.0%,  $n = 83$ ), but not species richness (10.2%,  $n = 4$ ). Hypopomidae were represented by 35 individuals (12.2%) and three species (7.7%). Aptereronotidae and Serrasalminae were represented by two species each, while the 14 remaining families were represented by only one species and most commonly by less than four individuals each. The most common species were *Apistogramma agassizi* (Cichlidae), *Hyphessobrycon eques* (Characidae) and *Pyrhulina* sp. (*brevis* species group) (Lebiasinidae), with respectively 49 (17.1%), 24 (8.4%) and 23 (8.0%) individuals. Fish species richness was significantly related to raft area (number of species =  $3.87 + 0.81 \times \text{area}$ ;  $r^2 = 0.70$ ,  $P = 0.01$ ). No significant relationship was found between fish abundance and raft area. Nearly half (49.30%) of individual fish collected were juveniles.

Amphibians occurred on six rafts. We found 42 individuals of nine species belonging to five families, four of Anura (Hylidae, Leptodactylidae, Pseudidae and Bufonidae) and one of Gymnophiona (Typhlonectidae) (Table 1). Hylidae were the most abundant (83.3% of all individuals,  $n = 35$ ) and species-rich (55.5% of all species,  $n = 5$ ) family. The four remaining families were each represented by a single species and 1–3 individuals. The tree-frog *Hyla walfordi* was by far the most abundant species (45.2% of all individuals,  $n = 19$ ). In one single raft, with an area of 8.48 m<sup>2</sup>, we found 11 individuals of *H. walfordi*, two of them gravid females. A gravid female of *Hyla leucophyllata* was found in another raft. Only five anurans (11.9%) collected were immatures (three tadpoles and two metamorphs). We did not find any significant relationship between raft area and amphibian species richness or abundance.

A rich and abundant invertebrate fauna was found associated to these rafts, including: a leech, pulmonate molluscs, spiders, crustaceans (shrimps and crabs), and insects such as odonates (naiads and adults), blattarians, orthopterans (ensiferans and caeliferans), homopterans, belostomatid heteropterans, hymenopterans (ants), lepid-

opterans (larvae and adults) and coleopterans (larvae, pupae and adults). Particularly abundant were shrimps, spiders and orthopterans.

Aquatic macrophyte rafts have been suggested to act as dispersal vectors in large lakes (Lake Mamiraua: Henderson & Hamilton 1995; Lake Malawi: Oliver & McKaye 1982) and rivers (Parana River: Achaval *et al.* 1979) for both aquatic and terrestrial faunas, although the effectiveness can vary depending on the fish group considered (Henderson & Hamilton 1995). Albeit preliminary, our sampling in the Solimões River indicates that floating macrophyte rafts can transport a remarkably diverse and abundant vertebrate assemblage. In addition to the 39 species of fishes and nine of amphibians found (Table 1), at least 13 other species of fish and seven of amphibians live in aquatic macrophyte stands (Hödl 1977, Junk 1973). Many of the fish and amphibian species that we found in the floating rafts show very wide distributional ranges in the floodplains of the Amazon basin (Burgess 1989, Frost 1985, Géry 1977, Kullander 1986). Dispersal of vertebrates through the Amazon River seems to be a common phenomenon and floating macrophyte rafts may be acting as important vectors. Moreover, transportation by floating rafts may represent an unusually predictable vector of long-distance dispersal both from spatial (unidirectional) and temporal (seasonal) standpoints (see also Henderson & Hamilton 1995).

This predictability could have favoured selection for the utilization of macrophyte stands as breeding sites and nursery grounds for several fish species. Rafts could transport juveniles and therefore return individuals to the populations of the several fish species that migrate upriver to breed. Consistent with these hypotheses, Sazima & Zampogno (1985) observed that young piranhas sheltering among the roots of water hyacinths had no fin damage, in contrast to larger juveniles and adults. They also suggested that, apart from shelter, young piranhas might profit from water hyacinth transportation. This could explain why piranhas, which spawn during the annual floods, have a wide distribution despite the absence of known migratory movements by the adults in Amazonian rivers (Goulding 1980).

The effectiveness of the rafts as a long-distance dispersal vector is further reinforced by the observation that average current velocities in the Amazon River range from 1.0 to 1.5 m s<sup>-1</sup> in the wet season (Sioli 1975). Under such conditions a raft could travel 86–130 km per day, or, in other words, travel the whole extension of the Solimões/Amazonas River system (4000 km) in as little as 31 d. The probability of establishing viable populations upon arrival could be enhanced both by the synchronization of the drift and reproductive period, and by the large observed densities of fishes (up to 10.5 m<sup>-2</sup>) and amphibians (up to 1.65 m<sup>-2</sup>). In summary, drifting of many rafts containing dense vertebrate populations in

**Table 1.** Absolute frequency of fish and amphibian families and species found in eight macrophyte rafts in the lower Solimões river, Amazonas, Brazil. Individuals classified as adults (A) or juveniles (J) (<sup>a</sup> = tadpoles; <sup>b</sup> = metamorphs).

Family	Species	Number of individuals			Number of rafts in which the species was found	Individuals per family	
		J	A	Total			
FISHES							
Ageneiosidae	<i>Ageneiosus</i> sp.	1	–	1	1	1	
Anostomidae	<i>Leporinus fasciatus</i> (Bloch, 1794)	1	–	1	1	1	
Apteronotidae	<i>Apteronotus albifrons</i> (Linnaeus, 1766)	–	1	1	1	2	
	<i>Apteronotus hasemani</i> (Ellis, 1913)	–	1	1	1		
Auchenipteridae	<i>Parauchenipterus galeatus</i> (Linnaeus, 1766)	3	1	4	4	4	
Callichthyidae	<i>Megalechis thoracata</i> (Valenciennes, 1840)	1	–	1	1	1	
Characidae	<i>Aphyocharax</i> sp.	8	5	13	3	94	
	<i>Charax</i> sp.	13	–	13	1		
	<i>Ctenobrycon spilurus</i> (Valenciennes, 1850)	3	–	3	3		
	<i>Hemigrammus</i> sp. 1	3	5	8	2		
	<i>Hemigrammus</i> sp. 2	–	1	1	1		
	<i>Hemigrammus</i> sp. 3	7	1	8	3		
	<i>Hemigrammus</i> sp. 4	1	–	1	1		
	<i>Hyphessobrycon eques</i> (Steindachner, 1882)	10	14	24	3		
	<i>Hyphessobrycon</i> sp. 1	5	5	10	5		
	<i>Hyphessobrycon</i> sp. 2	–	1	1	1		
	<i>Hyphessobrycon</i> sp. 3	–	2	2	1		
	<i>Moenkhausia</i> sp. ( <i>intermedia</i> group)	–	7	7	2		
	<i>Prionobrama filigera</i> (Cope, 1870)	–	1	1	1		
	<i>Roeboides</i> sp.	2	–	2	2		
	Crenuchidae	<i>Crenuchus spilurus</i> Günther, 1863	–	1	1	1	2
		<i>Klausewitzia</i> sp.	–	1	1	1	
	Cichlidae	<i>Apistogramma agassizi</i> (Steindachner, 1875)	27	22	49	3	83
<i>Apistogramma</i> sp.		6	12	18	3		
<i>Cichlasoma amazonarum</i> (Kullander, 1883)		6	6	12	6		
<i>Crenicichla inpa</i> (Ploeg, 1991)		3	1	4	4		
Erythrinidae	<i>Hoplias</i> cf. <i>malabaricus</i>	3	–	3	3	3	
Gymnotidae	<i>Gymnotus</i> aff. <i>stenoleucus</i>	2	–	2	1	2	
Hypopomidae	<i>Brachyhypopomus brevirostris</i> (Steindachner, 1868)	6	12	18	3	35	
	<i>Brachyhypopomus pinnicaudatus</i> (Hopkins, 1991)	3	13	16	3		
	<i>Brachyhypopomus</i> sp. n.	1	–	1	1		
	<i>Pyrrhulina</i> sp. ( <i>brevis</i> group)	2	21	23	5	23	
Lepidosirenidae	<i>Lepidosiren paradoxa</i> (Fitzinger, 1837)	–	1	1	1	1	
Pimelodidae	<i>Paulicea luetkeni</i> (Steindachner, 1875)	1	–	1	1	1	
Rivulidae	<i>Rivulus</i> aff. <i>ornatus</i>	1	1	2	2	2	
Serrasalminidae	<i>Metynnis</i> sp.	2	–	2	2	13	
	<i>Serrasalmus</i> sp.	11	–	11	4		
Sternopygidae	<i>Eigenmannia</i> sp.	5	5	10	3	10	
Synbranchidae	<i>Synbranchus</i> sp.	4	4	8	5	8	
Total	19	39	141	145	286	–	
AMPHIBIANS							
Bufonidae	<i>Bufo marinus</i> (Linnaeus, 1758)	2 <sup>b</sup>	–	2	1	2	
Leptodactylidae	<i>Leptodactylus</i> aff. <i>leptodactyloides</i>	–	1	1	1	1	
Hylidae	<i>Hyla walfordi</i> Bokermann, 1962	–	19	19	3	35	
	<i>Hyla leucophyllata</i> (Beireis, 1783)	–	4	4	2		
	<i>Hyla raniceps</i> (Cope, 1862)	–	2	2	2		
	<i>Hyla punctata</i> (Schneider, 1799)	2 <sup>a</sup>	2	4	3		
	<i>Sphaenorhynchus carneus</i> (Cope, 1868)	–	5	5	1		
	Unidentified species	1 <sup>a</sup>	–	1	1		
Pseudidae	<i>Lysapsus laevis</i> Parker, 1935	–	3	3	3	3	
Typhlonectidae	<i>Typhlonectes compressicauda</i> (Duméril and Bibron, 1841)	–	1	1	1	1	
Total	5	9–10	5	37	42	–	

breeding condition provides a scenario consistent with the hypothesis of an effective long-distance dispersal vector. This reported dispersal system could be in part responsible for the wide distribution of many of the amphibian and fish species found associated to Amazonian white-water riverine systems.

## ACKNOWLEDGEMENTS

This manuscript has been improved with the criticism of Walter Hödl, Richard Lehtinen, Woodruff Benson, Miguel Trefaut Rodrigues, Antonio Carlos Marques and Martin Henzl. This study was conducted during the field course Ecology of the Amazonian Forest, promoted by the Organization of Tropical Studies, Universidade Estadual de Campinas, Smithsonian Institution and Instituto Nacional de Pesquisas da Amazônia. We are indebted to these institutions for financial and logistical support. LCS was funded by a grant of Conselho Nacional de Desenvolvimento Científico e Tecnológico (200093/97-5) during the final preparation of this manuscript.

## LITERATURE CITED

- ACHAVAL, F., GONZÁLES, J. G., MENEGHEL, M. & MELGA-REJO, A. 1979. Lista comentada del material recogido en costas uruguayas, transportado por camalotes desde el río Parana. *Acta Zoologica Lilloana* 35:195–200.
- BOTERO, J. I. S. & ARAÚJO-LIMA, C. A. R. M. 2001. As macrófitas aquáticas como berçário para a ictiofauna da várzea do rio Amazonas. *Acta Amazonica* 31:437–447.
- BROWN, J. H. & LOMOLINO, M. V. 1998. *Biogeography*. Second edition. Sinauer Associates, Sunderland. 691 pp.
- BURGESS, W. E. 1989. *An atlas of freshwater and marine catfishes*. TFH Publications, Neptune. 784 pp.
- FROST, D. R. 1985. *Amphibian species of the world: a taxonomic and geographical reference*. Allen Press and Association of Systematics Collections, Lawrence. 732 pp.
- GÉRY, J. 1977. *Characoids of the world*. TFH Publications, Neptune. 672 pp.
- GOULDING, M. 1980. *The fishes and the forest: explorations in Amazonian natural history*. University of California Press, Berkeley. 280 pp.
- HENDERSON, P. A. & HAMILTON, H. F. 1995. Standing crop and distribution of fish in drifting and attached floating meadow within an Upper Amazonian varzea lake. *Journal of Fish Biology* 47:266–276.
- HÖDL, W. 1977. Call differences and calling site segregation in anuran species from Central Amazonian floating meadows. *Oecologia* 28:351–363.
- IRION, G., JUNK, W. J. & DE MELLO, J. A. S. N. 1997. The large Central Amazonian River floodplains near Manaus: geological, climatological, hydrological, and geomorphological aspects. Pp. 23–46 in Junk, W. (ed.). *Ecological Studies 126, The Central Amazon floodplain – ecology of a pulsing system*. Springer-Verlag, Berlin.
- JUNK, W. 1970. Investigations on the ecology and production-biology of the 'floating meadows' (*Paspalum-Echinochloetum*) on the middle Amazon. Part I: the floating vegetation and its ecology. *Amazoniana* 2:449–495.
- JUNK, W. 1973. Investigations on the ecology and production-biology of the 'floating meadows' (*Paspalum-Echinochloetum*) on the middle Amazon. Part II: The aquatic fauna in the root zone of floating vegetation. *Amazoniana* 4:9–102.
- JUNK, W., BAYLEY, P. B. & SPARKS, R. E. 1989. The flood pulse concept in river-floodplain systems. Pp. 110–127 in Dodge, D. P. (ed.). *Proceedings of the International Large River Symposium (LARS)*. Canadian special publication of fisheries and aquatic sciences 106.
- JUNK, W. & PIEDADE, M. T. F. 1993. Herbaceous plants of the Amazon floodplain near Manaus: species diversity and adaptations to the flood pulse. *Amazoniana* 12:467–484.
- JUNK, W. & PIEDADE, M. T. F. 1997. Plant life in the floodplain with special reference to herbaceous plants. Pp. 147–186 in Junk, W. (ed.). *Ecological Studies 126, The Central Amazon floodplain – ecology of a pulsing system*. Springer-Verlag, Berlin.
- JUNK, W. J. & ROBERTSON, B. A. 1997. Aquatic invertebrates. Pp. 279–298 in Junk, W. (ed.). *Ecological Studies 126, The Central Amazon floodplain – ecology of a pulsing system*. Springer-Verlag, Berlin.
- JUNK, W., SOARES, M. G. M. & SAINT-PAUL, U. 1997. The fish. Pp. 385–408 in Junk, W. (ed.). *Ecological Studies 126, The Central Amazon floodplain – ecology of a pulsing system*. Springer-Verlag, Berlin.
- KULLANDER, S. O. 1986. *Cichlid fishes of the Amazon River drainage of Peru*. Swedish Museum of Natural History, Stockholm. 431 pp.
- OLIVER, M. K. & MCKAYE, K. R. 1982. Floating islands – a means of fish dispersal in Lake Malawi, Africa. *Copeia* 1982:748–754.
- PETERMANN, P. 1997. The birds. Pp. 419–452 in Junk, W. (ed.). *Ecological Studies 126, The Central Amazon floodplain – ecology of a pulsing system*. Springer-Verlag, Berlin.
- SAZIMA, I. & ZAMPROGNO, C. 1985. Use of water hyacinths as shelter, foraging place, and transport by young piranhas, *Serrasalmus spilopleura*. *Environmental Biology of Fishes* 12:237–240.
- SIOLI, H. 1975. Tropical rivers: the Amazon. Pp. 461–488 in Whitton, B. A. (ed.). *Studies in Ecology 2, River ecology*. University of California Press, Los Angeles.