

Measurement of $t\bar{t}$ production in the τ + jets topology using $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV

V. M. Abazov,³⁵ B. Abbott,⁷³ M. Abolins,⁶² B. S. Acharya,²⁹ M. Adams,⁴⁸ T. Adams,⁴⁶ G. D. Alexeev,³⁵ G. Alkhazov,³⁹ A. Alton,^{61,*} G. Alverson,⁶⁰ G. A. Alves,² L. S. Ancu,³⁴ M. Aoki,⁴⁷ Y. Arnaud,¹⁴ M. Arov,⁵⁷ A. Askew,⁴⁶ B. Åsman,⁴⁰ O. Atramentov,⁶⁵ C. Avila,⁸ J. BackusMayes,⁸⁰ F. Badaud,¹³ L. Bagby,⁴⁷ B. Baldin,⁴⁷ D. V. Bandurin,⁴⁶ S. Banerjee,²⁹ E. Barberis,⁶⁰ P. Baringer,⁵⁵ J. Barreto,² J. F. Bartlett,⁴⁷ U. Bassler,¹⁸ V. Bazterra,⁴⁸ S. Beale,⁶ A. Bean,⁵⁵ M. Begalli,³ M. Begel,⁷¹ C. Belanger-Champagne,⁴⁰ L. Bellantoni,⁴⁷ S. B. Beri,²⁷ G. Bernardi,¹⁷ R. Bernhard,²² I. Bertram,⁴¹ M. Besançon,¹⁸ R. Beuselinck,⁴² V. A. Bezzubov,³⁸ P. C. Bhat,⁴⁷ V. Bhatnagar,²⁷ G. Blazey,⁴⁹ S. Blessing,⁴⁶ K. Bloom,⁶⁴ A. Boehnlein,⁴⁷ D. Boline,⁷⁰ T. A. Bolton,⁵⁶ E. E. Boos,³⁷ G. Borissoy,⁴¹ T. Bose,⁵⁹ A. Brandt,⁷⁶ O. Brandt,²³ R. Brock,⁶² G. Brooijmans,⁶⁸ A. Bross,⁴⁷ D. Brown,¹⁷ J. Brown,¹⁷ X. B. Bu,⁷ D. Buchholz,⁵⁰ M. Buehler,⁷⁹ V. Buescher,²⁴ V. Bunichev,³⁷ S. Burdin,^{41,†} T. H. Burnett,⁸⁰ C. P. Buszello,⁴² B. Calpas,¹⁵ E. Camacho-Pérez,³² M. A. Carrasco-Lizarraga,³² B. C. K. Casey,⁴⁷ H. Castilla-Valdez,³² S. Chakrabarti,⁷⁰ D. Chakraborty,⁴⁹ K. M. Chan,⁵³ A. Chandra,⁷⁸ G. Chen,⁵⁵ S. Chevalier-Théry,¹⁸ D. K. Cho,⁷⁵ S. W. Cho,³¹ S. Choi,³¹ B. Choudhary,²⁸ T. Christoudias,⁴² S. Cihangir,⁴⁷ D. Claes,⁶⁴ J. Clutter,⁵⁵ M. Cooke,⁴⁷ W. E. Cooper,⁴⁷ M. Corcoran,⁷⁸ F. Couderc,¹⁸ M.-C. Cousinou,¹⁵ A. Croc,¹⁸ D. Cutts,⁷⁵ M. Ćwiok,³⁰ A. Das,⁴⁴ G. Davies,⁴² K. De,⁷⁶ S. J. de Jong,³⁴ E. De La Cruz-Burelo,³² F. Déliot,¹⁸ M. Demarteau,⁴⁷ R. Demina,⁶⁹ D. Denisov,⁴⁷ S. P. Denisov,³⁸ S. Desai,⁴⁷ K. DeVaughan,⁶⁴ H. T. Diehl,⁴⁷ M. Diesburg,⁴⁷ A. Dominguez,⁶⁴ T. Dorland,⁸⁰ A. Dubey,²⁸ L. V. Dudko,³⁷ D. Duggan,⁶⁵ A. Duperrin,¹⁵ S. Dutt,²⁷ A. Dyshkant,⁴⁹ M. Eads,⁶⁴ D. Edmunds,⁶² J. Ellison,⁴⁵ V. D. Elvira,⁴⁷ Y. Enari,¹⁷ S. Eno,⁵⁸ H. Evans,⁵¹ A. Evdokimov,⁷¹ V. N. Evdokimov,³⁸ G. Facini,⁶⁰ T. Ferbel,^{58,69} F. Fiedler,²⁴ F. Filthaut,³⁴ W. Fisher,⁶² H. E. Fisk,⁴⁷ M. Fortner,⁴⁹ H. Fox,⁴¹ S. Fuess,⁴⁷ T. Gadfort,⁷¹ A. Garcia-Bellido,⁶⁹ V. Gavrilov,³⁶ P. Gay,¹³ W. Geist,¹⁹ W. Geng,^{15,62} D. Gerbaudo,⁶⁶ C. E. Gerber,⁴⁸ Y. Gershtein,⁶⁵ G. Ginther,^{47,69} G. Golovanov,³⁵ A. Goussiou,⁸⁰ P. D. Grannis,⁷⁰ S. Greder,¹⁹ H. Greenlee,⁴⁷ Z. D. Greenwood,⁵⁷ E. M. Gregores,⁴ G. Grenier,²⁰ Ph. Gris,¹³ J.-F. Grivaz,¹⁶ A. Grohsjean,¹⁸ S. Grünendahl,⁴⁷ M. W. Grünewald,³⁰ F. Guo,⁷⁰ J. Guo,⁷⁰ G. Gutierrez,⁴⁷ P. Gutierrez,⁷³ A. Haas,^{68,‡} S. Hagopian,⁴⁶ J. Haley,⁶⁰ L. Han,⁷ K. Harder,⁴³ A. Harel,⁶⁹ J. M. Hauptman,⁵⁴ J. Hays,⁴² T. Head,⁴³ T. Hebbeker,²¹ D. Hedin,⁴⁹ H. Hegab,⁷⁴ A. P. Heinson,⁴⁵ U. Heintz,⁷⁵ C. Hensel,²³ I. Heredia-De La Cruz,³² K. Herner,⁶¹ G. Hesketh,⁶⁰ M. D. Hildreth,⁵³ R. Hirosky,⁷⁹ T. Hoang,⁴⁶ J. D. Hobbs,⁷⁰ B. Hoeneisen,¹² M. Hohlfield,²⁴ S. Hossain,⁷³ Z. Hubacek,¹⁰ N. Huske,¹⁷ V. Hynek,¹⁰ I. Iashvili,⁶⁷ R. Illingworth,⁴⁷ A. S. Ito,⁴⁷ S. Jabeen,⁷⁵ M. Jaffré,¹⁶ S. Jain,⁶⁷ D. Jamin,¹⁵ R. Jesik,⁴² K. Johns,⁴⁴ M. Johnson,⁴⁷ D. Johnston,⁶⁴ A. Jonckheere,⁴⁷ P. Jonsson,⁴² J. Joshi,²⁷ A. Juste,^{47,§} K. Kaadze,⁵⁶ E. Kajfasz,¹⁵ D. Karmanov,³⁷ P. A. Kasper,⁴⁷ I. Katsanos,⁶⁴ R. Kehoe,⁷⁷ S. Kermiche,¹⁵ N. Khalatyan,⁴⁷ A. Khanov,⁷⁴ A. Kharchilava,⁶⁷ Y. N. Kharzheev,³⁵ D. Khatidze,⁷⁵ M. H. Kirby,⁵⁰ J. M. Kohli,²⁷ A. V. Kozelov,³⁸ J. Kraus,⁶² A. Kumar,⁶⁷ A. Kupco,¹¹ T. Kurča,²⁰ V. A. Kuzmin,³⁷ J. Kvita,⁹ S. Lammers,⁵¹ G. Landsberg,⁷⁵ P. Lebrun,²⁰ H. S. Lee,³¹ S. W. Lee,⁵⁴ W. M. Lee,⁴⁷ J. Lellouch,¹⁷ L. Li,⁴⁵ Q. Z. Li,⁴⁷ S. M. Lietti,⁵ J. K. Lim,³¹ D. Lincoln,⁴⁷ J. Linnemann,⁶² V. V. Lipaev,³⁸ R. Lipton,⁴⁷ Y. Liu,⁷ Z. Liu,⁶ A. Lobodenko,³⁹ M. Lokajicek,¹¹ P. Love,⁴¹ H. J. Lubatti,⁸⁰ R. Luna-Garcia,^{32,||} A. L. Lyon,⁴⁷ A. K. A. Maciel,² D. Mackin,⁷⁸ R. Madar,¹⁸ R. Magaña-Villalba,³² S. Malik,⁶⁴ V. L. Malyshev,³⁵ Y. Maravin,⁵⁶ J. Martínez-Ortega,³² R. McCarthy,⁷⁰ C. L. McGivern,⁵⁵ M. M. Meijer,³⁴ A. Melnitchouk,⁶³ D. Menezes,⁴⁹ P. G. Mercadante,⁴ M. Merkin,³⁷ A. Meyer,²¹ J. Meyer,²³ N. K. Mondal,²⁹ G. S. Muanza,¹⁵ M. Mulhearn,⁷⁹ E. Nagy,¹⁵ M. Naimuddin,²⁸ M. Narain,⁷⁵ R. Nayyar,²⁸ H. A. Neal,⁶¹ J. P. Negret,⁸ P. Neustroev,³⁹ S. F. Novaes,⁵ T. Nunnemann,²⁵ G. Obrant,³⁹ J. Orduna,³² N. Osman,⁴² J. Osta,⁵³ G. J. Otero y Garzón,¹ M. Owen,⁴³ M. Padilla,⁴⁵ M. Pangilinan,⁷⁵ N. Parashar,⁵² V. Parihar,⁷⁵ S. K. Park,³¹ J. Parsons,⁶⁸ R. Partridge,^{75,‡} N. Parua,⁵¹ A. Patwa,⁷¹ B. Penning,⁴⁷ M. Perfilov,³⁷ K. Peters,⁴³ Y. Peters,⁴³ G. Petrillo,⁶⁹ P. Pétrouff,¹⁶ R. Piegai,¹ J. Piper,⁶² M.-A. Pleier,⁷¹ P. L. M. Podesta-Lerma,^{32,¶} V. M. Podstavkov,⁴⁷ M.-E. Pol,² P. Polozov,³⁶ A. V. Popov,³⁸ M. Prewitt,⁷⁸ D. Price,⁵¹ S. Protopopescu,⁷¹ J. Qian,⁶¹ A. Quadt,²³ B. Quinn,⁶³ M. S. Rangel,² K. Ranjan,²⁸ P. N. Ratoff,⁴¹ I. Razumov,³⁸ P. Renkel,⁷⁷ P. Rich,⁴³ M. Rijssenbeek,⁷⁰ I. Ripp-Baudot,¹⁹ F. Rizatdinova,⁷⁴ M. Rominsky,⁴⁷ C. Royon,¹⁸ P. Rubinov,⁴⁷ R. Ruchti,⁵³ G. Safronov,³⁶ G. Sajot,¹⁴ A. Sánchez-Hernández,³² M. P. Sanders,²⁵ B. Sanghi,⁴⁷ A. S. Santos,⁵ G. Savage,⁴⁷ L. Sawyer,⁵⁷ T. Scanlon,⁴² R. D. Schamberger,⁷⁰ Y. Scheglov,³⁹ H. Schellman,⁵⁰ T. Schliephake,²⁶ S. Schlobohm,⁸⁰ C. Schwanenberger,⁴³ R. Schwienhorst,⁶² J. Sekaric,⁵⁵ H. Severini,⁷³ E. Shabalina,²³ V. Shary,¹⁸ A. A. Shchukin,³⁸ R. K. Shivpuri,²⁸ V. Simak,¹⁰ V. Sirotenko,⁴⁷ P. Skubic,⁷³ P. Slattery,⁶⁹ D. Smirnov,⁵³ K. J. Smith,⁶⁷ G. R. Snow,⁶⁴ J. Snow,⁷² S. Snyder,⁷¹ S. Söldner-Rembold,⁴³ L. Sonnenschein,²¹ A. Sopczak,⁴¹ M. Sosebee,⁷⁶ K. Soustruznik,⁹ B. Spurlock,⁷⁶ J. Stark,¹⁴ V. Stolin,³⁶ D. A. Stoyanova,³⁸ E. Strauss,⁷⁰ M. Strauss,⁷³ D. Strom,⁴⁸ L. Stutte,⁴⁷ P. Svoisky,⁷³ M. Takahashi,⁴³ A. Tanasijczuk,¹ W. Taylor,⁶ M. Titov,¹⁸ V. V. Tokmenin,³⁵ D. Tsybychev,⁷⁰ B. Tuchming,¹⁸ C. Tully,⁶⁶ P. M. Tuts,⁶⁸ L. Uvarov,³⁹ S. Uvarov,³⁹ S. Uzunyan,⁴⁹ R. Van Kooten,⁵¹ W. M. van Leeuwen,³³ N. Varelas,⁴⁸ E. W. Varnes,⁴⁴

I. A. Vasilyev,³⁸ P. Verdier,²⁰ L. S. Vertogradov,³⁵ M. Verzocchi,⁴⁷ M. Vesterinen,⁴³ D. Vilanova,¹⁸ P. Vint,⁴² P. Vokac,¹⁰ H. D. Wahl,⁴⁶ M. H. L. S. Wang,⁶⁹ J. Warchol,⁵³ G. Watts,⁸⁰ M. Wayne,⁵³ M. Weber,^{47,**} L. Welty-Rieger,⁵⁰ M. Wetstein,⁵⁸ A. White,⁷⁶ D. Wicke,²⁴ M. R. J. Williams,⁴¹ G. W. Wilson,⁵⁵ S. J. Wimpenny,⁴⁵ M. Wobisch,⁵⁷ D. R. Wood,⁶⁰ T. R. Wyatt,⁴³ Y. Xie,⁴⁷ C. Xu,⁶¹ S. Yacoob,⁵⁰ R. Yamada,⁴⁷ W.-C. Yang,⁴³ T. Yasuda,⁴⁷ Y. A. Yatsunenko,³⁵ Z. Ye,⁴⁷ H. Yin,⁷ K. Yip,⁷¹ H. D. Yoo,⁷⁵ S. W. Youn,⁴⁷ J. Yu,⁷⁶ S. Zelitch,⁷⁹ T. Zhao,⁸⁰ B. Zhou,⁶¹ J. Zhu,⁶¹ M. Zielinski,⁶⁹ D. Zieminska,⁵¹ and L. Zivkovic⁶⁸

(D0 Collaboration)

¹Universidad de Buenos Aires, Buenos Aires, Argentina

²LAFEX, Centro Brasileiro de Pesquisas Físicas, Rio de Janeiro, Brazil

³Universidade do Estado do Rio de Janeiro, Rio de Janeiro, Brazil

⁴Universidade Federal do ABC, Santo André, Brazil

⁵Instituto de Física Teórica, Universidade Estadual Paulista, São Paulo, Brazil

⁶Simon Fraser University, Vancouver, British Columbia, and York University, Toronto, Ontario, Canada

⁷University of Science and Technology of China, Hefei, People's Republic of China

⁸Universidad de los Andes, Bogotá, Colombia

⁹Charles University, Faculty of Mathematics and Physics, Center for Particle Physics, Prague, Czech Republic

¹⁰Czech Technical University in Prague, Prague, Czech Republic

¹¹Center for Particle Physics, Institute of Physics, Academy of Sciences of the Czech Republic, Prague, Czech Republic

¹²Universidad San Francisco de Quito, Quito, Ecuador

¹³LPC, Université Blaise Pascal, CNRS/IN2P3, Clermont, France

¹⁴LPSC, Université Joseph Fourier Grenoble 1, CNRS/IN2P3, Institut National Polytechnique de Grenoble, Grenoble, France

¹⁵CPPM, Aix-Marseille Université, CNRS/IN2P3, Marseille, France

¹⁶LAL, Université Paris-Sud, CNRS/IN2P3, Orsay, France

¹⁷LPNHE, Universités Paris VI and VII, CNRS/IN2P3, Paris, France

¹⁸CEA, Ifu, SPP, Saclay, France

¹⁹IPHC, Université de Strasbourg, CNRS/IN2P3, Strasbourg, France

²⁰IPNL, Université Lyon 1, CNRS/IN2P3, Villeurbanne, France, and Université de Lyon, Lyon, France

²¹III. Physikalisches Institut A, RWTH Aachen University, Aachen, Germany

²²Physikalisches Institut, Universität Freiburg, Freiburg, Germany

²³II. Physikalisches Institut, Georg-August-Universität Göttingen, Göttingen, Germany

²⁴Institut für Physik, Universität Mainz, Mainz, Germany

²⁵Ludwig-Maximilians-Universität München, München, Germany

²⁶Fachbereich Physik, Bergische Universität Wuppertal, Wuppertal, Germany

²⁷Panjab University, Chandigarh, India

²⁸Delhi University, Delhi, India

²⁹Tata Institute of Fundamental Research, Mumbai, India

³⁰University College Dublin, Dublin, Ireland

³¹Korea Detector Laboratory, Korea University, Seoul, Korea

³²CINVESTAV, Mexico City, Mexico

³³FOM-Institute NIKHEF, and University of Amsterdam/NIKHEF, Amsterdam, The Netherlands

³⁴Radboud University Nijmegen/NIKHEF, Nijmegen, The Netherlands

³⁵Joint Institute for Nuclear Research, Dubna, Russia

³⁶Institute for Theoretical and Experimental Physics, Moscow, Russia

³⁷Moscow State University, Moscow, Russia

³⁸Institute for High Energy Physics, Protvino, Russia

³⁹Petersburg Nuclear Physics Institute, St. Petersburg, Russia

⁴⁰Stockholm University, Stockholm, and Uppsala University, Uppsala, Sweden

⁴¹Lancaster University, Lancaster LA1 4YB, United Kingdom

⁴²Imperial College London, London SW7 2AZ, United Kingdom

⁴³The University of Manchester, Manchester M13 9PL, United Kingdom

⁴⁴University of Arizona, Tucson, Arizona 85721, USA

⁴⁵University of California Riverside, Riverside, California 92521, USA

⁴⁶Florida State University, Tallahassee, Florida 32306, USA

⁴⁷Fermi National Accelerator Laboratory, Batavia, Illinois 60510, USA

⁴⁸University of Illinois at Chicago, Chicago, Illinois 60607, USA

⁴⁹Northern Illinois University, DeKalb, Illinois 60115, USA

⁵⁰Northwestern University, Evanston, Illinois 60208, USA

- ⁵¹Indiana University, Bloomington, Indiana 47405, USA
⁵²Purdue University Calumet, Hammond, Indiana 46323, USA
⁵³University of Notre Dame, Notre Dame, Indiana 46556, USA
⁵⁴Iowa State University, Ames, Iowa 50011, USA
⁵⁵University of Kansas, Lawrence, Kansas 66045, USA
⁵⁶Kansas State University, Manhattan, Kansas 66506, USA
⁵⁷Louisiana Tech University, Ruston, Louisiana 71272, USA
⁵⁸University of Maryland, College Park, Maryland 20742, USA
⁵⁹Boston University, Boston, Massachusetts 02215, USA
⁶⁰Northeastern University, Boston, Massachusetts 02115, USA
⁶¹University of Michigan, Ann Arbor, Michigan 48109, USA
⁶²Michigan State University, East Lansing, Michigan 48824, USA
⁶³University of Mississippi, University, Mississippi 38677, USA
⁶⁴University of Nebraska, Lincoln, Nebraska 68588, USA
⁶⁵Rutgers University, Piscataway, New Jersey 08855, USA
⁶⁶Princeton University, Princeton, New Jersey 08544, USA
⁶⁷State University of New York, Buffalo, New York 14260, USA
⁶⁸Columbia University, New York, New York 10027, USA
⁶⁹University of Rochester, Rochester, New York 14627, USA
⁷⁰State University of New York, Stony Brook, New York 11794, USA
⁷¹Brookhaven National Laboratory, Upton, New York 11973, USA
⁷²Langston University, Langston, Oklahoma 73050, USA
⁷³University of Oklahoma, Norman, Oklahoma 73019, USA
⁷⁴Oklahoma State University, Stillwater, Oklahoma 74078, USA
⁷⁵Brown University, Providence, Rhode Island 02912, USA
⁷⁶University of Texas, Arlington, Texas 76019, USA
⁷⁷Southern Methodist University, Dallas, Texas 75275, USA
⁷⁸Rice University, Houston, Texas 77005, USA
⁷⁹University of Virginia, Charlottesville, Virginia 22901, USA
⁸⁰University of Washington, Seattle, Washington 98195, USA
(Received 25 August 2010; published 27 October 2010)

We present a measurement of the $t\bar{t}$ production cross section multiplied by the branching ratio to tau lepton decaying semihadronically (τ_h) plus jets, $\sigma(p\bar{p} \rightarrow t\bar{t} + X) \cdot \text{BR}(t\bar{t} \rightarrow \tau_h + \text{jets})$, at a center of mass energy $\sqrt{s} = 1.96$ TeV using 1 fb^{-1} of integrated luminosity collected with the D0 detector. Assuming a top quark mass of 170 GeV, we measure $\sigma_{t\bar{t}} \cdot \text{BR}_{\tau_h+j} = 0.60^{+0.23}_{-0.22}(\text{stat})^{+0.15}_{-0.14}(\text{syst}) \pm 0.04(\text{lumi})$ pb. In addition, we extract the $t\bar{t}$ production cross section using the $t\bar{t} \rightarrow \tau_h + \text{jets}$ topology, with the result $\sigma_{t\bar{t}} = 6.9^{+1.2}_{-1.2}(\text{stat})^{+0.8}_{-0.7}(\text{syst}) \pm 0.4(\text{lumi})$ pb. These findings are in good agreement with standard model predictions and measurements performed using other top quark decay channels.

DOI: 10.1103/PhysRevD.82.071102

PACS numbers: 13.85.Lg, 13.85.Ni, 13.85.Qk, 14.65.Ha

The decay $t \rightarrow Wb \rightarrow \tau\nu_\tau b$ provides a unique laboratory to investigate the properties of the third generation fermions—the top (t) and bottom (b) quarks, the tau lepton (τ), and the tau neutrino (ν_τ)—in a single process. In the standard model (SM), the t quark branching ratio (BR) to a W boson and a b quark is $\approx 100\%$, and the final state is determined by the SM BR of the W boson. Since the t is the heaviest quark and the τ the heaviest lepton, any non-SM

mass- or flavor-dependent couplings could change the t quark decay rate into final states with τ leptons. Therefore, any deviation in the BR of $t \rightarrow \tau\nu_\tau b$ from that predicted by the SM can be an indication of non-SM physics. For example, in the Type 2 two-Higgs doublet model [1], such as required by the minimal supersymmetric standard model [2], the t quark can have a significant BR to a charged Higgs boson (H^\pm) and a b quark if $m_{H^\pm} < m_t - m_b$. For large values of $\tan\beta$, the ratio of the vacuum expectation values of the two-Higgs doublets, the charged Higgs boson preferentially decays to $\tau\nu_\tau$, thereby increasing the BR of $t \rightarrow \tau\nu_\tau b$ relative to the SM expectation and leading to a larger measured $\sigma(p\bar{p} \rightarrow t\bar{t} + X) \cdot \text{BR}(t\bar{t} \rightarrow \tau + \text{jets})$ compared to the value expected from SM assumptions for the BRs and the production cross section [3–5]. Other possible non-SM processes that can enhance the t quark to τ lepton BR are R -parity violating decays of

*Visitor from Augustana College, Sioux Falls, SD, USA.

†Visitor from The University of Liverpool, Liverpool, UK.

‡Visitor from SLAC, Menlo Park, CA, USA.

§Visitor from ICREA/IFAE, Barcelona, Spain.

||Visitor from Centro de Investigacion en Computacion - IPN, Mexico City, Mexico.

¶Visitor from ECFM, Universidad Autonoma de Sinaloa, Culiacán, Mexico.

**Visitor from Universität Bern, Bern, Switzerland.

the t quark in supersymmetric models [6] and new Z' bosons with nonuniversal couplings [7].

In this article, we present the first measurement of $t\bar{t}$ production in the τ + jets final state using a data sample corresponding to an integrated luminosity of 1 fb^{-1} collected with the D0 detector [8] at the Fermilab Tevatron $p\bar{p}$ Collider operating at a center of mass energy $\sqrt{s} = 1.96 \text{ TeV}$. This measurement uses semihadronic τ lepton decays, with $\text{BR} \approx 65\%$, as secondary electrons and muons from τ lepton decays are difficult to distinguish from primary electrons and muons resulting from W decays. Previous measurements of $t\bar{t}$ production using τ leptons in the final state have been performed by the D0 [9] and CDF [10] collaborations in the $\tau_h + \ell$ channel, where τ_h represents semihadronic τ lepton decay modes and ℓ represents either an electron or a muon.

We apply the following preselection requirements: events must satisfy a multijet trigger requiring at least four jets; this is the same trigger used in the $t\bar{t}$ cross section measurement in the all-hadronic decay mode [11]. Reconstructed events are required to have missing transverse energy $E_T \geq 15 \text{ GeV}$ and E_T significance > 3 , where the E_T significance is a measure of the likelihood that the E_T arises from physical sources rather than fluctuations in the measurement of the energies of the physics objects (jets, muons, electrons, and unclustered energy) [12]. Each event must also have at least four reconstructed jets with pseudorapidity $|\eta| < 2.5$ and transverse momentum $p_T > 15 \text{ GeV}$ using an iterative jet cone algorithm [13] with a cone size $\Delta\mathcal{R} = \sqrt{(\Delta\eta)^2 + (\Delta\phi)^2} = 0.5$ [14]. The jet energies are corrected for the energy response of the calorimeter, the cone size, multiple $p\bar{p}$ interactions, event pileup, and calorimeter noise [15]. At least one jet is required to have $p_T > 35 \text{ GeV}$, and at least two jets are required to have $p_T > 25 \text{ GeV}$. Each event is also required to have at least one τ_h candidate with $p_T > 10 \text{ GeV}$, $|\eta| < 2.5$, and tau neural network output, $NN_\tau > 0.3$ [16]. Finally, to ensure this analysis is statistically independent of other D0 $t\bar{t}$ cross section measurements so that it can be included in a combined cross section measurement, events satisfying the requirements of the $t\bar{t} \rightarrow e(\mu) + \text{jets}$ channel [17], which include an isolated electron (muon) with $p_T > 20 \text{ GeV}$, are rejected, as are events satisfying the requirements of the $t\bar{t}$ cross section measurement in the all-hadronic channel [11].

A semihadronic τ lepton candidate is a calorimeter cluster of cone size $\Delta\mathcal{R} = 0.5$ that includes any subclusters that might be present with $E > 800 \text{ MeV}$ constructed from cells in the electromagnetic (EM) section of the calorimeter and the associated tracks with $p_T > 1.5 \text{ GeV}$ in a cone $\Delta\mathcal{R} = 0.3$ contained within the calorimeter cluster. These τ candidates are classified according to one of three types based on the number of tracks and activity in the EM calorimeter, motivated by the semihadronic τ lepton decays: (1) $\tau^\pm \rightarrow \pi^\pm \nu_\tau$, (2) $\tau^\pm \rightarrow \pi^\pm \pi^0 \nu_\tau$,

(3) $\tau^\pm \rightarrow \pi^\pm \pi^\pm \pi^\mp (\pi^0) \nu_\tau$. We define the three tau-types as follows: a single track with no EM subclusters (tau-type 1); a single track and ≥ 1 EM subclusters (tau-type 2); and at least two tracks and ≥ 0 EM subclusters (tau-type 3).

To further reduce the number of quark and gluon jets reconstructed as τ leptons, we train separate neural networks for each τ_h lepton decay type to improve the discrimination of τ lepton candidates from the jet background. The input variables to NN_τ are chosen to be minimally dependent on the τ lepton energy and to exploit the low track multiplicity and the narrow width of the calorimeter cluster produced by τ leptons decaying semihadronically, the low mass of the τ lepton, and the differences in longitudinal and transverse shower shapes between τ leptons and jets [16]. A total of 12 NN_τ input variables are used to characterize the presence and properties of τ_h leptons, with seven of these variables in common for all three tau-types. The 12 variables are classified as follows: isolation variables, shower shape variables, and correlation variables between the calorimeter cluster and the associated charged particle tracks. Each NN_τ is trained on $Z \rightarrow \tau^+ \tau^-$ Monte Carlo (MC) events for signal and jets from data, where a jet and a nonisolated muon are back-to-back in ϕ , for background. These are the same training samples used in Ref. [18].

To measure the number of $t\bar{t} \rightarrow \tau_h + \text{jets}$ signal events in data, the physics and instrumental backgrounds must be determined. The main physics backgrounds are $W + \text{jets}$ events, where the W boson decays to a τ lepton, and to a smaller extent $Z + \text{jets}$ events, where the Z boson decays to a pair of τ leptons with one misidentified as a jet and the E_T is due to the neutrinos from the decays of the τ leptons. The main instrumental background is multijet production where a jet is misidentified as a τ lepton and the energy is mismeasured leading to a net E_T .

The preselection efficiencies and SM BRs for $t\bar{t}$ to final states with leptons [19] are given in Table I. These, as well as the final efficiencies, are calculated using a MC simulation of the experiment. The $t\bar{t}$ signal with leptons in the final state and $W(Z) + \text{jets}$ background are simulated using the ALPGEN 1.2 [20] matrix element generator assuming a t quark mass of 170 GeV and using the CTEQ6L1 [21] parton

TABLE I. A summary of the SM BRs of the various $t\bar{t}$ subprocesses and the preselection efficiencies, where the uncertainties are derived from MC statistics. The leptonic τ lepton decays are included in the e and μ channels, and l^\pm represents an e , μ , or τ lepton.

	BR (%)	$\epsilon_{\text{preselection}}$ (%)
$t\bar{t} \rightarrow \tau_h + \text{jets}$	9.75	40.5 ± 0.2
$t\bar{t} \rightarrow e + \text{jets}$	17.7	17.0 ± 0.2
$t\bar{t} \rightarrow \mu + \text{jets}$	17.6	11.1 ± 0.1
$t\bar{t} \rightarrow l^+ l^- + \text{jets}$	11.1	4.04 ± 0.03

distribution function set. These events are then processed through PYTHIA 6.2 [22] to simulate parton showering, fragmentation, hadronization, and decays of short lived particles, except for b hadrons and τ leptons. EVTGEN [23] is used to model the decays of b hadrons, while τ leptons are decayed using TAUOLA [24]. To avoid double counting final states generated by the leading-order parton-level calculation of ALPGEN and the parton-level shower evolution of PYTHIA, a matching algorithm is used [25]. The generated events are then processed through the GEANT-based [26] simulation of the D0 detector providing tracking hits, calorimeter cell energies and muon hit information. The same reconstruction algorithm is applied to data and simulated events.

The preselected data sample is used to extract the signal and to study the multijet background after additional selection criteria are applied. To extract the signal sample, we require $NN_\tau > 0.95$. The selected events are then separated on the basis of tau-type according to the τ lepton candidate with the highest value of NN_τ . This is done primarily to separate tau-type 3 events from the tau-type 1 and 2 events, since the former has a much higher misidentification rate and thus result in larger uncertainties on the $t\bar{t}$ cross section. In addition, we require that each event have at least one identified b jet using the b -tag neural network (NN_b) with the requirement $NN_b > 0.775$. The NN_b uses nine input variables that characterize the presence and properties of secondary vertices and track impact parameters within the jet [27]. The efficiencies of these selections are shown in Table II.

The expected fraction of $t\bar{t}$ events in the signal sample is $\approx 15\%$ for tau-type 1 and 2, and $\approx 3\%$ for tau-type 3 assuming $\sigma_{t\bar{t}} = 6.9$ pb as measured in this analysis. In addition, the signal sample contains $W(Z) +$ jets and multijet background events that must be subtracted. The $W(Z) +$ jets contamination is determined using MC events, while the multijet background is determined from data. We start with the preselected sample and apply a loose τ lepton veto, $NN_\tau < 0.9$. Using MC events, we expect that the resulting sample contains $< 2\%$ $t\bar{t} \rightarrow \tau_h +$ jets events and $< 3\%$ $W(Z) +$ jets events, and therefore provides a good representation of the multijet background. To further improve the modeling, the $W(Z) +$ jets expectation is subtracted from the multijet background data sample.

The numbers of signal and background events are extracted from the final selected sample using a neural network (NN_{sb}) event discriminant with the following input variables: (1) the scalar sum of the p_T of all jets and the τ lepton candidate in the event; (2) the aplanarity [28]; (3) the E_T significance; (4) the invariant mass of all jets and the τ lepton candidate in the event; and (5) a χ^2 representing how well the 2 and 3 jet invariant masses agree with values expected for hadronic t quark decays, $\chi^2 = (M_{3\text{jet}} - m_t)/\sigma_t^2 + (M_{2\text{jet}} - m_W)^2/\sigma_W^2$, with $M_{3\text{jet}}$ ($M_{2\text{jet}}$) being the 2 (3) jet invariant mass, $m_t = 170$ GeV, $\sigma_t = 45$ GeV, and $m_W = 80$ GeV, $\sigma_W = 10$ GeV are the mass and its resolution in the all-hadronic final state for the t quark and W boson, respectively. The jet combination minimizing the χ^2 is used. The NN_{sb} is trained using a generated $t\bar{t} \rightarrow \tau_h +$ jets MC sample for signal and half the multijet data sample for background.

We apply the trained NN_{sb} to the signal data sample, the remaining half of the multijet sample, a $t\bar{t}$ MC sample with leptons in the final state that is independent of the NN_{sb} training sample, and a $W(Z) +$ jets MC sample. The application of NN_{sb} on the multijet and MC samples is used to generate templates, as shown in Fig. 1, that are used to determine the fraction of $t\bar{t}$ and multijet events using a negative log-likelihood fit. The normalization of the $W(Z) +$ jets MC sample is derived by scaling the $W(Z)$ transverse (dilepton) mass distribution to data. The normalization for $t\bar{t} \rightarrow e(\mu) +$ jets is fixed to the theoretical cross section [5] and BRs.

The number of $t\bar{t} \rightarrow \tau_h +$ jets events extracted from the fit to data are $25.1^{+11.2}_{-10.5}$ (stat) and $18.0^{+11.3}_{-10.5}$ (stat) for channels with tau-types 1 and 2 together, and with tau-type 3, respectively. The fitted numbers of the multijet background events are $336.4^{+11.2}_{-10.5}$ (stat) and $1083.2^{+11.3}_{-10.3}$ (stat), for the two channels, respectively. The numbers of $t\bar{t}$ events are comparable to the expected values given in Table III.

To minimize the statistical uncertainty of the measurement of $\sigma(p\bar{p} \rightarrow t\bar{t} + X) \cdot \text{BR}(t\bar{t} \rightarrow \tau_h + \text{jets})$, which we denote as $\sigma_{t\bar{t}} \cdot \text{BR}_{\tau_h+j}$, we fit the entire NN_{sb} output distribution rather than counting events above a given value. The value of $\sigma_{t\bar{t}} \cdot \text{BR}_{\tau_h+j}$ and the fraction of multijet background in the sample are obtained from a negative log-likelihood fit to the NN_{sb} distributions for tau-types 1 and 2 and tau-type 3, independently:

TABLE II. The efficiencies for the tight τ lepton candidate ($NN_\tau > 0.95$) and b -tagging selections for tau-type 1 and 2, and tau-type 3 channels. The uncertainties are based on MC statistics.

	Tau-types 1 and 2 Trigger (%)	Tau-types 1 and 2 $NN_\tau > 0.95$ (%)	Tau-types 1 and 2 b -tag (%)	Tau-type 3 Trigger (%)	Tau-type 3 $NN_\tau > 0.95$ (%)	Tau-type 3 b -tag (%)
$t\bar{t} \rightarrow \tau_h +$ jets	$74.8^{+1.7}_{-0.1}$	23.7 ± 0.3	$60.1^{+2.8}_{-2.7}$	$73.6^{+1.6}_{-0.1}$	19.4 ± 0.2	$59.9^{+2.8}_{-2.7}$
$t\bar{t} \rightarrow e +$ jets	$69.9^{+1.5}_{-0.1}$	33.1 ± 0.4	$58.7^{+2.8}_{-2.7}$	$66.0^{+1.5}_{-0.1}$	8.1 ± 0.2	$58.9^{+2.8}_{-2.7}$
$t\bar{t} \rightarrow \mu +$ jets	$65.9^{+1.5}_{-0.1}$	3.8 ± 0.1	$60.3^{+2.8}_{-2.7}$	$66.1^{+1.4}_{-0.1}$	7.7 ± 0.2	$59.0^{+2.8}_{-2.7}$
$t\bar{t} \rightarrow l^+ l^- +$ jets	$50.5^{+1.1}_{-0.2}$	43.7 ± 0.4	$60.2^{+2.8}_{-2.7}$	$50.7^{+1.1}_{-0.2}$	20.6 ± 0.3	$61.4^{+2.9}_{-2.8}$




FIG. 1 (color online). The output of NN_{sb} for (a) the tau-type 1 and 2 channel, (b) the tau-type 3 channel. The χ^2 per degree of freedom between data and templates is 0.6 for (a) and 0.5 for (b).

$$L(\sigma_{t\bar{t}}, \tilde{N}_i, N_i^{\text{obs}}) = -\log\left(\prod_i \frac{\tilde{N}_i^{N_i^{\text{obs}}}}{N_i^{\text{obs}}!} e^{-\tilde{N}_i}\right), \quad (1)$$

where $\tilde{N}_i = \sigma_{t\bar{t}} \times \sum_j \epsilon_{t\bar{t}(j)}^i \times \text{BR}_{t\bar{t}(j)} \times \mathcal{L} + N_{\text{bkg},i}$ is the expected number of events in the i^{th} bin of the NN_{sb} histogram for a given $\sigma_{t\bar{t}}$, with integrated luminosity \mathcal{L} , number of background events $N_{\text{bkg},i}$, and the efficiency (BR) for the j^{th} $t\bar{t}$ leptonic channel $\epsilon_{t\bar{t}(j)}$ ($\text{BR}_{t\bar{t}(j)}$), and N_i^{obs} is the observed number of events in the i^{th} bin.

The measured value of $\sigma_{t\bar{t}} \cdot \text{BR}_{\tau_h+j}$ is

$$0.60_{-0.22}^{+0.23}(\text{stat})_{-0.14}^{+0.15}(\text{syst}) \pm 0.04(\text{lumi}) \text{ pb},$$

where we combine the tau-type 1 and 2 measurement with the tau-type 3 measurement. Using the theoretical cross section $\sigma_{t\bar{t}} = 8.06_{-0.73}^{+0.52}$ pb for $m_t = 170$ GeV from

TABLE III. Expected event yields in the two analysis channels assuming the measured $t\bar{t}$ production cross section of 6.9 pb. The uncertainties are derived from MC statistics.

	Tau-type 1 and 2	Tau-type 3
$t\bar{t} \rightarrow \tau_h + \text{jets}$	27.6 ± 0.4	22.1 ± 0.3
$t\bar{t} \rightarrow e + \text{jets}$	26.3 ± 0.4	5.9 ± 0.2
$t\bar{t} \rightarrow \mu + \text{jets}$	2.0 ± 0.1	3.7 ± 0.1
$t\bar{t} \rightarrow l^+ l^- + \text{jets}$	4.1 ± 0.1	2.0 ± 0.1
Total $t\bar{t} \rightarrow \text{leptons}$	61.3 ± 0.6	34.4 ± 0.4
$W + \text{jets}$	13.5 ± 0.3	5.9 ± 0.2
$Z + \text{jets}$	3.4 ± 0.4	1.9 ± 0.1

Ref. [5], we measure $\text{BR}_{\tau_h+j} = 0.074_{-0.027}^{+0.029}$ which is consistent with the SM value given in Table I.

Table IV summarizes the systematic uncertainties on $\sigma_{t\bar{t}} \cdot \text{BR}_{\tau_h+j}$. These are calculated by varying the source by plus and minus one standard deviation, and propagating the uncertainty to the final $\sigma_{t\bar{t}} \cdot \text{BR}_{\tau_h+j}$. The jet energy corrections account for the effect of the jet energy scale and resolution. Jet identification takes account of the difference in the jet finding efficiency in data and MC. The b -tagging entry accounts for the systematic uncertainties on its efficiency. The τ lepton identification uncertainty is derived by fluctuating the value of each input variable within its statistical uncertainty and observing its effect on the NN_{τ} output. The trigger category accounts for the uncertainty in the multijet trigger turn-on and also takes into account the possibility that a multijet event with a τ lepton can have a different trigger turn-on. Multijet modeling accounts for the uncertainty of the multijet sample to model the $t\bar{t} \rightarrow \tau_h + \text{jets}$ background and its limited statistics. The category MC modeling accounts for the $W + \text{jets}$ modeling, the uncertainty in the scale factor both for light flavor jets and heavy flavor jets, and the parton distribution function uncertainty. The $t\bar{t}$ cross section systematic uncertainty represents the effect of the normalization of the non-tau lepton $t\bar{t}$ background, which is normalized to the theoretical value of the cross section. In addition to the sources listed in Table IV, there is a $\pm 6.1\%$ uncertainty in the luminosity measurement [29].

In addition, we present the combined measurement of the production cross section for $t\bar{t}$ using all measured $t\bar{t}$ channels with leptons in the final state listed in Table III that satisfy the selection criteria described above. We repeat the negative log-likelihood fit for the number of $t\bar{t}$ signal and multijet background events fixing the $t\bar{t}$ BRs to their SM values, but this time fit for all $t\bar{t}$ channels arriving at 60.5 ± 11.8 (stat) events and 24.0 ± 11.4 (stat) events for channels with tau-types 1 and 2 and with tau-type 3 characteristics, respectively. The fitted multijet backgrounds in this case are 336.7 ± 11.8 (stat) events and 1083.2 ± 11.4 (stat) events, for the two channels, respectively. The production cross section is calculated using the negative log-likelihood defined in Eq. (1) for tau-types 1 and 2 and tau-type 3 separately. The two cross sections are then combined to give

$$\sigma_{t\bar{t}} = 6.9_{-1.2}^{+1.2}(\text{stat})_{-0.7}^{+0.8}(\text{syst}) \pm 0.4(\text{lumi}) \text{ pb}.$$

To estimate the dependence on m_t , we reevaluate the efficiencies and templates using $m_t = 175$ GeV and find

$$\sigma_{t\bar{t}} = 6.3_{-1.1}^{+1.2}(\text{stat}) \pm_{-0.7}^{+0.7}(\text{syst}) \pm 0.4(\text{lumi}) \text{ pb}.$$

In summary, we have performed a measurement of $\sigma_{t\bar{t}} \cdot \text{BR}_{\tau_h+j} = 0.60_{-0.26}^{+0.28}$ pb and, using the theoretical $t\bar{t}$ production cross section, extracted $\text{BR}_{\tau_h+j} = 0.074_{-0.027}^{+0.029}$, which agrees with the SM expectation. In addition, we have performed a measurement of the $p\bar{p} \rightarrow t\bar{t} + X$

TABLE IV. Systematic uncertainties on $\sigma_{t\bar{t}} \cdot \text{BR}_{\tau_h+j}$ (in pb) as measured for the $t\bar{t} \rightarrow \tau_h + \text{jets}$ channel.

Source	$\tau_h + \text{jets}$ (types 1 and 2)		$\tau_h + \text{jets}$ (type 3)		Combined	
Jet energy corrections	-0.078	+0.081	-0.047	+0.047	-0.068	+0.069
Jet identification	-0.019	+0.019	-0.012	+0.012	-0.016	+0.016
b tagging	-0.074	+0.084	-0.035	+0.041	-0.060	+0.068
Tau identification	-0.035	+0.035	-0.020	+0.021	-0.029	+0.029
Trigger	-0.002	+0.053	-0.000	+0.027	-0.002	+0.043
Multijet modeling	-0.090	+0.090	-0.169	+0.169	-0.083	+0.083
MC modeling	-0.028	+0.028	-0.012	+0.013	-0.023	+0.022
$t\bar{t}$ cross section	-0.064	+0.068	-0.029	+0.030	-0.052	+0.055
Total systematic uncertainty	-0.16	+0.15	-0.18	+0.15	-0.14	+0.15

production cross section, $\sigma_{t\bar{t}} = 6.9_{-1.4}^{+1.5}$ pb, using the $t\bar{t} \rightarrow \tau_h + \text{jets}$ topology. The measurement is in agreement with the SM [3–5] and previous experimental measurements using other $t\bar{t}$ channels [19] at the Tevatron.

We thank the staffs at Fermilab and collaborating institutions, and acknowledge support from the DOE and NSF (USA); CEA and CNRS/IN2P3 (France); FASI, Rosatom and RFBR (Russia); CNPq, FAPERJ, FAPESP, and

FUNDUNESP (Brazil); DAE and DST (India); Colciencias (Colombia); CONACyT (Mexico); KRF and KOSEF (Korea); CONICET and UBACyT (Argentina); FOM (The Netherlands); STFC and the Royal Society (United Kingdom); MSMT and GACR (Czech Republic); CRC Program and NSERC (Canada); BMBF and DFG (Germany); SFI (Ireland); The Swedish Research Council (Sweden); and CAS and CNSF (China).

-
- [1] V. Barger and R.J.N. Phillips, *Phys. Rev. D* **41**, 884 (1990).
- [2] J. Guasch and J. Sola, *Phys. Lett. B* **416**, 353 (1998).
- [3] M. Cacciari *et al.*, *J. High Energy Phys.* 09 (2008) 127.
- [4] N. Kidonakis and R. Vogt, *Phys. Rev. D* **78**, 074005 (2008).
- [5] S. Moch and P. Uwer, *Phys. Rev. D* **78**, 034003 (2008).
- [6] T. Han and M.B. Magro, *Phys. Lett. B* **476**, 79 (2000).
- [7] C. Yue, H. Zong, and L. Liu, *Mod. Phys. Lett. A* **18**, 2187 (2003).
- [8] V.M. Abazov *et al.* (D0 Collaboration), *Nucl. Instrum. Methods Phys. Res., Sect. A* **565**, 463 (2006).
- [9] V.M. Abazov *et al.* (D0 Collaboration), *Phys. Lett. B* **679**, 177 (2009).
- [10] A. Abulencia *et al.* (CDF Collaboration), *Phys. Lett. B* **639**, 172 (2006).
- [11] V.M. Abazov *et al.* (D0 Collaboration), *Phys. Rev. D* **82**, 032002 (2010).
- [12] A. Schwartzman, Report No. FERMILAB-THESIS-2004-2.
- [13] G.C. Blazey *et al.*, in *Proceedings of the Workshop: QCD and Weak Boson Physics in Run II*, edited by U. Baur, R.K. Ellis, and D. Zeppenfeld (Fermilab Report No. Fermilab-Pub-00/297, 2000).
- [14] The D0 coordinate system has the positive z axis along the proton beamline, and $z = 0$ at the center of the detector. The polar and azimuthal angles are denoted as θ and ϕ , respectively. The pseudorapidity is defined as $\eta = -\ln(\tan\frac{\theta}{2})$.
- [15] J. Hegeman, *J. Phys. Conf. Ser.* **160**, 012024 (2009).
- [16] V.M. Abazov *et al.* (D0 Collaboration), *Phys. Lett. B* **670**, 292 (2009); C.F. Galea, *Acta Phys. Pol. B* **38**, 769 (2007).
- [17] V.M. Abazov *et al.* (D0 Collaboration), *Phys. Rev. D* **76**, 092007 (2007).
- [18] V.M. Abazov *et al.* (D0 Collaboration), *Phys. Lett. B* **670**, 292 (2009).
- [19] C. Amsler *et al.* (Particle Data Group), *Phys. Lett. B* **667**, 1 (2008).
- [20] M.L. Mangano *et al.*, *J. High Energy Phys.* 07 (2003) 001.
- [21] J. Pumplin *et al.*, *J. High Energy Phys.* 07 (2002) 012; D. Stump *et al.*, *J. High Energy Phys.* 10 (2003) 046.
- [22] T. Sjöstrand *et al.*, *Comput. Phys. Commun.* **135**, 238 (2001).
- [23] D.J. Lange, *Nucl. Instrum. Methods Phys. Res., Sect. A* **462**, 152 (2001).
- [24] S. Jadach, Z. Was, R. Decker, and J.H. Kuehn, *Comput. Phys. Commun.* **76**, 361 (1993).
- [25] S. Höche *et al.*, arXiv:hep-ph/0602031.
- [26] R. Brun and F. Carminati, CERN Report No. W5013 (unpublished).
- [27] V.M. Abazov *et al.* (D0 Collaboration), *Nucl. Instrum. Methods Phys. Res., Sect. AA&E* **620**, 490 (2010).
- [28] The aplanarity is $3/2\lambda_3$, with λ_3 being the smallest eigenvalue of the momentum tensor $M^{\alpha\beta} = \sum_i p_i^\alpha p_i^\beta / \sum_i |\vec{p}_i|^2$, where i runs over the number of jets and the τ lepton candidate, and $\alpha, \beta = 1, 2, 3$ specifies the three spatial components of the momentum.
- [29] T. Andeen *et al.*, Report No. FERMILAB-TM-2365, 2007.