



## Molecular characterization of fire ants, *Solenopsis* spp., from Brazil based on analysis of mtDNA gene cytochrome oxidase I

Cintia Martins<sup>1a</sup>, Rodrigo Fernando de Souza<sup>2b</sup>, Odair Correa Bueno<sup>3c</sup>

<sup>1</sup>Universidade Federal do Piauí, Campus de Parnaíba, Avenida São Sebastião, 2819, 64202-020, Parnaíba, Piauí, Brazil

<sup>2</sup>UNESP Universidade Estadual Paulista Julio de Mesquita Filho/ CEIS Centro de Estudos de Insetos Sociais. Avenida 24A, 1515. CEP 13506-900 Rio Claro, São Paulo, Brazil

<sup>3</sup>UNESP Universidade Estadual Paulista Julio de Mesquita Filho/ CEIS Centro de Estudos de Insetos Sociais. Avenida 24A, 1515. CEP 13506-900 Rio Claro, São Paulo, Brazil

### Abstract

Species from the *Solenopsis saevissima* (Smith) (Hymenoptera: Formicidae) species group are native to South America and have a cosmopolitan distribution because they have been accidentally introduced in many countries around the world. In Brazil, they have a wide distribution, including urban areas. The present study was conducted to investigate the characterization of *Solenopsis* genus populations associated with urban/human interference sites in Brazil by analyzing the mitochondrial gene cytochrome oxidase I and estimating the degree of relatedness of these populations to make inferences about their phylogeny and also observe the patterns of mitochondrial haplotype (mitotype) distribution across their range. The results revealed complete geographical coherence and polyphyly for the *Solenopsis invicta* Buren and *Solenopsis saevissima* species groups, which confirms the diversity of the genera. It also suggests the possibility that reproductively-isolated populations occur, resulting in the evolutionary process of speciation. No predominant haplotype was found in the populations analyzed, but some were more prevalent.

**Abbreviations:** COI, cytochrome oxidase I

**Keywords:** mitochondrial DNA, phylogeny, *Solenopsis invicta*, *Solenopsis saevissima*

**Correspondence:** <sup>a</sup> [martins.c@ufpi.edu.br](mailto:martins.c@ufpi.edu.br), <sup>b</sup> [souza\\_bio@yahoo.com.br](mailto:souza_bio@yahoo.com.br), <sup>c</sup> [odaircb@rc.unesp.br](mailto:odaircb@rc.unesp.br)

**Received:** 7 October 2011 **Accepted:** 2 December 2013 **Published:** 10 April 2014

**Editor:** Henry Hagedorn was editor of this paper.

**Copyright:** This is an open access paper. We use the Creative Commons Attribution 3.0 license that permits unrestricted use, provided that the paper is properly attributed.

**ISSN:** 1536-2442 | Vol. 14, Number 50

#### Cite this paper as:

Martins C, Fernando de Souza R, Bueno OC. 2014. Molecular characterization of fire ants, *Solenopsis* spp., from Brazil based on analysis of mtDNA gene cytochrome oxidase I. *Journal of Insect Science* 14:50. Available online: <http://www.insectscience.org/14.50>

## Introduction

---

Ants are highly adaptive insects and are distributed in most terrestrial environments in great abundance and diversity. Many species have aggressive behavior, which in turn may displace other species. Invasive species can handle several types of habitats, such as urban and agricultural areas, that are of great social and economic importance to humans. In this invasive group are the ants of the genus *Solenopsis* (including the well-known fire ant), which occur worldwide and have a wide distribution in Brazil, including in urban areas. They are highly aggressive in defense of their colony and during foraging.

The species *Solenopsis invicta* Buren (Hymenoptera: Formicidae) was spread from South America to various places around the world via wood export (Taber 2000). Their presence has been documented in the United States, the West Indies, New Zealand, Puerto Rico, Australia (Henschaw et al. 2005; Tschinkel 2006), Taiwan (Chen et al. 2006), and China (Yijuan et al. 2007).

In South America, a place of high ant genera diversity, the distinction between *Solenopsis* species is difficult due to a reduced number of diagnostic characteristics (Pitts et al. 2005). An important study by Pitts et al. (2005) about cladistic analysis of the *Solenopsis saevissima* (Smith) species-group represented a step towards understanding the group, but some important unresolved issues remained (see Shoemaker et al. 2006).

Wilson (1952) considered that there may be only three species of fire ants, with South American fire ants comprising a large hybrid ant colony with several variants of hybridizations of parapatric regional areas.

Ross and Shoemaker (2005) conducted a genetic analysis to delineate species of fire ants in South America and found that *S. invicta* and *S. richteri* were reproductively isolated, in contrast to previous findings suggesting the existence of regions in which hybrid colonies existed in the USA (Shoemaker et al. 1996). In addition, it was proposed that the existence of cryptic species in *S. invicta* and *S. richteri* indicated that the group was undergoing radiation and morphological differences that were not leading to reproductive isolation or neutral genetic divergence. It is noteworthy that before Ross and Shoemaker (2005), the occurrence of cryptic species of *S. invicta* were found by observed divergences in mitochondrial DNA by Shoemaker et al. 2003b, and more recently Delsinne et al. (2012) also suggested the existence of cryptic species in *Solenopsis* genus inferred by cytochrome oxidase I (COI) and nuclear wingless genetic markers.

Phylogenetic analysis carried out by Shoemaker et al. (2006) for the species group *Solenopsis saevissima* based on mitochondrial DNA sequences of samples from Brazil and Argentina imply that the group should be monophyletic. However, they found an occurrence of divergent mitochondrial DNA lineages in several species, suggesting a polyphyletic pattern for the invasive *S. invicta*.

According to Ross et al. (2009), high levels of evolutionary divergence and differentiation between regional populations of *S. saevissima* do occur. As these two widely distributed populations are connected by substantial levels of recent gene flow, other groups are evolutionarily independent or on the way to becoming such. Several of these lineages are parapatric with other populations, suggesting that intrinsic barriers to pre-mating and post-mating gene flow are occurring. Ross et al.

(2009) also suggested that genetic differences found in *S. saevissima* might be due to inter-specific hybridization with other regional species that occur in sympatry or parapatry, including *S. geminata*.

Considering that South America is the focus of fire ant occurrence, two aspects are relevant: 1) the Pantanal region of South America is considered the nucleus of dispersion of *S. invicta*, and 2) the other regions of Brazil are dominated by *S. saevissima* (Ahrens et al. 2005; Ross and Shoemaker 2005; Shoemaker et al. 2006; Ross et al. 2009).

Despite the wide distribution of fire ants throughout the world and several studies focusing on understanding their evolution and distribution aspects, there are no specific studies of their distribution in urban or human interfered habitats in Brazil, which is part of their place of origin.

The aims of this study were to characterize the populations of fire ants (*Solenopsis* spp.) from several regions of Brazil and Corrientes, Argentina, through analysis of mitochondrial DNA gene sequences, including part of the COI gene. The focus populations from this analysis were fire ants associated with urban or human-interfered habitats. Through phylogenetic analysis, the degree of relatedness of these populations was determined and their phylogeny was inferred. We first expected to find a prevalent haplotype associated with urban habitats. However, geographical coherence was found in *S. invicta* and *S. saevissima*, but no predominant haplotype was found in the populations analyzed, which clearly illustrates the diversity of the genera in Brazil.

## Materials and Methods

### Specimen collection, identification, and material preservation

The 114 analyzed sample nests were collected by the authors at 42 locations in Brazil, in the states of Amapa, Amazonas, Para, Tocantins, Mato Grosso do Sul, Minas Gerais, Parana, Rio Grande do Sul, Santa Catarina, and Sao Paulo, as well as samples from Corrientes, Argentina. The collected samples were associated with habitats that had been interfered with or disturbed by humans. Table 1 includes the collection codes, locations, species, mitochondrial DNA haplotype, geographic coordinates, and correspondent GenBank accession numbers. The collection sites are shown in Figure 1.

The samples contained workers of various sizes that were fixed and maintained in 80% ethanol under freezing to prevent degradation of DNA until the moment of use. The identification was made based on Trager (1991) and Pitts (2002).

The visual differentiation between different species of *Solenopsis* is hampered due to poor definition of morphological characteristics (Pitts et al. 2005). In this sense, molecular data can clarify the doubts created by morphological identifications and may even be the main tool used to differentiate species by allowing for the creation of a DNA barcode (Hebert et al. 2003a; Hebert et al. 2003b; Ratnasingham and Hebert 2007; Hebert et al. 2010). According to Pitts (2002), there seems to be higher-level concordance between mtDNA data and the morphological data in some *Solenopsis* species. In this sense, we used mitochondrial DNA data, more specifically COI, for species identification confirmation. By sequencing part of the COI gene, fragments were generated for all popula-

tions. Then, using the NCBI Blast (National Center for Biotechnology Information, [www.ncbi.nlm.nih.gov](http://www.ncbi.nlm.nih.gov)), we compared our data with sequences deposited in GenBank. Species identity was confirmed when there was great similarity of the experimental and database sequences; this was defined as either high score values or E-values equal to or close to 0 or very close to those deposited in the database.

### DNA extraction

Total DNA was extracted using a non-phenolic method. Five whole ant workers (pooled) were used. The extraction protocol was the same as used in Martins et al. (2010).

### PCR amplification

Mitochondrial DNA fragments of approximately 920 bp were amplified by PCR. These fragments were part of the COI gene (approximately 780 bp), leucine transfer RNA (70 bp), and part of the cytochrome oxidase II (approximately 60 bp). The amplifications were carried out with a final volume of 25  $\mu$ L, containing 250 to 500 ng of DNA template and 0.2–0.4  $\mu$ M (5–10 pmol) of each primer, using the Ready-to-go kit (Amersham Pharmacia Biotech, GE Healthcare Life Sciences, [www.gelifesciences.com](http://www.gelifesciences.com)).

The thermal cycler was programmed as proposed by Ross and Shoemaker (1997): 1 min at 94°C (initial denaturation) and 35 cycles at 94°C for 1 min, annealing temperature of 48°C for 1 min, and extension temperature of 68°C for 2 min, followed by a final extension step at 72°C for 5 min.

The primers used were: C1-J-2195 (COI-RLR) (5' – TTGATTTTTTGGTCATCCAG AAGT – 3') and DDS-COII-4 (5' – TAAGAT GGTTAATGAAGAGTAG – 3') (Ahrens et al. 2005; Ross and Shoemaker 1997). When

the combination of primers did not amplify the desired fragment, a second primer was used instead of DDS-COII-4, named JerryGarcia-CI (5' – GGG AATTAGAATTTTG AAGAG – 3') (Shoemaker et al. 2006), which produces fragments of approximately 780 bp that include only the gene COI.

### DNA sequencing

DNA was sequenced with the BigDye Terminator Kit (Applied Biosystems, Life Technologies, [www.lifetechnologies.com](http://www.lifetechnologies.com)). Both DNA chains of each sample were sequenced separately with the corresponding primers using an automatic sequencer ABI Prism 377 (Applied Biosystem). DNA sequencing was carried out according to standard protocols. The final volume was 10  $\mu$ L. The extension products were precipitated with 75% isopropanol.

### Phylogenetic analysis

The sequences were initially analyzed separately with BioEdit software ([www.mbio.ncsu.edu/BioEdit/bioedit.html](http://www.mbio.ncsu.edu/BioEdit/bioedit.html)) and aligned using ClustalW software (Higgins et al. 1992) followed by manual modifications. A second and more refined alignment was performed with MUSCLE3.6 software (Edgar 2004).

After all sequences were aligned with the sequences retrieved from GenBank (Table 4), some bases at the end of the fragment were excluded due to unsatisfactory alignment. The resulting matrix consisted of approximately 700 bp comprising only the COI.

The resulting alignment was used for the construction of the network of strains using DnaSP4.90 software (Rozas et al. 2003) and Network4.5 ([www.fluxus-engineering.com](http://www.fluxus-engineering.com)) using the median joining parameter (Bandelt et al. 1999).

The reconstruction of the phylogeny based on maximum parsimony analysis was conducted using PAUP 4.0 software (Swofford 2003). The data set was analyzed using setting 1 for gap and setting 3 for substitutions. One thousand replicates were used to generate bootstrap values.

MrModeltest 2.2 (Nylander 2002) was used before carrying out Bayesian analyses, appropriate models of sequence evolution were chosen via the Akaike information criterion, and the model selected was GTR+I+G. The reconstruction of the phylogeny based on the Bayesian analysis was carried out using MrBayes software (Huelsenbeck et al. 2001). A Markov chain was run for 1,000,000 generations and sampled at each 100 generations. To summarize the parametric values and the trees generated, the first 10% of the trees were excluded as burn-in, and the probability values were then calculated with the remaining trees.

Considering the clade division found in phylogenetics analysis, we analyzed clade 3, clade 5, clade 6, and clade 7 in terms of haplotype diversity, nucleotide diversity, Tajima's D, polymorphic sites, G+C content, and average number of nucleotide differences between populations with DnaSP4.90 software (Rozas et al. 2003).

## Results

Of the 114 analyzed colonies, 72 had a unique haplotype sequence of the COI mitochondrial DNA, which are illustrated in the network (Figure 2). Table 1 includes the species identification, the collecting locales (georeferenced), and the corresponding haplotypes. All COI sequences generated in this study have been deposited in the GenBank

database under accession numbers JN808775 to JN808838 (see Table 1).

The prevalent haplotypes were H59 (Argentina); H7, H9, and H11 (in southeastern Brazil); H31 (midwest Brazil); H68 and H3 (southern Brazil); and H13, H64, and H75 (northern Brazil). Furthermore, the remaining haplotypes did not seem to have derived from the most prevalent ones (see Figure 2). Moreover, the haplotype distribution in the network indicates that there was no shared haplotypes between different localities, suggesting great diversity of *Solenopsis* in Brazil, but now seen in a view from those ants associated with human-disturbed habitats.

Of the 726 characters used in maximum parsimony analysis, 482 were constant and 206 were informative characters (parsimony-informative). Forty-eight equally parsimonious trees were found as a result of phylogenetic analysis of different *Solenopsis* populations based on a portion of the COI gene. Both analyses (maximum parsimony and Bayesian inference) were nearly the same, and only the Bayesian tree analysis was illustrated with posterior probability values (Figure 3).

The phylogenetic tree was rooted with a representative of the species *Monomorium pharaonis* recovered from GenBank. Several internal sequences from GenBank of some species of *Solenopsis* were also incorporated into the analysis (Table 2). The clusters in the phylogenetic tree (Figure 3) revealed the occurrence of several clades, and most are well supported. Important clades include the following:

Clade 1: The presence of diverging species grouped as closely-related species disagrees with the phylogeny proposed by Shoemaker et

al. (2006) and Tschinkel et al. (2006). The two species *S. geminata* and *S. saevissima* are morphologically distinct, even though they have been grouped in this clade.

Clade 2: Representatives of *S. pusilligni*, collected in Ladário, Mato Grosso do Sul, cluster with the representative of the same species recovered from GenBank (AY950775).

Clade 3: Representatives of *S. saevissima* populations from the northern region of Brazil along with representatives listed in GenBank (AY950715, FJ467557, FJ467537). This clade appears to differ from populations of *S. saevissima* from southern Brazil (clade 7).

Clade 4: Once again, the presence of diverging species grouped as closely-related species (*S. saevissima*, *S. megergates*, and *S. invicta*).

Clade 5: Representatives of *S. invicta* populations from the states of São Paulo, Paraná, Mato Grosso, and Mato Grosso do Sul form a well-supported clade of specimens of *S. invicta* with restricted occurrence to this geographical area.

Clade 6: A second clade of *S. invicta* occurs in populations from the state of Rio Grande do Sul and Santa Catarina.

Clades 8, 9, and 10: Representatives of the *S. invicta* and *S. saevissima* species form an isolated group of representatives of these species that are not allocated in previous clades. The terminal clade contains the representative of *S. invicta* from the Rio de Janeiro.

The results of this analysis reveal the existence of well-supported clades of *S. invicta* and *S. saevissima* from different geographical regions that are split between *S. saevissima* that occur in the North, South, southern, and Mid-

west Brazil. *S. invicta* also has separate representatives from the South and Southeast.

G+C content of the samples was approximately 30%, corroborating the high A+T frequencies expected for insects (Simon et al. 1994). The average number of nucleotide differences between clades 5 and 6 was 18,420 and between clades 3 and 7 was 65,229.

The number of haplotypes found in clade 7 was the lowest of all. As for haplotype diversity, clade 3 was the lowest, followed by clade 5, 7, and 6. Regarding nucleotide diversity ( $\pi$ ), clade 7 was the lowest, followed by 5, 6, and 3. Tajima's D for clade 3 and 6 was negative (-0.02978 and -0.18128 respectively) and for clade 5 and 7 was positive (0.13862 and 0.36991 respectively). Clade 7 was the one with the lowest polymorphic sites, followed by clade 5, 6, and 3 (see Table 3 to summarize results).

## Discussion

The results show complete consistency when grouping populations according to geographical distribution and even polyphyly for *S. invicta* and *S. saevissima*, which reveals diversity of this ant genus in Brazil. However grouping of divergent species (see clades 1, 4, 6, 8, and 9 in Figure 3) could be due to a lack of data collection in a region where representatives of these species occur, which could bring together representatives to form new clades, such as those found by Shoemaker et al. (2006), or indicative that those haplotypes that could form clades not supported here rarely occur in urban areas.

The polyphyly found in *S. invicta* (Figure 3) was also observed by Shoemaker et al. 2000, 2003, and 2006. Shoemaker et al. (2006) found discordance between the phylogeny re-

constructed with mtDNA haplotypes and those constructed using morphology, and they reported seven well-supported clades of the species *S. invicta*. They suggested that the polyphyly of the mitochondrial DNA sequence of these species appears to result in the crowding of multiple morphological characteristics that represent genetic lineages that are evolutionarily indistinguishable and independent (cryptic species). They concluded that current morphological boundaries overestimate the distribution of fire ants and assumed that the mtDNA tree they reconstructed faithfully categorized the development of reproductive isolation and patterns of ancestral populations of the *S. saevissima* species group.

The geographical grouping of *S. invicta* (clades 5 and 6) and *S. saevissima* (clades 3 and 7) supports the hypothesis that regional populations of each species are derived from refuges or large isolated areas of earlier endemism from which expansion has occurred (Ahrens et al. 2005; Shoemaker et al. 2006). Because our focus was populations from *Solenopsis* from habitats that were affected by humans, this expansion could be driven by human activities.

Climate patterns could also be the cause of the presence of divergent clades of *S. saevissima* (clades 3 and 7). The third clade has representatives from the states of Tocantins, Amazonas, Pará, and Amapá, which are characterized by hot climates from semi-humid (Tocantins) to wet (Amazonas, Pará, and Amapá) located within the Amazon biome; the seventh clade has representatives from the states in southern Brazil where the climate is predominantly hot (São Paulo) and mid-mesothermal (Rio Grande do Sul and Santa Catarina). On the other hand, the high nucleotide diversity observed in clade 3 ( $D < 0$ )

could be related to geographic distribution because the northern region of Brazil is less populated, so human interference is reduced, which in turn can reduce the pressure against divergence of those ants populations consequently experiencing rapid growth.

The presence of the endosymbiont *Wolbachia* can influence cytoplasmic genome selection, such as on host mtDNA evolution (Shoemaker et al. 2003). The presence of *Wolbachia* may be related to the divergence of the *S. invicta* and *S. saevissima* clades in the present study. It may indicate traces of increased substitution rates in mtDNA associated with recurrent *Wolbachia* infections in affected lineages (Shoemaker et al. 2004). For this scenario to be possible, separate clades of concerned species should be fully isolated or have minimal evidence of migration, which should be the case for the populations studied herein that are geographically separated by many miles. Additionally, *Solenopsis* populations from Brazil with high rates of *Wolbachia* infection offset other populations in which the infection rate is low or absent (Martins et al. 2012).

Populations from *S. saevissima* comprising clade 3 show  $D < 0$ , indicating that this population has experienced rapid growth, and those from clade 7 show  $D > 0$ , indicating that this population has experienced recent bottleneck. *S. invicta* populations from clade 5 presented  $D > 0$  and from clade 6  $D < 0$ , indicating respectively recent bottleneck and rapid growth. In clade 3, as already discussed, populations may be under less human interference and are rarely infected by *Wolbachia*. The populations in clade 6 are probably more affected or influenced by human activity and are also most infected by *Wolbachia*, however clade 3 and 6 are representatives of different species, which may indicate different responses to different selective pressures.

The absence of gene flow may occur due to the existence of a few mtDNA haplotypes that are shared by colonies of different locations. Of the total 72 haplotypes found, only 13 were shared by colonies of different locations. This is evidenced by the size of the circles that indicate the network in Figure 2 (and Table 1 for reference locations). The absence of gene flow may be due to the Paraná River acting as a physical barrier.

The Paraná River is a natural barrier and can restrict gene flow, which could affect the structure and evolution of populations of native fire ants (Ross et al. 1997), but according to Ahrens et al. (2005) there is no explicit information that supports this hypothesis. Likewise, the present study also provides no information to support this hypothesis, although clade 5 included *S. invicta* from northwestern Brazil separated from those of southeastern Brazil. In clade 6, *S. invicta* from Campo Grande (Mato Grosso do Sul) grouped with others from southern Brazil and are therefore separated by the Paraná River, which may mean that mitochondrial gene flow between these regions can occur.

The effects of geographical barriers can be minimized by the constant transport of ants by human activity, resulting in subsequent gene flow. Ahrens et al. (2005) considered that these movements should not be so frequent, as to prevent the continued geographic genetic divergence of populations, but it is important to note that in addition to several species of the genus *Solenopsis* having status as invaders, there is intense trade between certain regions of Brazil, which could explain the occurrence of populations that appear northwest of the Paraná River and in southern Brazil. This may hinder our understanding of the true distribu-

tion and evolutionary history of the genus in its region of origin.

This hypothesis finds support in the proposal by Lofgren (1986) and is also emphasized by Tschinkel (2006), as they reported that human activity was a factor in *S. invicta* dispersal the USA after its introduction. Despite the great diversity of ants in South America, including the *Solenopsis* genus, some species may be favored over others by the reduction of native forests and establishment of monocultures.

Because fire ants have become a global pest, resolving interspecific relationships and species limits is important in understanding the patterns of diversification in South America, their place of origin, and the dispersal of fire ants, especially their distribution in urban or human-influenced habitats. This study shows the need to expand studies using molecular markers in populations of fire ants that occur in urban areas in order to understand the mechanisms these populations are going through and the relationship between rapid urbanization and its relationship with natural populations in these urban areas.

### Acknowledgements

Thanks are due to CNPq, National Council for Scientific and Technological Development.

### References

- 
- Ahrens ME, Ross KG, Shoemaker DD. 2005. Phylogeographic structure of the fire ant *Solenopsis invicta* in its native South American range: roles of the natural barriers and habitat connectivity. *Evolution* 59: 1733-1743.
- Bandelt H-J, Foster P, Röhl A. 1999. Median-



joining networks for inferring intraspecific phylogenies. *Molecular Biology and Evolution* 16: 37-48.

Chen JSC, Shen C-H, Lee H-J. 2006. Monogynous and polygynous red imported fire ants, *Solenopsis invicta* Buren (Hymenoptera: Formicidae), in Taiwan. *Environmental Entomology* 35: 167-172.

Delsinne T, Sonet G, Nagy ZT, Wauters N, Jacquemin J, Leponce M. 2012. High species turnover of the genus *Solenopsis* (Hymenoptera: Formicidae) along an altitudinal gradient in the Ecuadorian Andes, indicated by a combined DNA sequencing and morphological approach. *Invertebrate Systematics* 26(6): 457-469.

Edgar RC. 2004. MUSCLE: Multiple sequence alignment with high accuracy and high throughput. *Nucleic Acid Research* 32: 1792-97.

Hebert PD, Deward JR, Landry JF. 2010. DNA barcodes for 1/1000 of the animal kingdom. *Biology Letters* 6(3): 359-62.

Hebert PDN, Cywinska A, Ball SL, deWaard JR. 2003a. Biological identifications through DNA barcodes. *Proceedings of the Royal Society of London Series B: Biological Sciences* 270: 313-321.

Hebert PDN, Ratnasingham S, deWaard JR. 2003b. Barcoding animal life: cytochrome *c* oxidase subunit 1 divergences among closely related species. *Proceedings of the Royal Society of London Series B: Biological Sciences* 270: 596-599.

Henschaw MT, Kunzmann N, Vanderwoude C, Senetra M, Crozier RH. 2005. Population genetics and history of the introduced fire ant,

*Solenopsis invicta* Buren (Hymenoptera: Formicidae) in Australia. *Austral Journal of Entomology* 44: 37-44.

Higgins DG, Bleasby AJ, Fuchs R. 1992. Clustal V: Improved software for multiple sequence alignment. *Computer Applications in the Biosciences* 8: 189-191.

Huelsenbeck JP, Ronquist F, Nielsen R, Bollback JP. 2001. Evolution: Bayesian inference of phylogeny and its impact on evolutionary biology. *Science* 294: 2310-2314.

Lofgren, CS. 1986. History of imported fire ants in the United States. In: Lofgren CS, Vander Meer RK, Editors. *Fire ants and leaf cutting ants: biology and management*. Westview Press.

Martins C, Fontes LR, Bueno OC, Martins VG. 2010. *Coptotermes gestroi* (Isoptera: Rhinotermitidae) in Brazil: possible origins inferred by mitochondrial cytochrome oxidase II gene sequences. *Genome* 53(9): 651-657.

Martins C, Souza RF, Bueno OC. 2012. Presence and distribution of the endosymbiont *Wolbachia* among *Solenopsis* spp. (Hymenoptera: Formicidae) from Brazil and its evolutionary history. *Journal of Invertebrate Pathology* 109: 287-296.


Nylander JAA. 2002. *MrModeltest 2.2*. Evolutionary Biology Centre, Uppsala University, Uppsala, Sweden.

Pitts JP. 2002. *A cladistic analysis of the Solenopsis saevissima species-group* (Hymenoptera: Formicidae). PhD thesis, University of Georgia, USA.


- Pitts JP, McHugh JV, Ross KG. 2005. Cladistic analysis of the fire ants of the *Solenopsis saevissima* species-group (Hymenoptera: Formicidae). *Zoologica Scripta* 34: 493-505.
- Ratnasingham S, Hebert PDN. 2007. BOLD: The Barcode of life system ([www.barcodinglife.org](http://www.barcodinglife.org)). *Molecular Ecology Notes* 1-10.
- Ross KG, Gotzek D, Ascunce MS, Shoemaker DD. 2009. Species delimitation: a case study in a problematic ant taxon. *Systematic Biology* 59: 1-23.
- Ross KG, Krieger MJB, Shoemaker DD, Vargo EL, Keller L. 1997. Hierarchical analysis of genetic structure in native fire ant populations: results from three classes of molecular markers. *Genetics* 147: 643-655.
- Ross KG, Shoemaker DD. 1997. Nuclear and mitochondrial genetic structure in two social forms of the fire ant *Solenopsis invicta*: insights into transitions to an alternate social organization. *Heredity* 78: 590-602.
- Ross KG, Shoemaker DD. 2005. Species delimitation in native South American fire ants. *Molecular Ecology* 14: 3419-3438.
- Rozas J, Sánchez-Delbarrio JC, Messeguer X, Rozas R. 2003. DnaSP, DNA polymorphism analyses by the coalescent and other methods. *Bioinformatics* 19: 2496-2497.
- Shoemaker DD, Ahrens ME, Ross KG. 2006. Molecular phylogeny of the fire ants of the *Solenopsis saevissima* species-group based on mtDNA sequences. *Molecular Phylogenetics and Evolution* 38: 200-215.
- Shoemaker DD, Dyer KA, Ahrens M, McAbee K, Jaenike J. 2004. Decreased diversity but increased substitution rate in host mtDNA as a consequence of *Wolbachia* endosymbiont infection. *Genetics* 168: 2049-2058.
- Shoemaker DD, Keller G, Ross KG. 2003. Effects of *Wolbachia* on mtDNA variation in two fire ant species. *Molecular Ecology* 12: 1757-1772.
- Shoemaker DD, Ross KG, Arnold ML. 1996. Genetic structure and evolutions of the fire ant hybrid zone. *Evolution* 50: 1958-1976.
- Shoemaker DD, Ross KG, Keller L, Vargo EL, Werren JH. 2000. *Wolbachia* infections in native and introduced populations of the fire ants (*Solenopsis* spp.). *Insect Molecular Biology* 9: 661-673.
- Simon C, Frati F, Beckenbach A, Crespi B, Liu H, Flook P. 1994. Evolution, weighting, and phylogenetic utility of mitochondrial gene sequences and a compilation of conserved polymerase chain reaction primers. *Annals of the Entomological Society of America* 87:651-701.
- Swofford DL. 2003. *Phylogenetic analyses using parsimony, version 4.0b10*. Smithsonian Institution, Washington DC.
- Taber SW. 2000. *Fire ants*, Texas A&M University Press.
- Trager JC. 1991. A revision of the fire ants, *Solenopsis geminata* group (Hymenoptera: Formicidae: Myrmicinae). *Journal of the New York Entomological Society* 99: 141-198.
- Tschinkel WR. 2006. *The fire ants*, The Belknap Press of Harvard University Press.


Wilson EO. 1952. O complexo *Solenopsis saevissima* na América do Sul (Hymenoptera: Formicidae). *Memórias do Instituto Oswaldo Cruz* 50: 49-68.

Yijuan X, Youngyue L, Ling Z, Guangwen L. 2007. Foraging behavior and recruitment of red imported fire ant *Solenopsis invicta* Buren in typical habitats of South China. *Acta Ecologica Sinica* 27: 855-860.



**Figure 1.** South America map showing the 42 localities of collection of *Solenopsis* spp. Table 1 shows details. Map created with speciesMapper (<http://smlink.cria.org.br/mapper?criaLANG=pt>). High quality figures are available online.





**Figure 3.** Bayesian inference tree reconstructed from the COI gene in populations of *Solenopsis* from different regions of Brazil and part of Argentina. Posterior probability values are shown above branches. The tree was rooted with a representative of *Monomorium pharaonis*. High quality figures are available online.

**Table 1.** Ant species, collection codes and locations, mitochondrial DNA haplotype, geographic coordinates, GenBank accession numbers.

Ant species	Collection codes and locations	Mitochondrial DNA haplotype	Geographic coordinates	GenBank accession numbers
<i>S. geminata</i>	E1820	H63	S03°06'25"	JN808821
	Manaus, AM		W60°01'34"	
<i>S. geminata</i>	E1825	H64	S03°06'25"	JN808822
	Manaus, AM		W60°01'34"	
<i>S. geminata</i>	E1818	H70	S03°06'25"	JN808828
	Manaus, AM		W60°01'34"	
<i>S. geminata</i>	E1823	H73	S03°06'25"	JN808822
	Manaus, AM		W60°01'34"	
<i>S. geminata</i>	E1830	H77	S03°06'25"	JN808822
	Manaus, AM		W60°01'34"	
<i>S. geminata</i>	E1826	H75	S03°06'25"	JN808832
	Manaus, AM		W60°01'34"	
<i>S. geminata</i>	E1827	H75	S03°06'25"	JN808832
	Manaus, AM		W60°01'34"	
<i>S. geminata</i>	E1832	H78	S03°06'25"	JN808832
	Manaus, AM		W60°01'34"	
<i>S. geminata</i>	E1822	H78	S03°06'25"	JN808832
	Manaus, AM		W60°01'34"	
<i>S. invicta</i>	E1710	H11	S22°58'51"	JN808784
	Rio de Janeiro, RJ		W43°16'75"	
<i>S. invicta</i>	E1628	H4	S20°27'59"	JN808778
	Campo Grande, MS		W54°35'33"	
<i>S. invicta</i>	E1648	H5	S26°47'06"	JN808779
	Caçador, SC		W50°59'27"	
<i>S. invicta</i>	E1652	H2	S20°14'29"	JN808776
	Miranda, MS		W56°22'43"	
<i>S. invicta</i>	E1680	H3	S29°07'22"	JN808777
	Bento Gonçalves, RS		W51°20'58"	
<i>S. invicta</i>	E1683	H8	S23°30'21"	JN808782
	Ubatuba, SP		W45°07'55"	
<i>S. invicta</i>	E1684	H9	S23°30'21"	JN808783
	Ubatuba, SP		W45°07'55"	
<i>S. invicta</i>	E1685	H9	S23°30'21"	JN808783
	Ubatuba, SPB		W45°07'55"	
<i>S. invicta</i>	E1686	H11	S23°19'02"	JN808784
	Picinguaba, SP		W44°54'04"	
<i>S. invicta</i>	E1704	H30	S19°30'31"	JN808792
	Corumbá, MS		W57°20'05"	
<i>S. invicta</i>	E1705	H31	S18°45'11"	JN808793
	Corumbá, MS		W57°07'09"	
<i>S. invicta</i>	E1706	H32	S18°50'00"	JN808794
	Corumbá, MS		W57°18'55"	
<i>S. invicta</i>	E1707	H33	S18°50'00"	JN808795
	Corumbá, MS		W57°18'55"	
<i>S. invicta</i>	E1709	H11	S22°58'51"	JN808784
	Rio de Janeiro, RJ		W43°16'75"	
<i>S. invicta</i>	E1720	H37	S25°25'46"	JN808799
	Paraná		W49°16'18"	
<i>S. invicta</i>	E1721	H31	S19°00'23"	JN808793
	Corumbá, MS		W57°39'10"	
<i>S. invicta</i>	E1722	H31	S19°00'23"	JN808793
	Corumbá, MS		W57°39'10"	
<i>S. invicta</i>	E1723	H31	S14°09'40"	JN808793
	Porto Esperança, MT		W56°04'38"	
<i>S. invicta</i>	E1724	H40	S14°09'40"	JN808801
	Porto Esperança, MT		W56°04'38"	
<i>S. invicta</i>	E1725	H41	S29°59'14"	JN808802
	Porto Alegre, RS		W51°09'580"	
<i>S. invicta</i>	E1726	H41	S29°59'14"	JN808802
	Porto Alegre, RS		W51°09'580"	
<i>S. invicta</i>	E1727	H42	S29°59'14"	JN808803
	Porto Alegre, RS		W51°09'580"	
<i>S. invicta</i>	E1737	H36	S22°23'34"	JN808798
	Rio Claro, SP		W47°33'21"	
<i>S. invicta</i>	E1739	H68	S22°58'51"	JN808826
	Rio de Janeiro, RJ		W43°16'75"	
<i>S. invicta</i>	E1741	H11	S22°58'51"	JN808784
	Rio de Janeiro, RJ		W43°16'75"	
<i>S. invicta</i>	E1744	H43	S23°30'21"	JN808804
	Ubatuba, SP		W45°07'55"	
<i>S. invicta</i>	E1748	H46	S23°30'21"	JN808782
	Ubatuba, SP		W45°07'55"	
<i>S. invicta</i>	E1749	H9	S23°30'21"	JN808783
	Ubatuba, SP		W45°07'55"	
<i>S. invicta</i>	E1752	H9	S23°30'21"	JN808783
	Ubatuba, SP		W45°07'55"	

Table 1 continued.

<i>S. invicta</i>	E1754	H50	S23°12'22"	JN808783
	São Luiz do Paraitinga, SP		W45°20'43"	
<i>S. invicta</i>	E1768	H51	S20°28'46"	JN808809
	Anastácio, MS		W55°48'08"	
<i>S. invicta</i>	E1770	H51	S20°28'42"	JN808809
	Aquidauana, MS		W55°47'03"	
<i>S. invicta</i>	E1771	H53	S20°28'42"	JN808811
	Aquidauana, MS		W55°47'03"	
<i>S. invicta</i>	E1780	H9	S24°31'46"	JN808783
	Registro, SP		W47°51'24"	
<i>S. invicta</i>	E1781	H9	S24°31'46"	JN808783
	Registro, SP		W47°51'24"	
<i>S. invicta</i>	E1783	H54	S27°41'42"	JN808812
	Alfredo Wagner, SC		W49°19'53"	
<i>S. invicta</i>	E1786	H55	S29°09'48"	JN808813
	Bento Gonçalves, RS		W51°31'54"	
<i>S. invicta</i>	E1788	H56	S29°07'21"	JN808814
	Pinto Bandeira, RS		W51°26'56"	
<i>S. invicta</i>	E1789	H56	S29°07'21"	JN808814
	Pinto Bandeira, RS		W51°26'56"	
<i>S. invicta</i>	E1794	H58	S25°25'42"	JN808816
	Curitiba, PR		W49°16'25"	
<i>S. invicta</i>	E1808	H59	S27°18'39"	JN808817
	Corrientes, Argentina		W58°33'44"	
<i>S. invicta</i>	E1807	H59	S27°18'39"	JN808817
	Corrientes, Argentina		W58°33'44"	
<i>S. invicta</i>	E1805	H59	S27°18'39"	JN808817
	Corrientes, Argentina		W58°33'44"	
<i>S. invicta</i>	E1801	H59	S27°18'39"	JN808817
	Corrientes, Argentina		W58°33'44"	
<i>S. invicta</i>	E1803	H59	S27°18'39"	JN808817
	Corrientes, Argentina		W58°33'44"	
<i>S. invicta</i>	E1802	H59	S27°18'39"	JN808817
	Corrientes, Argentina		W58°33'44"	
<i>S. invicta</i>	E1806	H59	S27°18'39"	JN808817
	Corrientes, Argentina		W58°33'44"	
<i>S. invicta</i>	E1810	H59	S27°18'39"	JN808817
	Corrientes, Argentina		W58°33'44"	
<i>S. invicta</i>	E1798	H43	S08°07'49"	JN808804
	Recife, PE		W34°54'09"	
<i>S. invicta</i>	E1799	H37	S23°25'35"	JN808799
	Maringá, PR		W51°56'46"	
<i>S. invicta</i>	E1800	H37	S23°25'35"	JN808799
	Maringá, PR		W51°56'46"	
<i>S. invicta</i>	E1784	H61	S27°48'57"	JN808819
	Lages, SC		W50°22'17"	
<i>S. invicta</i>	E1787	H56	S29°07'21"	JN808814
	Pinto Bandeira, RS		W51°26'56"	
<i>S. invicta</i>	E1790	H62	S28°00'23"	JN808820
	Capão Alto, RS		W50°32'26"	
<i>S. invicta</i>	E1815	H69	S31°46'33"	JN808827
	Pelotas, RS		W52°20'33"	
<i>S. invicta</i>	E1816	H79	S31°46'33"	JN808827
	Pelotas, RS		W52°20'33"	
<i>S. invicta</i>	E1646	H85	S26°46'32"	JN808838
	Caçador, SC		W51°00'56"	
<i>S. invicta</i>	E1645	H84	S26°46'32"	JN808837
	Caçador, SC		W51°00'56"	
<i>S. megergates</i>	E1782	H68	S26°33'53"	JN808826
	São Francisco, SC		W48°43'10"	
<i>S. megergates</i>	E1793	H68	S25°51'45"	JN808826
	Areia Branca, PR		W19°21'45"	
<i>S. megergates</i>	E1644	H68	S26°46'32"	JN808826
	Caçador, SC		W51°00'56"	
<i>S. megergates</i>	E1643	H68	S26°46'32"	JN808826
	Caçador, SC		W51°00'56"	
<i>S. pusillignis</i>	E1657	H17	S19°01'05"	JN808790
	Ladário, MS		W57°33'04"	
<i>S. pusillignis</i>	E1708	H34	S19°01'03"	JN808796
	Ladário, MS		W57°34'11"	
<i>S. saevissima</i>	E1608	H16	S6°38'55"	JN808789
	Eirunepé, AM		W69°52'32"	
<i>S. saevissima</i>	E1615	H6	S22°23'34"	JN808780
	Rio Claro, SP		W47°33'44"	
<i>S. saevissima</i>	E1631	H7	S22°26'11"	JN808781
	Ipeúna, SP		W47°43'10"	

Table I continued.

<i>S. saevissima</i>	E1640	H15	S01°23'28"	JN808788
	Belém, PA		W48°28'43"	
<i>S. saevissima</i>	E1650	H39	S21°45'51"	JN808800
	Juiz de Fora, MG		W43°20'56"	
<i>S. saevissima</i>	E1662	H13	S00°00'23"	JN808786
	Macapá, AP UFAP		W51°05'06"	
<i>S. saevissima</i>	E1666	H14	S00°02'19"	JN808787
	Macapá, AP IEPA		W51°05'39"	
<i>S. saevissima</i>	E1671	H12	S00°02'19"	JN808785
	Macapá, AP		W51°05'39"	
<i>S. saevissima</i>	E1682	H1	S29°04'31"	JN808775
	Bento Gonçalves, RS		W51°14'13"	
<i>S. saevissima</i>	E1712	H29	S22°23'47"	JN808791
	Rio Claro, SP		W47°32'51"	
<i>S. saevissima</i>	E1713	H35	S17°25'20"	JN808797
	Buritizeiro, MG		W44°56'54"	
<i>S. saevissima</i>	E1714	H35	S17°25'20"	JN808797
	Buritizeiro, MG		W44°56'54"	
<i>S. saevissima</i>	E1738	H60	S22°58'51"	JN808818
	Rio de Janeiro, RJ		W43°16'75"	
<i>S. saevissima</i>	E1740	H60	S22°58'51"	JN808818
	Rio de Janeiro, RJ		W43°16'75"	
<i>S. saevissima</i>	E1742	H7	S23°32'53"	JN808781
	São Paulo, SP		W46°38'11"	
<i>S. saevissima</i>	E1743	H7	S23°30'21"	JN808781
	Ubatuba, SP		W45°07'55"	
<i>S. saevissima</i>	E1746	H44	S23°30'21"	JN808805
	Ubatuba, SP		W45°07'55"	
<i>S. saevissima</i>	E1747	H45	S23°30'21"	JN808806
	Ubatuba, SP		W45°07'55"	
<i>S. saevissima</i>	E1750	H47	S23°30'21"	JN808791
	Ubatuba, SP		W45°07'55"	
<i>S. saevissima</i>	E1751	H48	S23°30'21"	JN808807
	Ubatuba, SP		W45°07'55"	
<i>S. saevissima</i>	E1753	H49	S23°30'21"	JN808808
	Ubatuba, SP		W45°07'55"	
<i>S. saevissima</i>	E1769	H52	S20°47'37"	JN808810
	Três Lagoas, MS		W51°37'59"	
<i>S. saevissima</i>	E1791	H57	S27°15'32"	JN808815
	São Cristóvão do Sul, SC		W50°26'50"	
<i>S. saevissima</i>	E1792	H57	S27°15'32"	JN808815
	São Cristóvão do Sul, SC		W50°26'50"	
<i>S. saevissima</i>	E1738	H60	S22°58'51"	JN808818
	Rio de Janeiro, RJ		W43°16'75"	
<i>S. saevissima</i>	E1740	H60	S22°58'51"	JN808818
	Rio de Janeiro, RJ		W43°16'75"	
<i>S. saevissima</i>	E1716	H65	S10°42'37"	JN808823
	Porto Nacional, TO		W48°24'34"	
<i>S. saevissima</i>	E1717	H66	S10°19'07"	JN808824
	Taquaraçu, TO		W48°09'22"	
<i>S. saevissima</i>	E1718	H81	S10°19'07"	JN808834
	Palmas, TO		W48°09'22"	
<i>S. saevissima</i>	E1719	H67	S10°19'07"	JN808825
	Palmas, TO		W48°09'22"	
<i>S. saevissima</i>	E1819	H71	S03°06'25"	JN808829
	Manaus, AM		W60°01'34"	
<i>S. saevissima</i>	E1821	H72	S03°06'25"	JN808830
	Manaus, AM		W60°01'34"	
<i>S. saevissima</i>	E1824	H74	S03°06'25"	JN808831
	Manaus, AM		W60°01'34"	
<i>S. saevissima</i>	E1829	H76	S03°06'25"	JN808829
	Manaus, AM		W60°01'34"	
<i>S. saevissima</i>	E1831	H74	S03°06'25"	JN808831
	Manaus, AM		W60°01'34"	
<i>S. saevissima</i>	E1833	H80	S04°01'33"	JN808833
	Pindoretama, CE		W38°18'24"	
<i>S. saevissima</i>	E1718	H81	S10°12'46"	JN808834
	Palmas, TO		W48°21'37"	
<i>S. saevissima</i>	E1785	H82	S27°48'57"	JN808835
	Lages, SC		W50°22'17"	
<i>S. saevissima</i>	E1828	H71	S03°06'25"	JN808829
	Manaus, AM		W60°01'34"	
<i>S. saevissima</i>	E1795	H83	S08°07'44"	JN808836
	Recife, PE		W34°54'13"	



**Table 2.** Ant species used as out-group and in-group in phylogenetic analyses and respective GenBank accession numbers, designed haplotypes from these analyses, and collection location retrieved from data in GenBank.

Species	GenBank accession numbers	mtDNA haplotype	Location
<i>Monomorium pharaonis</i>	DQ023095	H26	-
<i>S. geminata</i>	AY254489	H27	USA
<i>S. invicta</i>	AY950708	H89	Rio Negro: Brazil
<i>S. invicta</i>	AY950734	H91	Céu Azul: Brazil
<i>S. invicta</i>	AY950768	H93	Corumbá: Brazil
<i>S. invicta</i>	AY950747	H92	Rinco dos Cabrais: Brazil
<i>S. invicta</i>	EU352607	H95	Mississippi: USA
<i>S. invicta</i>	DQ831672	H95	China
<i>S. invicta</i>	AY950750	H98	Corrientes: Argentina
<i>S. invicta</i>	EU373819	H97	Louisiana: USA
<i>S. invicta</i>	EU352608	H100	Mississippi: USA
<i>S. invicta</i>	AY950765	H10	Comodoro: Brazil
<i>S. megergates</i>	AY950773	H20	Curitiba: Brazil
<i>S. pusillignis</i>	AY950775	H25	Corumbá: Brazil
<i>S. saevissima</i>	FJ467529	H103	Nordeste: Brazil
<i>S. saevissima</i>	AY950715	H86	Pará: Brazil
<i>S. saevissima</i>	FJ467557	H88	Nordeste: Brazil
<i>S. saevissima</i>	FJ467537	H87	Norte: Brazil
<i>S. saevissima</i>	AY950711	H90	Rio Negro: Brazil
<i>S. saevissima</i>	FJ467520	H101	Sudeste: Brazil
<i>S. saevissima</i>	AY950783	H38	Água Clara: Brazil
<i>S. saevissima</i>	FJ467540	H94	Norte: Brazil
<i>S. saevissima</i>	FJ467547	H99	Nordeste: Brazil
<i>S. saevissima</i>	FJ467527	H102	Nordeste: Brazil
<i>S. saevissima</i>	AY950783	H38	Água Clara: Brazil
<i>S. saevissima</i>	FJ467520	H82	Sudeste: Brazil
<i>Solenopsis</i> sp.	AY950784	H22	Santa Fé: Argentina

**Table 3.** Number of haplotypes for each clade, haplotype diversity, nucleotide diversity, Tajima's D, and polymorphic sites.

Clade	Number of haplotypes	Haplotype diversity	Nucleotide diversity ( $\pi$ )	Tajima's D	Polymorphic sites
Clade 3 (Ssae)	10	0.955	0.02859	-0.02978	58
Clade 5 (Sinv)	11	0.962	0.01789	0.13862	29
Clade 6 (Sinv)	10	0.982	0.01966	-0.18128	31
Clade 7 (Ssae)	7	0.964	0.01709	0.36991	26

Ssae = *S. saevissima*, Sinv = *S. invicta*.