



Pharmacology of Antitubercular Drugs: Challenges and Advances

Jean Leandro dos Santos* and Chung Man Chin

Department of Drugs and Medicines, School of Pharmaceutical Science, State University of São Paulo, Brazil

Keywords: Tuberculosis; Drug discovery; Antitubercular drugs; New drugs; Multidrug-resistant

Abbreviations: TB: Tuberculosis; MTB: *Mycobacterium tuberculosis*; WHO: World Health Organization; MRD-TB: Multidrug-Resistant Tuberculosis; XDR-TB: Extensively Drug-Resistant Tuberculosis; HIV: Human Immunodeficiency Virus; FDA: Food and Drug Administration

Tuberculosis (TB) is a highly prevalent infectious disease caused by *Mycobacterium tuberculosis* (MTB). The World Health Organization (WHO) recently estimated that one-third of the world's population is infected with MTB and that 13.7 million people worldwide have active TB [1]. Long-term therapy consists of four drugs (isoniazid, rifampin, ethambutol, and pyrazinamide), which are administered for 2 months, followed by two drugs (isoniazid and rifampin), which are used in combination for 4 months [2]. Although this treatment regimen has reduced the mortality rate worldwide, the emergence of multidrug-resistant TB (MDR-TB), extensively drug-resistant TB (XDR-TB), and TB/HIV co-infection are challenges that need to be overcome [3]. The mechanism involved in drug resistance is still not fully understood, but several factors are involved, including changes in membrane permeability, mutations in or modifications to cellular targets, and increased expression of efflux pumps [4].

In recent years, the number of new drugs approved for treating TB has not increased proportionally to the investment made in this clinical field. Since the discovery of rifampicin in the 1950s, the last drug approved by United States Food and Drug Administration for treating TB was bedaquiline (Sirturo; Janssen Pharmaceuticals Inc./ Johnson & Johnson) in 2012 [5]. Despite the lack of new anti-TB drugs, it is clear from the scientific literature that researchers are still pursuing new potent and safe anti-TB drugs. We recently performed a review of articles published between 2008 and 2013 in several databases, including Science Direct, Medline, Scopus, and EMBASE, to identify novel anti-TB compounds with activity against MTB strain H37Rv, a minimum inhibitory concentration (MIC) of <10 μ M, and a selective index (SI) of >10. The SI was calculated as the median inhibitory concentration (IC₅₀)/MIC. This literature search revealed 110 different chemical scaffolds, but <10% were evaluated against resistant strains [6].

All of the anti-TB compounds reported in the literature were derived from three main sources: 1) natural products, 2) synthetic organic compounds, and 3) synthetic inorganic compounds. In particular, several organic compounds have been obtained using molecular modification strategies [7]. Koul et al. reported that the anti-TB compounds used in clinical trials, including SQ109, PNU-100480, PA-824, and OPC-67683, were derived from known antimicrobial compounds [8]. Our review revealed that fluoroquinolones and quinoxaline derivatives were the most active compounds. The fluoroquinolone derivative 7-(3-(diethylcarbamoyl)piperidin-1-yl)-1-cyclopropyl-6-fluoro-1,4-dihydro-4-oxoquinoline-3-carboxylic acid was active against MTB (strain H37Rv) and MDR-TB *in vitro*, with an MIC of 0.09 μ M in both strains. This compound also reduced the mycobacterial load in the lung and spleen of mice *in vivo* [9]. Quinoxaline derivatives were also potent against MTB. The high activities of some quinoxalines, mainly against resistant MTB strains, are well known and their biological effects are

related to an induction of hypoxia, which causes damage to genetic material [8]. Vicente et al. reported that the compound 7-methyl-3-(4'-fluoro)phenylquinoxaline-2-carbonitrile 1,4-di-N-oxide had potent anti-TB effects with an MIC of <0.2 μ g/mL and an SI of >500 [10].

The absence of new drugs to treat latent TB is another challenge that demands investigation. The next milestone for treating TB is to identify new drugs with potent sterilizing activity against replicating and non-replicating strains that can reduce the duration of chemotherapy [11,12]. Additionally, determining the association with other drugs that can decrease or inhibit TB's main drug-resistance mechanisms will improve treatment efficiency and cure rates.

References

1. World Health Organization, Global tuberculosis report, 2012, 8.
2. Lienhardt C, Cook SV, Burgos M, Yorke-Edwards V, Rigouts L, et al. (2011) Efficacy and safety of a 4-drug fixed-dose combination regimen compared with separate drugs for treatment of pulmonary tuberculosis: the Study C randomized controlled trial. *JAMA* 305: 1415-1423.
3. Caminero JA, Sotgiu G, Zumla A, Migliori GB (2010) Best drug treatment for multidrug-resistant and extensively drug-resistant tuberculosis. *Lancet Infect Dis* 10: 621-629.
4. Zumla A, Nahid P, Cole ST (2013) Advances in the development of new tuberculosis drugs and treatment regimens. *Nat Rev Drug Discov* 12: 388-404.
5. Cohen J (2013) Infectious disease. Approval of novel TB drug celebrated—with restraint. *Science* 339: 130.
6. Santos JL, Fernandes GFS, Bosquesi PL, Marino LB, Leite CQF et al. (2014) Synthetic Organic Compounds as Potential Antitubercular Drugs: A Review of the Progress Made in the Last 5 Years. In: Beatriz A, Lima DP (Org) *Recent Advances in the Synthesis of Organic Compounds to Combat Neglected Tropical Diseases*. Bentham Science Publishers.
7. Dutra LA, Melo TRF, Chin CM, Dos Santos JL (2012) Antitubercular drug discovery: the molecular modification as promise tool. *Int Res Pharm Pharmacol* 21: 1-9.
8. Koul A, Arnout E, Lounis N, Guillemont J, Andries K (2011) The challenge of new drug discovery for tuberculosis. *Nature* 469: 483-490.
9. Senthilkumar P, Dinakaran M, Yogeewari P, China A, Nagaraja V, et al. (2009) Antimycobacterial activities of novel fluoroquinolones. *Biomed Pharmacother* 63: 27-35.
10. Vieira M, Ferraz R, Fernandes R, Prudêncio C (2014) New quinoxalines with biological applications. *Biochem Pharmacol* 3: e152.
11. Vicente E, Pérez-Silanes S, Lima LM, Ancizu S, Burguete A, et al. (2009) Selective activity against *Mycobacterium tuberculosis* of new quinoxaline 1,4-di-N-oxides. *Bioorg Med Chem* 17: 385-389.
12. Duncan K, Barry CE 3rd (2004) Prospects for new antitubercular drugs. *Curr Opin Microbiol* 7: 460-465.

*Corresponding author: Jean Leandro dos Santos, Department of Drugs and Medicines, School of Pharmaceutical Science, State University of São Paulo, São Paulo, Brazil, Tel: 55-16-3301-6972; Fax: +55-16-3301-6960, E-mail: santosjl@cfar.unesp.br

Received May 28, 2014; Accepted May 29, 2014; Published June 03, 2014

Citation: dos Santos JL, Chin CM (2014) Pharmacology of Antitubercular Drugs: Challenges and Advances. *Biochem Pharmacol* 3: e164. doi:10.4172/2167-0501.1000e164

Copyright: © 2014 dos Santos JL, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.