



Draft Genome Sequence of *Bradyrhizobium elkanii* Tnp ϕ A 33, a Producer of Polyhydroxyalkanoates

Erica M. Lopes,^a Luciano T. Kishi,^a Camila C. Fernandes,^{a,d}
Fernanda Laroza Paganelli,^b Lucia M. C. Alves,^a Eliana G. M. Lemos,^{a,d}
Jackson A. Marcondes de Souza^c

Departamento de Tecnologia, Faculdade de Ciências Agrárias e Veterinárias, UNESP–Universidade Estadual Paulista, Jaboticabal, São Paulo, Brazil^a; University Medical Center Utrecht, Utrecht, the Netherlands^b; Departamento de Biologia Aplicada à Agropecuária, Faculdade de Ciências Agrárias e Veterinárias, UNESP–Universidade Estadual Paulista, Jaboticabal, São Paulo, Brazil^c; Faculdade de Ciências Agrárias e Veterinárias, Departamento de Tecnologia, Laboratório Multiusuário de Sequenciamento em Larga Escala e Expressão Gênica, UNESP–Universidade Estadual Paulista, Jaboticabal, São Paulo, Brazil^d

ABSTRACT The genus *Bradyrhizobium* comprises bacteria with the ability to form nitrogen-fixing symbioses with legumes. They are of great interest in agriculture, as well as for the production of biopolymers such as polyhydroxyalkanoates. Here, we report the draft genome assembly of *Bradyrhizobium elkanii* Tnp ϕ A 33 comprising 9 Mb, 1,124 contigs, and 9,418 open reading frames.

Members of the *Rhizobiaceae* and *Bradyrhizobiaceae* families have the ability to produce degradable biopolymers such as polyhydroxybutyrate (PHB), which can be used in the manufacture of biodegradable plastics (1). Studies based on the genome of the wild-type strain *B. elkanii* SEMIA 587 (2) have shown the potential for the synthesis and accumulation of PHB. The Tnp ϕ A transposon was randomly introduced in the genome of *B. elkanii* SEMIA 587, and the production of PHB was evaluated in comparison with the wild type. Cells were lysed for PHB extraction from *B. elkanii* Tnp ϕ A 33 (3). Marcondes et al. (4) viewed the expression of *phbC* and *glgA* genes and indicated the synthesis and accumulation of reserves of PHB and glycogen simultaneously. The connection between the storage of PHB and the tricarboxylic acid cycle demonstrates the importance of this polymer in the regulation of carbon balance. PHB probably also plays an important role during symbiosis with plants regarding: (i) energy-reserve offer for cell division during plant infection, (ii) protection for the nitrogenase complex enzyme, (iii) providing the reducing power for maintenance of O₂ diffusion in the absence of photosynthesis, (iv) recovering bacterial cells after their release from plants to the rhizosphere, and (v) increasing the survival time of the bacteria in the soil rhizosphere (1).

Genomic DNA from *B. elkanii* Tnp ϕ A 33 was extracted using a Wizard genomic DNA purification kit (Promega). The library was constructed using the Ion Xpress Plus fragment library kit (Thermo Fisher Scientific), using conditions recommended by the manufacturer. Single-read sequencing (1 × 200 bp) was performed on an Ion Torrent proton sequencer (Thermo Fisher Scientific) on a separate Ion PI chip, according to the manufacturer's protocols.

The DNA sequencing analysis allowed the partial identification of the *B. elkanii* mutant genome based on 22,962,787 reads sequenced and 18,907,631 reads showing Phred quality scores greater than 20 using PRINSEQ (<http://prinseq.sourceforge.net>). The clustering of contigs was performed with the SPAdes program (5), resulting in 1,124 contigs (9,542,959 bp), 63.7% GC content, 9,418 annotated open reading frames, and 77

Received 9 November 2016 Accepted 14 November 2016 Published 23 February 2017

Citation Lopes EM, Kishi LT, Fernandes CC, Paganelli FL, Alves LMC, Lemos EGM, de Souza JAM. 2017. Draft genome sequence of *Bradyrhizobium elkanii* Tnp ϕ A 33, a producer of polyhydroxyalkanoates. *Genome Announc* 5:e01502-16. <https://doi.org/10.1128/genomeA.01502-16>.

Copyright © 2017 Lopes et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Eliana G. M. Lemos, egerle@fcav.unesp.br.

E.M.L. and L.T.K. contributed equally to this article.

RNAs. Gene prediction and functional annotation of the draft genome were performed using the RAST server (6).

This mutant strain has revealed important genes for (i) the synthesis and accumulation of PHB, mostly *phbA*, *phbB*, and *phbC*, which represent biotechnological application (7); (ii) the processes of nodulation, *nodABC* (8, 9), and biological nitrogen fixation, *nifHDK* and *fixN*; and (iii) the amino acid metabolism, *glgA*, *glnA*, and *glnB* (4), among others. The group of genes described is very important for biopolymer industry and agriculture applications corroborating the versatility of fates for this mutant strain. Finally, this partial genome analysis has shown great similarity to the wild-type genome of *Bradyrhizobium elkanii* hosted in the NCBI databases (2).

Accession number(s). The draft genome sequence of *Bradyrhizobium elkanii* Tn*phoA* 33 was deposited at GenBank, NCBI, under accession number [MOXO00000000](https://doi.org/10.1128/JB.00563-12).

ACKNOWLEDGMENTS

This research was supported by the Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP) and by the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq).

REFERENCES

- Gjalt W, Huisman LLM 1999. Metabolic engineering of poly(3-hydroxyalkanoates): from DNA to plastic. *Microbiol Mol Biol Rev* 63:21–53.
- de Souza JA, Tieppo E, Magnani Gde S, Alves LM, Cardoso RL, Cruz LM, de Oliveira LF, Raittz RT, de Souza EM, Pedrosa Fde O, Lemos EG. 2012. Draft genome sequence of the nitrogen-fixing symbiotic bacterium *Bradyrhizobium elkanii* 587. *J Bacteriol* 194:3547–3548. <https://doi.org/10.1128/JB.00563-12>.
- Paganelli FL. 2009. Cloning and heterologous expression of genes responsible for synthesis of polyhydroxybutyrate em *Bradyrhizobium elkanii*. PhD Thesis. University of Campinas, Campinas, São Paulo, Brazil.
- Marcondes J, Lemos EGM 2011. DNA microarray applied to data mining of *Bradyrhizobium elkanii* genome and prospection of active genes, p 229–244. In Funatsu K (ed.), *Knowledge-oriented applications in data mining*. Intech, São Paulo, Brazil.
- Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Prjibelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <https://doi.org/10.1089/cmb.2012.0021>.
- Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, Formsma K, Gerdes S, Glass EM, Kubal M, Meyer F, Olsen GJ, Olson R, Osterman AL, Overbeek RA, McNeil LK, Paarmann D, Paczian T, Parrello B, Pusch GD, Reich C, Stevens R, Vassieva O, Vonstein V, Wilke A, Zagnitko O. 2008. The RAST server: rapid annotations using subsystems technology. *BMC Genomics* 9:75. <https://doi.org/10.1186/1471-2164-9-75>.
- Quelas JI, Mesa S, Mongiardini EJ, Jendrosseck D, Lodeiro AR. 2016. Regulation of polyhydroxybutyrate synthesis in the soil bacterium *Bradyrhizobium diazoefficiens*. *Appl Environ Microbiol* 82:4299–4308. <https://doi.org/10.1128/AEM.00757-16>.
- Tan HW, Weir BS, Carter N, Heenan PB, Ridgway HJ, James EK, Sprent JI, Young JPW, Andrews M. 2012. Rhizobia with 16S rRNA and nifH similar to *Mesorhizobium huakuii* but novel *recA*, *glnII*, *nodA* and *nodC* genes are symbionts of New Zealand *Carmichaelinae*. *PLoS One* 7:e47677. <https://doi.org/10.1371/journal.pone.0047677>.
- Handelsman J, Rondon MR, Brady SF, Clardy J, Goodman RM. 1998. Molecular biological access to the chemistry of unknown soil microbes: a new frontier for natural products. *Chem Biol* 5:R245–R249. [https://doi.org/10.1016/S1074-5521\(98\)90108-9](https://doi.org/10.1016/S1074-5521(98)90108-9).