

UNIVERSIDADE ESTADUAL PAULISTA "JÚLIO DE MESQUITA FILHO"

Campus de São José dos Campos Instituto de Ciência e Tecnologia







DANIELLE BLAZYS CORREA

INCREASED BURNED AREA IN THE PANTANAL OVER THE PAST TWO DECADES

DANIELLE BLAZYS CORREA

INCREASED BURNED AREA IN THE PANTANAL OVER THE PAST TWO

DECADES

Master's Thesis presented to the São Paulo State

University (Unesp), Institute of Science and

Technology, São José dos Campos; National

Center for Monitoring and Early Warning of

Natural Disasters (Cemaden) as part of the

requirements of the master's degree for the

Graduate Program in NATURAL DISASTERS.

Area: Natural Disasters. Research Line:

Instrumentation and data analysis.

Advisor: Prof. Enner Alcântara

Co-Advisor: Prof. Renata Libonati

São José dos Campos

2022

Instituto de Ciência e Tecnologia [internet]. Normalização de tese e dissertação [acesso em 2022]. Disponível em http://www.ict.unesp.br/biblioteca/normalizacao

Apresentação gráfica e normalização de acordo com as normas estabelecidas pelo Serviço de Normalização de Documentos da Seção Técnica de Referência e Atendimento ao Usuário e Documentação (STRAUD).

```
Correa, Danielle Blazys
Increased burned area in the Pantanal over the past two decades. / Danielle Blazys Correa. - São José dos Campos : [s.n.], 2022.
```

Dissertação (Mestrado em Desastres Naturais) - Pós-graduação em Desastres Naturais - Universidade Estadual Paulista (Unesp), Instituto de Ciência e Tecnologia; Centro Nacional de Monitoramento e Alertas de Desastres Naturais (Cemaden), São José dos Campos, 2022.

Orientador: Enner Herenio de Alcântara

Coorientadora: Renata Libonati

Burned areas. 2. Land use and land cover changes. 3. Tropical wetland.
 Extreme events. I. Alcântara, Enner Herenio de , orient. II. Libonati, Renata, coorient. III. Universidade Estadual Paulista (Unesp), Instituto de Ciência e Tecnologia, São José dos Campos. IV. Universidade Estadual Paulista 'Júlio de Mesquita Filho' - Unesp. V. Universidade Estadual Paulista (Unesp). VI. Centro Nacional de Monitoramento e Alertas de Desastres Naturais (Cemaden). VII. Título.

EXAMINATION BOARD

Enner Alcântara (Advisor)

São Paulo State University (Unesp)
Institute of Science and Technology
São José dos Campos

José Antônio Marengo Orsini (Board Member)

São Paulo State University (Cemaden)
Institute of Science and Technology
São José dos Campos

Fábio de Oliveira Roque (Board Member)
Federal University of Mato Grosso do Sul (UFMS)
Institute of Biosciences
Cidade Universitária, Campo Grande

São José dos Campos, March 2022

AKNOWLEDGEMENTS

The intensity of the last two years of my master's trajectory, especially when occurring during a hard and prolonged Pandemic, showed me once again how important it is to have special people around us, even if not physically. Therefore, I could not be more thankful for the key pieces that, in different ways, strengthened me to complete this stage of life.

I'm thankful to the UNESP and all its employees, that even during such a delicate moment for Brazilian science and education, stood up to keep the high-quality education standards, offering full support to students. In this context, a special thank you to my Advisor, Enner, who betted and trusted in me since the beginning, supported and encouraged my ideas, pushed me to keep ahead when hard moments appeared and were always available to help me solve my questions.

I also want to thank my fiancé Victor Hugo, my love, who is present during all my steps ever since we met, during my achievements and frustrations, celebrations and relaxing moments. He helps me to take life easier and appreciate it even more. Thank you for letting me share with you my dreams, insecurities, the daily small happy moments, and for encouraging me to face challenges with my head up saying how much you believe in my potential.

I want to thank my parents, Eliane and William, for the unconditional love and care, that through my whole life made everything easier and more comfortable, removed significant invisible stones from my path so that I could dedicate myself to studying what I chose and building my career with commitment the way I desired.

At last, I am grateful for all the sincere moments of listening and exchange shared with my friends and family. You are amazing.

All we have are us.

SUMMARY

ABSTRACT		
1	INTRODUCTION	5
2	ARTICLE	8
	2.1 Article – Blazys, D. C., Alcantara, E., Libonati, R., Massi, K. G., Park, E.	
	Increased burned area in the Pantanal over the past two decades.	8
	FINAL CONSIDERATIONS	
R	REFERENCES	25

BLAZYS, D.C. Increased burned area in the Pantanal over the past two decades. São José dos Campos. São Paulo State University (Unesp), Institute of Science and Technology; National Center for Monitoring and Early Warning of Natural Disasters (Cemaden), 2022.

ABSTRACT

Wildfires are behaving differently now if compared to other time in history in relation to frequency, intensity and affected ecosystems. In Brazil, unprecedent fires are being experienced in the last 10 years and to prevent and minimize similar disasters we must understand more about the natural and human influences on fires on each ecosystem as well as the gaps to fire combat strategies. The Brazilian Pantanal is the largest contiguous wetland in the world and a complex environmental system. In 2020 Pantanal experienced a catastrophic wildfire which burned out of control that shocked the world. The fire in Pantanal is expected to escalate given the increase in drought episodes, inadequate fire management strategies and weak environment regulations. In this study, we analyzed recent patterns and changes in fire frequency across the Pantanal based on land use and cover classes. The inter-annual variability of the fire and land cover changes between 2000 and 2021 was assessed using burned area (BA) from MCD64A1 V.6 product and land use and land cover (LULC) data from Landsat imagery. Our results showed that while smaller BA were more frequent for all LULC, forest and grassland classes represented much larger BA with lower frequencies. Cropland showed the smallest burned area among the LULC. Given the differences in the rates of recovery and regeneration after fire for different classes, Pantanal should be systematically monitored to develop a more effective fire combat strategy. We understand that our work demonstrates fundamental spatiotemporal clues to managers to strategically approach unusual fires with result of what periods were exceptional burned according to LULC classes, as observed for Forests in 2020 and an exponential burning growth in cropland during late dry (LD) season since 2014.

Keywords: Land use and land cover changes, Burned areas, Tropical wetland, Extreme events

1 INTRODUCTION

On 13th October 2020, The New York Times wrote: "The World's Largest Tropical Wetland Has Become an Inferno". The mentioned wetland was the Brazilian Pantanal, while the inferno refers to massive wildfires. This biome, globally known as a rich biodiversity neotropical wetland, had one third of its area affected by fires in 2020 (LIBONATI et al., 2020) and around 17 million vertebrates direct killed (TOMAS et al., 2022). It was considered the biggest fire event in history, with an increase of 376% in the last two decades and only 43% of the area unaffected (DAMASCENO-JUNIOR et al., 2021, GARCIA et al., 2021). The previous year (2019) was also marked by high numbers of burnings. In fact, an increasing trend of fire foci from 2000 to 2015 has been identified on South Pantanal (Mato Grosso do Sul State: Oliveira-Junior, 2020), with drought patterns also becoming more frequent (THIELEN et al. 2020, CARDOSO AND MARCUZZO, 2010).

However, fire is a phenomenon that has many faces and its effects can be beneficial or harmful to biodiversity conservancy and human lives depending on where and how they burn (HARDESTY et al. 2005). Fire shapes biogeochemical cycles in ecosystems, influencing soil nutrients availability, water quality and atmospheric composition (PEREIRA et al., 2018, DE OLIVEIRA et al., 2020), affecting directly and indirectly fauna and flora species diversity, abundance and distribution, beyond socioeconomic dimensions (PAUSAS AND PARR, 2018; STEENVOORDEN et al., 2019).

Fire prone ecosystems presents a framework where plants' physiology, phenology and metabolism has evolved through millions of years in order to keep its features with the respective fire regime, after affecting the corresponding secondary organisms in the food net (HARDESTY et al. 2005, PAUSAS AND PARR, 2018, DURIGAN et al., 2020). Reinforcing natural fire cycles are beneficial and life-sustaining in ecosystems that evolved with fire, as occurs with Brazilian savannah (Cerrado) (HARDESTY et al. 2005). According to Sommers et al, 2011, these fire regimes can be characterized by attributes that can be listed as patterns in burning season, fire interval and return, fire frequency, burned layer (ground, crown, understory/ sub-canopy), burned area, fire-intensity, duration, among others and depending on the context, only some of these aspects can be predominant in a fire regime characterization (ARRUDA et al., 2016). Grasslands, woodlands, savannas and wetlands ecosystems, for

example, are characterized by frequent, low-intensity surface fires that act to maintain an open structure allowing sunlight to penetrate (HARDESTY et al. 2005, FIDELIS et al. 2007).

On the other hand, fire sensitive environments, as tropical moist broadleaf forests of Amazon basin, do not present sufficient evolutionary traces to respond to this disturbance frequently. Therefore, wildfires (rare events of fire that usually takes large proportions) reflect changing phytophysiognomic aspects and environment's functions through time. It can entail some species diminishing and extinguishing, being replaced by others and leading to degradation and land cover conversion. Frequent, large, and intense fires were, until recently, rare events in these kinds of ecosystems (HARDESTY et al. 2005, ALENCAR et al, 2015). The current more frequent fires are primarily associated with human activity, used as a tool to clear land after deforestation or maintaining existing farmland and pasture (HARDESTY et al. 2005, BOWMAN et al., 2020). In turn when it occurs in drought periods, there is an increase in risk of fires (LIBONATI et al, 2021).

In this sense, general reasons for fire occurrences and flame intensity are related to the availability of elements involved in the combustion process being oxygen, flammable fuel load and heat. Climate conditions and its variability that leverage to oxygen depletion or availability or to dry biomass accumulation due evapotranspiration (as wind speed, air temperature, precipitation, pressure), beyond natural or anthropic sources of ignition, are key parameters to be monitored pre, during and after fires (JOLLY et al., 2015, LIBONATI et al., 2020). Air pressure and temperature gradients can boost wind speed (KARNAUSKAS et al., 2018) which can spread flames through biomass available as fuel, for example, expanding burned areas. Ecologists believe that fires are behavering differently now if compared with other time in history due ignitions by humans and human-induced climate change, especially in fire-sensitive tropical rainforests, retrofitting climate change by contributing with GHG emissions (HARDESTY et al. 2005, BOWMAN et al., 2020).

For a better delimitation of the multiple consequences of 2020's megafire in Pantanal, what exactly burned and has been burned inside Pantanal can be further explored. Once fire affects fauna and flora with different intensities and timescales, how it distributes organisms in Pantanal is not yet fully understood (JUNK AND CUNHA, 2005). More detailed research is needed on the weather conditions that fan fires, as well as the influences of ecology and management (LIBONATI et al., 2020). Information concerning fire in tropical wetlands are scant and the major fire studies come from savannahs and forest areas (DAMASCENO-

JUNIOR et al., 2021). In Brazil, most of fire research has been developed in the Cerrado Biome (PIVELLO et al., 2021)

Yet 2020 fires in Pantanal resulted in recent important advances in explaining fire dynamics in the biome and some of the crucial structural gaps in local fire management could then be already spotted, knowledge about the region's fire regime need to shore up regarding a multidisciplinary approach (LIBONATI et al., 2020). Comparisons between fire over different land use and land covers, considering particularities of availability of fuel loads, and fire seasons with evident inter-annual projected trends were not found in the literature. In addition, Libonati et al. (2020), Garcia et al. (2021), indicates that studies about Pantanal fires and associations with deforestation remain unclear. Understanding more about past and current variability in Pantanal fire occurrence in different land cover is central to predicting future fire frequencies, extensions, intensity, seasonality, and severity and to better anticipate new trends and disasters, as already raised for other Brazilian biomes.

Due to the predicted increase in drought episodes (THIELEN et al., 2020), inadequate fire management strategies, lax laws, and weak environment regulations (LIBONATI et al., 2020) we hypothesized a growing BA trend in Pantanal. In different seasons and land use and land cover (LULC) classes, we hypothesized that BA are increasing in wetland, forest, savannah, grassland, pasture, and cropland areas inside the Pantanal and relatively, in terms of the respective class area, it remains stable in rural managed portions as grassland (also used by traditional cattle ranching), pasture, and cropland classes. This work aims to analyze de fire occurrence in the Pantanal in the last two decades by land use and land cover class. The objectives were:

- i. Get time series of LULC of Pantanal from 2000 to 2021;
- ii. Quantify and compare historical monthly BA for different LULC in Pantanal;
- iii. Track trends of burned area for each LULC in Pantanal;
- iv. Map burn frequencies by season in Pantanal;
- v. Compare burn frequency among LULC.

2 ARTICLE

2.1 Article – Blazys, D. C., Alcantara, E., Libonati, R., Massi, K. G., Park, E. Increased burned area in the Pantanal over the past two decades. Science of the Total Environment 835 (2022) 155386. https://doi.org/10.1016/j.scitotenv.2022.155386.

Science of the Total Environment 835 (2022) 155386



Contents lists available at ScienceDirect

Science of the Total Environment

journal homepage: www.elsevier.com/locate/scitotenv



Increased burned area in the Pantanal over the past two decades

Danielle Blazys Correa ^a, Enner Alcântara ^{a,*}, Renata Libonati ^{b,d,e}, Klécia Gili Massi ^a, Edward Park ^c

- ^a Graduate Program in Natural Disasters, Unesp/Cemaden, São Paulo State University, São José dos Campos, Brazil
- Department of Meteorology, Federal University of Rio de Janeiro, Rio de Janeiro, Brazil National Institute of Education and Earth Observatory of Singapore, Nanyang Technological University, Singapore Instituto Dom Luiz (IDL), Faculdade de Ciências, Universidade de Lisboa, 1749-016 Lisbon, Portugal
- Centro de Estudos Florestais, Universidade de Lisboa, 1349-017 Lisbon, Portugal

HIGHLIGHTS


- Fire patterns in the Pantanal wetland in the last two decades were analyzed by mapping burned area.
- Among land covers, grassland was the most vulnerable to fire.
- Mega-fires in 2020 have occurred exceptionally within forest.
 Large scale synoptic view is necessary for
- better management of fires in Pantanal.

Editor: Paulo Pereira

ARTICLE INFO

Keywords: LULC changes Burned areas Tropical wetland

GRAPHICAL ABSTRACT



ABSTRACT

Wildfires are behaving differently now compared to other time in history in relation to frequency, intensity and affected ecosystems. In Brazil, unprecedented fires are being experienced in the last decade. Thus, to prevent and minimize similar disasters, we must better understand the natural and human drivers of such extreme events. The Brazilian Pantanal is the largest contiguous wetland in the world and a complex environmental system. In 2020, Pantanal experienced catastrophic wildfires due to the synergy between climate, inadequate fire management strategies and weak environmental regulations. In this study, we analyzed recent patterns and changes in fire behavior across the Pantanal based on land use and cover (LULC) classes. The inter-annual variability of the fire and land cover changes between 2000 and 2021 was assessed using BA from MCD64A1 V.6 product and LULC data from Landsat satellite. Our work reveals that fires in the Pantanal over the last two decades tended to occur more frequently in grassland than in others land cover types, but the 2020 fires have preferentially burned forest regions. Large fire patches are more frequent in forest and grasslands; in contrast, croplands exhibit small patches. The results highlight that a broad scale analysis does not reflect distinct localized patterns, thus stratified and refined studies are required. Our work contributes as a first step to disentangling the role of anthropogenic-related drivers, namely LUIC changes, in shaping the fire regime in the Pantanal biome. This is crucial not only to predict future fire activity but also to guide appropriated fire management in the region.

1. Introduction

Over the past few decades, human activities and climate change have been contributing to more frequent and intense spikes in forest fire activity

* Corresponding author.

E-mail address: enner.alcantara@unesp.br (E. Alcântara).

around the world, leading to harmful impacts on the environment, wildlife, human health, economy, and infrastructure (Bowman et al., 2020). More recently, not only are fire-prone ecosystems experiencing more intense wildfires, but areas that typically burn less are also facing large-scale burning events (UNEP, 2022). The 2019 and 2020 fire seasons are being considered the worst in recent decades in many parts of the world-for instance, in Australia (Boer et al., 2020), California (Li and Banerjee, 2021), the Arctic

http://dx.doi.org/10.1016/j.scitotenv.2022.155386

Received 6 October 2021; Received in revised form 13 April 2022; Accepted 15 April 2022 Available online 21 April 2022

0048-9697/© 2022 Elsevier B.V. All rights reserved.

(Witze, 2020) and the Pantanal (Libonati et al., 2020). In 2020, the Brazilian Pantanal, the largest contiguous wetland in the world had one third of its area affected by fires (Libonati et al., 2020). It was considered the largest fire event in history, with a 376% increase in the extent of fires, compared to the annual average from the last two decades (Damasceno-Junior et al., 2021; Garcia et al., 2021).

Fire shapes biological and physical dimensions of ecosystems (Bowman et al., 2020), as well as socioeconomic frameworks (Pausas and Parr, 2018; Steenvoorden et al., 2019). Fire effects can be beneficial or harmful to biodiversity conservation and human lives depending on where, when, and how frequent ecosystems burn: if fires reinforce natural regimes, whether naturally or human ignited, they are recognized as good and lifesustaining in environments that have evolved with them (Hardesty et al., 2005). For instance, fires have been found to be important in maintaining vegetation diversity, structure, and functions in fire prone ecosystems such as savannas (Fidelis et al., 2007; Durigan et al., 2020).

Conversely, fires can be detrimental in fire sensitive formations (rainforests) (Prestes et al., 2020; Sansevero et al., 2020). Although the Brazilian Pantanal formations are classified as savannas, they are directly influenced by other biomes (IBGE, 2012), i.e., savannas of Cerrado and Atlantic and Amazon rainforests. For this reason, this biome is a mix between fire prone and fire sensitive ecosystems. Fire-sensitive sites in the Pantanal are therefore negatively affected by burnings through biodiversity loss (Miranda, 2010; Martins et al., 2022). As such, the future of Pantanal depends largely on proper fire management practices (Berlinck et al., 2022).

Fire occurrences can be attributed to both natural and anthropogenic causes with land use change being recently identified as a major driver in Brazil (Pivello et al., 2021). Native grasslands of Pantanal have been historically used for low intensity cattle ranching (embrapa.br/pantanal); since the 2000s, native grasses are being substituted by exotic grasses in pastures or being converted into croplands (mainly soybean) (Pott and Pott, 2004). The transition from natural vegetation to agriculture and pasture has led to high concentration of fire foci and increased risk of fire (Marques et al., 2021). However, fire studies focusing on the influence of land cover change remains scarce in this region, except for the recent work of Kumar et al. (2022) who used coarse land cover and burned area data to understand the main causes of the 2020 Pantanal fires.

A deeper understanding about past and current fire variability in the Pantanal considering different land covers is central to predicting future fire behavior and anticipating new trends and disasters (Pivello et al., 2021). Considering multiple land covers and land use changes over Pantanal, together with other important drive factors such as droughtheat waves episodes (Thielen et al., 2020; Thielen et al., 2021; Libonati et al., 2022) and regional fire management and regulations, is therefore crucial to developing an integrated understanding about current and future fire regime in the region (Libonati et al., 2020). However, such a comprehensive and long-term representation of fire occurrence per land use and land cover type (LULC) in the Pantanal is still not available.

The goal of this study is to analyze the recent observed patterns and changes in fire behavior over the Pantanal according to land use and land cover classes. Here, we analyzed long-term and annual variability of fire and land cover changes from the end of 2000 to the beginning of 2021 using burned area (BA) from MCD64A1 V.6 product and LULC classes from 2000 until 2019. First, we assessed fire frequency over the region by considering inter and intra-annual variability per LULC classes. Then, we examined trends to infer the role of LULC on Pantanal fire dynamics, paying particular attention to the year 2020 due the catastrophic fire events registered that year (Libonati et al., 2020; Damasceno-Junior et al., 2021; Garcia et al., 2021).

2. Materials and methods

2.1. Study site

Pantanal is the world's largest contiguous wetland at approximately $150,\!000\,\mathrm{km}^2$ and is located in the interior of the South America continent, covering part of Mato Grosso (35%) and Mato Grosso do Sul (65%) states in

Brazil (Fig. 1 a. and b.). Known as "Pantanal complex", this biome is influenced by the Amazon, Cerrado (Brazilian Savanna) and Atlantic Forest biomes in its north, east and south, respectively (Fig. 1 b., Marengo et al., 2021). On the west, the Brazilian Pantanal borders Bolivia and Paraguay in a region known as Chaco. The current LULC distribution (Fig. 1c.) shows pasture class advancing from east to west with small portions of cropland and other non-vegetated areas. The main continuous forestland and savanna classes are found in central to north portions while wetland class is concentrated in the west.

According to the Köppen's climate Classification (Koppen, 1948, Peel et al., 2007, Embrapa, 2020), the Tropical Wet-Dry or Tropical Savanna climate is predominant in the Pantanal, with cold and dry months varying between May to October, and hot and wet months varying between November and April. Different regions typically experience a flooding peak which lags the precipitation peak (Fig. 1 e.), and this timing can be as much as three months. As for rivers such as Miranda and Cuiabá, their flooding peaks occur soon after the peak of rain. Considering large-scale climate phenomena, Thielen et al. (2020) found oscillations in sea surface temperature (SST) of the Pacific and Atlantic Oceans as the main drivers for rainfall variability in these basins.

While the hydrology of the Pantanal landscape is driven by multiple factors, flood pulse is considered the main driver (Ivory et al., 2019). The Pantanal acts as a large reservoir in Paraguay River basin, where its flat terrain (Fig. 1 d.), varying soil classes (mainly Gleyic Podzols in the center, Luvic Planossols in the south and close to main rivers and Dystric Plintosols in the north (Embrapa Solos, 2017) and regional hydroclimate conditions influence floods in the area (Fernandes et al., 2007; Assine et al., 2015; Marengo et al., 2021). Pluriannual humidity and rainfall cycles can also result in severe floods or pronounced dry seasons that then influence flooding patterns (Damasceno-Junior et al., 2021; Garcia et al., 2021; Marengo et al., 2021). Processes that maintain Pantanal's landscape and wetland functioning is also likely to be disrupted by changes in rainfall regime and the flood pulse (Ivory et al., 2019). Increased sedimentation over a large area in the last 30 years that drastically modified the Taquari megafan, the largest alluvial fan comprised on Paraguay trunk river located in the central portion of the Pantanal and representing almost one third of the biome (Assine, 2005), is probably also associated with increased vulnerability to deforestation

The geographical location, difficult access, low soil nutrients availability, periodic floods, fires and traditional low-intensity cattle ranching in the natural plane grasslands since the 17th century (the current main economic activity) contributed to keeping the Pantanal's population density low, making it worthy of preservation (Junk and Cunha, 2005; Junk et al., 2006; Machado and da Costa, 2018). The implementation of development programs, crossroads, small hydroelectric reservoirs, and mining since the mid of 1970 and 1980, in addition to intensive farming in the Upper Paraguay Basin plateau, contributed to the degradation of natural vegetation and biophysical stability (Junk and Cunha, 2005; Alho and Sabino, 2011). It comprises almost 2000 km² of protected areas considered as UNESCO Natural World Heritage Site, and encompasses three Ramsar Sites, due to its ecological significance and high biodiversity. However, the expansion of agriculture frontiers in the Cerrado and Amazon biomes, exert a strong influence on natural resources on its surroundings (Nobre, 2010; Guerra et al., 2020) and the Pantanal biome susceptibility to changes appears again in this regional context (Hamilton, 2010).

2.2. Land use and land cover (LULC) data

The time series of LULC class data is an annual 30 m product from MapBiomas Project (Souza et al., 2020). The LULC maps from Collection 5 were made from Landsat imagery using the Random Forest machine learning classification process on Google Earth Engine cloud processing (see more at https://mapbiomas.org/en/methodology-overview). For Pantanal biome, the making of LULC mosaics prioritized Landsat scenes of dry period (May to August) by MapBiomas and the reported overall accuracy of Brazil classification is 89% (Souza et al., 2020).




Fig. 1. Overview of the Pantanal biome. (a) Location of the Pantanal in South America (b) Delimitation of the Upper Paraguay Basin, highlighting the Brazilian biomes. (c) Land use and land cover (LULC) studied classes in 2019 and gauge stations locations. (d) Elevation from ASTER-GDEM of the Pantanal and its surroundings. (e) Historical Monthly averages of Rainfall/pixel and water level at Leverger (light gray: 15°52′12.0″S; 56′04′37.2″W) and Ladario Gauge Stations (dark gray:19° 0′ 7.2″S; 57° 35″42.0 W), from November 2000 to January 2021.

The time range selected from MapBiomas was from 2000 to 2019, where 2000 was the earliest year available in the dataset. For the purpose of this research, the original LULC data was reclassified from 14 to 7 classes for each year, according to the amount of biomass availability or fire load material classification proposed by Shimabukuro et al., 2020. The converted classes with its original attributes' values are as follow: Forest (3 and 9), Humid Areas (11 and 33), Savanna-Shrubland (4), Grassland (12), Pasture (15) and Cropland (20, 39 and 41). Other non-vegetated areas (attributes 0, 24, 25 and 30) were not considered in this analysis. The meaning of the Attributes values can be found on https://mapbiomas.org/en/codigos-de-legenda?cama_set_language=en. Fig. 1 c. is an example of the final reclassified 2019 map.

2.3. Burned area (BA) data

Time series of BA data was acquired from MCD64A1 Version 6, a monthly global gridded 500 m product (Giglio et al., 2015), derived from a composition between processed MODIS Surface Reflectance imagery coupled with MODIS active fire observations. "BurnDate" band was used, and the original values assigned for each pixel are the Julian days of the year on which the burn emerged. The date range went from November 2000 to January 2021 (Fig. 2a).

Precipitation (PREC) data was obtained as a supporting variable as is considered an important parameter to infer drought magnitude and fuel load availability assessment (Gomes et al., 2017; Franke et al., 2018).

D.B. Correa et al.

Precipitation data was retrieved from ERA5-Land monthly average, a reanalysis dataset with 0.1 arc degrees resolution (approximately 11.1 km at equator) produced by the European Centre for Medium-Range Weather Forecasts (ECMWF) (Sabater, 2019). The selected date range for PREC is equal to MODIS BA and is also originally a global gridded product.

2.4. Remote sensing data processing

The Google Earth Engine (GEE) environment was used to obtain, prepare, and process the data in the Pantanal perimeter (IBGE, 2019). The global products were available as collections while National Mapbiomas dataset, as assets. Monthly variables (BA, PREC) were extracted for each LULC class by an iterative process. These monthly variables were subsequently filtered by and indexed with the LULC class and its area from November of a year to October of the following year (due to the methodology applied by MapBiomas on LULC classification). Beyond 2019, LULC map from year 2019 was used as data was unavailable. The PREC grid spatial layers were monthly reduced by each LULC using averages. The detailed authorial code for BA accountability by LULC over the months can be accessed through the following link: https://code.earthengine.google.com/

a0ab1632e4ae6cf36cc72d415a78be69. A similar code was employed to PREC variable (https://code.earthengine.google.com/521cfe6c3a9d9953bff2c52b86b8596c).

For the fire recurrence maps, the monthly BA data layers were stacked, and the pixel overlap was spatially accounted for. A reduced map covering the 243 burned months was divided by seasons (Koppen, 1948; Peel et al., 2007) late wet (LW), early dry (ED), late dry (LD) and early wet (EW) which correspond to the union of February to April, May to July, August to October and November to January, respectively. Seasons were defined based on total burning area in biome and Aw (or namely, Tropical Savanna/Wet-dry climate) Koppen's classification. Trimestral burn frequency maps are bivariate, including the year of last burn spatialized information in its viewing.

2.5. Statistical and spatial analysis

Monthly Absolute BA (ABA, km²) by LULC, as well as Relative BA (RBA, %) by the respective yearly LULC area class were represented by time series

graphics (zero values were masked). The ABA data corresponds to the monthly burned area by each LULC in each year of analysis and the RBA data is the ABA data divided by each LULC and then multiplied by 100.

Monthly BA time series (absolute and relative) by LULC were tested to identify for autocorrelation as well as seasonality characteristics. For these, Durbin Watson (D) and Augmented Dickey-Fuller (ADF) tests were used, respectively, considering p-value <0.05. D test is widely used to test autocorrelation in non-lagged dependent variables and when size sample is reduced. It ranges from 0 to 4, and when values are close to 2, there is no autocorrelation; when close to 0, there is a positive autocorrelation; when close to 4, there is a negative autocorrelation (Durbin and Watson, upon the close to 4, there is a negative autocorrelation (Durbin and Watson, and presence of seasonality over BA timeseries (Fuller, 1996). In this case, the null hypothesis is that the time series possesses a unit root and is non-stationary. When seasonality is presented, the series is not stationary.

ADF test was applied on time series ranging from November 2000 until January 2020 and also from November 2000 until January 2021. Since the year 2020 was a clear outlier (Garcia et al., 2021), we opted to exclude this year in the ADF tests in the Mann Kendall tests (more details of the last below). Hence, considering the three elements that defines a timeseries as stationary, if the shorter time series (until January 2020) was confirmed as stationary (p-value < 0.05), then it would be inferred that the non-stationarity in the longer time series (until January 2021) would be a result of the instability due to the influence of the mean and/or standard deviation - and not due to seasonality. Both tests, D and ADF, were supported by the Statsmodel library on Python (Fig. 2 b.).

Due to the series characteristics obtained from the aforementioned tests, BA (absolute and relative) trend analysis tests were performed using Yue and Wang Modified Mann Kendall test, appropriated for serial autocorrelation (Yue and Wang, 2004). The null hypothesis to be tested is that there are no trends, with p-value < 0.05. Lastly, series trends for the variables by LULC were confirmed by Theil-Sen Robust Linear Regression (TS) (Sen, 1968; Theil, 1992, Helsel and Hirsch, 2002).

Theil-Sen, also known as Kendall-Theil robust line, is a median-based and non-parametric estimator that does not make assumptions about the dataset distribution and is insensitive to outliers. It has been applied in




Fig. 2. Methodology flowchart. a. Temporal coverage of each used dataset: Land use and cover (LULC), burned area (BA) and Precipitation (PREC). b. Workflow on the used processing environments.

D.B. Correa et al.

many hydrological and environmental studies (Somorowska, 2016; Alves et al., 2021). As a linear fit estimator, it returns the slope (the degree of line decline of units/time, in this case) and y intercept values (studied monthly variables values in time zero), compounding the classical regression line equation ($y = \alpha x + b$). Both statistical analyses were performed with pyMannKendall Python package (Hussain and Mahmud, 2019). LULC classes' monthly variables were aggregated by trimester as averages for graphical representation.

3. Results

3.1. LULC changes

LULC in Pantanal composed mostly of forestland, humid areas, shrubland-savanna, grassland, pasture, and cropland (Fig. 1 c.); the LULC data from 2000 to 2019 showed a consistent increase (51%) in pasture area from 15,497.60 km² to 23,384.30 km² (Fig. 3) while croplands exhibited a small increase and savannah areas decreased. Other landcover types showed high variability between years. Forestland for instance, showed the lowest area variability.

Burned area time series by LULC.

The burned areas were not uniformly distributed across LULC classes, time, and seasons, across the past 20 years in Pantanal (Fig. 4). Major monthly ABA were observed in September of 2020, August of 2005, August of 2020, August of 2001 and September of 2007 in different areas of the biome, respectively ranging from 9.43% to 5.6% of the total area of the biome.

Grasslands had the largest ABA among land use classes for 2005 and 2001 (August) and in 2020 (September) with 6903, 5305 and 5887 km² respectively (Fig. 4 a. and b.). This represented 53.3, 54.9 and 39.9% of total burnings respectively, and made up between 3.7 and 4.6% of the biome area. Temporally, 2020 (August) and 2007 (September) were years with higher ABAs (especially forest and grasslands). In addition, 2002 and 2019 had longer burning seasons: from August to November, totaling to 26,767 and 19,600 km² respectively. Although September and October saw the highest ABA, fire was observed during the whole year in the Pantanal.

BA patterns look different when using a relative scale to assess BA patterns for each land use and land cover class. In 2020, forest, humid areas, and pasture responded with a higher proportion of burned areas (Fig. 4 b). In 2005, grassland, pasture, and savanna had a higher proportion of burned areas instead. Forest presented larger monthly burned area values in 2020 (August and September) and 2005 (August), humid areas in 2020 (from July to October), 2019 (August to October), 2012 (August to September) and 2002 (November), savanna in 2004 (September) and 2005 (August), grasslands in 2005, 2001 and 2020 (August for the first two and September for the last), pasture in 2020, followed by 2005 and



Fig. 3. LULC changes in Pantanal from 2000 to 2019. Source: MapBiomas

2002 (September for the first, August for the last two) and cropland in 2015 (November) and other months in the LW season in 2016, 2018 and 2019. Relatively, forests, grasslands and croplands were classes with higher turned areas (Fig. 4 c), especially during dry season months (August and September) and pastures, with lower relative burned area than the others LUIC.

The average yearly variation of absolute burned area, relative burned area, and monthly average precipitation during the ED season is shown in Fig. 5. In all LULC classes, burned areas from 2014 to 2016 were lower compared to other years. In the same years, the highest precipitation in the time series was observed. Shrubland-savanna, grassland and pasture showed larger burned area in 2002, 2005 and 2020. Forest showed low values of burned area (<75 km²) for most of the entire time series, except in 2020. Cropland on the other hand, did not show increased burned area in 2020.

During the LD season (Fig. 6) the burned areas increased, compared to the ED season (Fig. 5). For instance, in 2002, forest showed an increase from approximately 75 km² (ED season) of burned area to $900 \, \text{km}^2$ (LD season). During the LD season, precipitation variability was also larger compared to the ED season, i.e., from 2014 to 2016, precipitation in the LD season was lower than in the same period of ED season.

The mega fire of 2020 was pronounced for almost every LULC, except for shrubland-savanna, because the burned area in 2002 was higher than 2020. The same analysis for LW season and EW can be found in the S1 and S2.


For both shorter (Nov/2000 until Jan/2020) and longer time series (Nov/2000 until Jan/2021), only cropland presented a significant increase in absolute and relative burned areas (Table 1). The results in Table 1 shows that, only cropland indicated signs of increased burned area during the past twenty years

Over the last years, fire occurrences in forest, savanna, grassland and pasture classes have been decreasing, while the fire occurrences for the cropland have been increasing. However, in 2020, a different trend signal was observed for the forest class and as well as for the entire biome; the other classes (savanna, grassland and pasture) appeared less negative than the results excluding this year.

Most of Pantanal was burnt at least once in the last twenty years (Fig. 7). The map showed that across the 243 analyzed months, the maximum burn frequency reached 19 times that of fire recurrence in the same area. It can be also understood as a mean of almost one burn per year in most of the regions. Larger values were seen mainly at southwest Pantanal. The higher values were also mostly found over the current grassland. Conversely, smaller and low frequent burnt patches were seen in central and southeast portion of Pantanal. Some portions of Pantanal remain fire-free.

S3 illustrates the fire frequency per burned area (km²) per LULC. It revealed that smaller burned areas were frequent for all LULC classes. Large burned areas were observed in forest and grassland at a lower frequency. Cropland showed the smallest burned area among all LULC classes. For grassland, an area of 11,457.10 km², representing 20.8% of the entire Pantanal, burned once during the past twenty years while an area of 0.32 km² (0.001% of the grassland area) burned at least one time in each year. For cropland, an area of 40.11 km² (18.9% of the cropland area) burned only once during the entire time series and 0.02 km² (0.01% of the cropland area) burned 8 times during the past two decades. For humid class, the biggest area burned amounted to 5360.94 km² (18.7% of the cover class), which burned once in the entire study period; on the other hand, small burned areas of 0.09 km² (representing 0.0003%) in this class, burned 19 times.

Fires were concentrated on western side during the LW season and was observed to spread towards the east in the following seasons, reaching its peak in the LD season (Fig. 8). During LW season, recent burns appeared more extensive than other seasons, where older burned areas are greater. In LD season, higher values of burn frequency were present, especially in the southern and northeast portions of the biome. In this season, recent low frequency burns were seen in the north portion. In addition, old fires (from 2000) generally covered a smaller area than recent fires (2021), especially during the LD season. Overall, small burned areas were more frequent than large burned areas for all land use across all seasons. Forest and



D.B. Correa et al. Science of the Total Environment 835 (2022) 155386




Fig. 5. Yearly distribution of absolute BA average (left axis - green), relative BA average (right axis - yellow) and monthly precipitation average (right axis - blue) in ED season (from May to July) by LULC. Green hue of colour bars indicates the total area accounted yearly by LULC. Red bars are the constant Pantanal biome total area according to IBGE, 2019 dataset. Thin gray bars correspond to standard deviation (SD) of absolute BA. (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

grassland represented larger burned areas, while cropland represented the smaller ones (S3).

4. Discussion

Pantanal is a complex ecosystem composed by grasslands, savannas, forests and humid areas. Fire has played an important role in sustaining many of these ecosystems. For grasslands and savannas, burning is an important

ecological factor (Hardesty et al., 2005), and appropriate fire management protocol which needs to be undertaken (Pivello, 2011; Pivello et al., 2021). Grasslands usually accumulate an excess of dry grass and when an ignition occurs, naturally or by humans, fire easily spread around the LD season (Soriano et al., 2020).

As highlighted by Souza et al. (2020), cattle ranching and sugarcane expansion are driving the suppression of Pantanal natural vegetation grassland and extensive wetlands. The expansion of cattle ranches was favored

Fig. 4. Time series graphics with Monthly BA by LULC Class from November/2000 to January/2021 (zero values were masked). (a) Absolute values in km², legends scale common between classes. (b) Absolute values in km², maximum and minimum legends in relative scale for each class. (c) Relative values: relative to the respective LULC class series.




Fig. 6. As in Fig. 5 but for LD season (from August to October). Red bars are the constant Pantanal biome total area according to IBGE, 2019 dataset. (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

by a sequence of dry years from 1991 to 2004 (Mourão et al., 2010) and this pattern of pressure that resulted in LULC conversion is similar to the anthropic activities practiced on elevated basin portions (Guerra et al., 2020). Coutinho et al. (2016) showed that sugar cane production in the Pantanal expanded by 48% and agriculture by 39% from 2001 to 2013.

Fires can be detrimental for other parts of Pantanal, such as the tropical forest. Surface fires in tropical forests usually burn litter, duff and the understory grasses and herbs. They may cause extensive top-kill in small trees (Hoffmann et al., 2009) and change the composition, structure and function of forest ecosystems (Prestes et al., 2020; Sansevero at al., 2020). Undoubtedly, fires in the Pantanal forests peaked in 2020, and is mainly responsible for the increased burned area values showed in Table 1 (in the "total" column).

Our results of burned areas for the last 20 years across different LULC classes in the Pantanal showed that the fires of 2020 was a significant

event even though the biome has been continually burning for the past two decades. Most of the Pantanal has burnt at least once during the study period, especially during the dry season. In addition, the results suggest that small patches of burning were frequent in the last 20 years.

The year of 2020 was noted as one of the years in this study period that had a comparatively large burned area, which corroborates with previous studies (Garcia et al., 2021; Kumar et al., 2022). This is likely due to severe droughts in the region caused by a decrease in precipitation (Fig. 6; S1 and S2). This finding was supported by Marengo et al. (2021), Lázaro et al. (2020) and Thielen et al. (2021) which reported that precipitation and river levels have been decreasing in 2020 and its preceding years (2018 and 2019). Lower precipitation values after 2015 are also shown on our studies for ED season (Fig. 5) and after 2018 for LD season (Fig. 6).

Marengo et al. (2021) showed that in 2020, Pantanal experienced a persistent extreme drought with 60% less rain than a normal year, due to an

Table 1

Results of linear regression params BAs time series by LULC applied with SciPy library (95% confidence intervals considered). Slope red values indicates a positive slope (BA increasing), p-value < 0.05 are highlighted in bold. (a) time series ranging from Nov/2000 until Jan/2021. (b) time series ranging from Nov/2000 until Jan/2020.

(a)	Nov/2000-Jan/2021 Original Theil Sen trend Modified Mann Kendall Test Yue Wang (p-value < 0.05)								
		Forest	Savanna	Humid	Grassland	Pasture	Cropland	Total	
C1 (-)	ABA (km²/yr)	0.004	-0.018	0.700	-0.460	-0.069	0.000	0.157	
Slope (α)	RBA (%/yr)	0.00001	-0.00011	0.00215	-0.00077	-0.00047	0.00000	0.00018	
	ABA (km ²)	7.49	3.76	44.10	84.65	7.18	0.00	170.64	
Intercept (b)	RBA (%)	0.024	0.022	0.135	0.154	0.047	0.000	0.112	
p-Value	ABA	0.921	0.390	0.212	0.395	0.264	0.031	0.867	
	RBA	0.921	0.381	0.211	0.424	0.238	0.034	0.779	
(b)	Nov/2000-Jan/2020 Original Theil Sen trend Modified Mann Kendall Test Yue Wang (p-value < 0.05)								
		Forest	Savanna	Humid	Grassland	Pasture	Cropland	Total	
Gl ()	ABA (km ² /yr)	- 0.069	-0.040	0.019	-1.312	-0.125	0.000	-1.473	
Slope (α)	RBA (%/yr)	-0.0002	-0.0002	0.0001	-0.0024	-0.0008	0.0000	-0.0010	
	ABA (km ²)	6.83	3.53	46.91	83.51	7.50	0.00	165.22	
Intercept (b)	RBA (%)	0.022	0.021	0.144	0.152	0.048	0.000	0.109	
** 1	ABA	0.217	0.234	0.838	0.108	0.164	0.048	0.268	
p-Value	RBA	0.217	0.234	0.838	0.108	0.164	0.048	0.268	

anomalous warming of the tropical north Atlantic. As a result, unprecedented wildfires ripped through the region even in January and February, despite it being the wet season (Marengo et al., 2021). Garcia et al. (2021) also noted a reduced moisture inflow coming from the Amazon region in 2020 which could have exacerbated the drought. Our results showed that extreme droughts in 2020 and 2005 (Figs. 5 and 6) affecting all LULC classes coincided with great fire events reported in other studies (Kumar et al., 2022; Garcia et al., 2021). Coupled with 2020's climate and biomass conditions, wildfires in Pantanal reached huge proportions.

2020's fire season was exceptional, but conditions that led to this extreme event, such as increase of ignition sources, biomass availability, drier and warmer climate, are becoming increasingly common in the Pantanal (Libonati et al., 2020). For instance, Libonati et al. (2022) showed that the recent warming over this region since 1980 is around four times greater than the average global temperature increase. Decreasing trends in air humidity and in precipitation, together with higher temperatures, are increasing fire danger over the last 40 years. Recently, Kumar et al. (2022) showed that anthropogenic-caused fires are exacerbating climate effects on natural ecosystem within the Pantanal. On the Pantanal portion of Mato Grosso do Sul State, Oliveira-Júnior et al. (2020) had already identified a positive trend of fire foci from 2000 to 2015.

The Brazilian Pantanal mainly composes savannas (IBGE, 2012) and it is also closely situated to the other biomes, i.e., savannas of Cerrado and the Atlantic and the Amazon rainforests. Deforestation in the Amazon forests have affected rainfall and temperature regimes in the Pantanal region (Amaral e Silva et al., 2020), increasing the Pantanal's vulnerability to fire (Bergier et al., 2018).

However, long-term consequences of these fire occurrences on biodiversity and ecological services remain largely unknown (Libonati et al., 2020; Garcia et al., 2021; Damasceno-Junior et al., 2021). The implication of fires on the distribution of flora and fauna across Pantanal is yet not fully understood (Junk and Cunha, 2005). A recent effort revealed that around 17,000 million vertebrates were directly killed by the fires in the Pantanal during 2020 (Tomas et al., 2022). However, the authors point out that this value may be underestimated. For some areas affected by the fires during 2020 in the Pantanal, Martins et al. (2022) estimated the cost of post-fire restoration to be around 123 million USD.

Between other global and long coverage BA datasets available online, MCD64A1 shows important gains even with its gross spatial resolution of 500 m. Its inputs operate on short-wave infrared (SWIR) electromagnetic spectrum region and with a daily revisit. These qualities makes it a good data source, especially under tropical conditions where post-fire signals are swiftly lost by sensors, and cloud persistence limits imagery quality (Chuvieco et al., 2019). Further, MCD64 collections were robustly validated for Brazil under varying spatial and climate conditions (Rodrigues et al., 2019). Chuvieco et al. (2018) consider that these main different

global BA products complement each other as also highlighted by Long et al. (2019). Depending on the size of studied region, very high spatial resolution demands more computer capacity processing ant it can result also in impractical processing time (Long et al., 2019). MCD64A1 is broadly used over the globe, and it favors comparisons between regions around the world. Its cloud processing due the dataset availability on GEE platform also, makes convenient processing for our regional scale study.

According to Damasceno-Junior et al. (2021), fire regime in Pantanal is closely linked to the annual and pluriannual flood pulse and from August to October, the biomass produced during the flooding period becomes available as fuel for burning during the next dry season. As shown during the 2020 burns, Paraguay River, the main river of Pantanal (Fig. 1 b.), recorded the lowest water level during flooding periods in the last 47 years (Marengo et al., 2021). Consequently, most of the 2020 wildfires occurred in this flood zone (Damasceno-Junior et al., 2021; Garcia et al., 2021; Kumar et al., 2022). Nevertheless, an interannual to biannual/triannual scale burning cycles was also observed between grassland and humid areas formations. It coincides with respective LULC change due boundaries reshaping in transition zones and flooding dynamic alterations in drierwetter years seen in works of Damasceno-Junior et al. (2021) and Thielen et al. (2020).

The results observed in Fig. 8 highlights the combined influence of the flood pulse and rainfall regime in modulating the fire regime. During the LD season (Fig. 8c), for instance, it is highlighted that the occurrence of fire is increasingly frequent and consistent with the increase of more frequent extreme drought events. Based on our results, we hypothesized a growing trend in burned areas in all LULC in Pantanal, especially in grasslands, savannas and pastures. We have found an increase in burned areas, but in cropland land use class. In general, cropland expansion does not imply an increase in burnings since it takes place in older pasture areas, already converted from natural to anthropic stage (Chuvieco et al., 2019). The land use changes linked to deforestation of native vegetation traditionally involve fire for opening new pasture areas (Archibald et al., 2013). As such, deforestation and the increase in large agribusiness properties in Pantanal could be linked to these recent fire events (Mota et al., 2019; Margues et al., 2021). The higher BAs proportions found on native vegetation compared to managed areas also highlight the lack of adequate fire management.

Taking into consideration the Pantanal's flooding cycle along Paraguay River, where flooding reaches peak from May to July, near the months with lowest rainfall values (from June to August), the floodplain cannot catch fire during dry season except in dry years. In the case of dryness asynchrony, a larger amount of accumulated biomass produced during the last flooding will remain available and exposed as fuel – mainly the histosols (organic soils), that are wet and not available to fire in "normal" years (, Holden et al., 2018; Garcia et al., 2021).




Fig. 7. 20 years of burn frequency map from monthly MCD64A1 dataset, 500 m resolution.

Some works also detected that the drought patterns in Pantanal trended towards becoming more frequent (Thielen et al., 2020; Cardoso and Marcuzzo, 2010), amplifying the susceptibility of the biome to fires (Libonati et al., 2020). Relying directly on fire data, most studies in the Pantanal have focused on a single discipline, for example, plant ecology (Libonati et al., 2020). Integrated knowledge about how fire distribution occurs over Pantanal different phytophysiognomies and historical comparisons between land use and land cover were scarce until this present study.

Fire ignition by natural causes (lightning strikes) typically occurs during the summer (from December to February) and reaches small extents. As such, fires occurring during the dry season are mainly human related and accounts for the major amount of burned area (Menezes et al., 2022). Acording to Damasceno-Junior et al. (2021), integrated fire management during ED season helps to reduce biomass in grasslands, which reduces large, high-intensity fires during the LD season (Eloy et al., 2018).

4.1. Limitations and uncertainties

Greater spatial resolution than 500 m, for example, will be fundamental for more localized, small-scale studies (Chuvieco et al., 2018; Rodrigues et al., 2019). Omission of small fires are, as well, a limitation of daily burned area products, as MCD64A (Campagnolo et al., 2021) and filtering spatial data with such a different spatial resolution as used here can also reduce accuracy in edge areas and small patches, typical for cropland class in Pantanal, for example. On the other hand, with the use of higher resolution imageries, such and Landsat and Sentinel, it may improve burned areas detection. In LULC classes formed by small patches this higher resolution mapping can indicate burned areas that are not considered in gross LULC and BA classifications, particularly when more than a decade of burns time series is being analyzed.

LULC maps were only available until 2019, therefore we were not able to analyze changes in land cover in more recent history. LULC maps in 2019

D.B. Correa et al.

Science of the Total Environment 835 (2022) 155386




Fig. 8. Bivariate map (burn frequency versus last burned year) by season. (a) LW are the BA detections by MCD64A1 during Late Wet season months (February, March, April), summing 60 months of 20 years. (b) ED are the detections during Early Dry season months (May, June, and July), summing 60 months of 20 years. (c) LD are the detections during Late Dry season months (August, September, October), summing 60 months of 20 years. (d) EW shows detections over Early Wet Season months (November, December and January), summing 63 months of 21 years.

was used as a surrogate for 2020 and 2021 in the analysis here performed, which may lead to some overestimation (or underestimation) of burned areas in land cover that have changed in the last two years.

${\bf 4.2.}\ Implications\ for\ management\ and\ future\ perspectives$

Despite the severity of 2020 fires in Pantanal, it has led to a better understanding of fire dynamics in the biome and allowed the identification of crucial structural gaps in local fire knowledge (Libonati et al., 2020). As adopted in this study, comparisons of fire occurrences across different

LULC, the availability of fuel loads, and across different fire seasons have revealed interannual trends of fire in Pantanal. This is crucial to predicting future fire frequencies, extensions, intensity, seasonality, and severity. It also aids fire managers in anticipating new trends and disasters.

More detailed research is needed on the weather conditions that fan fires, as well as the influences of ecology and management (Libonati et al., 2020). Information concerning fire in tropical wetlands are scant and the major fire studies are based on savannas and forests (Damasceno-Junior et al., 2021). In Brazil, most fire research has been developed in the Cerrado and Amazonia biomes (Pivello et al., 2021). Long and

D.B. Correa et al.

consistent datasets available for analysis are also crucial for environmental studies in general, including for the understanding of drivers of fire activity and its consequences (Chuvieco et al., 2019).

While Pantanal has been historically subjected to much environmental stress with its recurrent dry-wet episodes, fires are likely to worsen in the future. Given our future climate change scenarios, the environmental conditions (low precipitation, soil moisture, high air temperature) optimal for fire occurrences is likely to persist (Thielen et al. (2020), Marques et al., 2021). Thus, knowing the vulnerability of each LULC class in this sensitive biome will help local policymakers to plan and manage Pantanal to improve resilience.

5. Conclusion

Despite the significant remaining natural vegetation in Pantanal, we should not negate the pressures brought about by land conversion and degradation and the Pantanal's susceptibility to major disasters. Growth of pasture areas in lower land reflects intensive anthropogenic activity on the basin's higher catchment areas. Burnings in croplands have also been on the rise, particularly during the LD season, despite still constituting a relatively small area. On the other hand, humid areas and grassland areas are more susceptible to burnings during extreme dry weather conditions. For instance, grassland has been the main area burned in the Pantanal over the past twenty years. But in 2020 fire season, when comparing by proportion to the respective class area (RBA), forest, humid areas, and pasture became greatly affected by fires. Still, our study showed that the fires of 2020 was the most important event even though the biome has been continually burning in the past two decades. We conclude that when accounting BAs at regional level, we must understand and compare values from a stratified approach since burns can have different drivers and implications to fauna. flora and climate depending on LULC and periods of the year. Also, different rates of recovery and regeneration of the multiple land covers in Pantanal post-fire must be closely monitored, to better fire management strategies for long-term mitigation of the negative consequences on the conservation of the world's largest wetland.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgments

E.A. acknowledge the Brazilian National Council for Scientific and Technological Development (CNPq) for research grant 303169/2018-4. R.L. was supported by CNPq [grant 311487/2021-1], FAPERJ [grant E26/202.714/2019]; and partially funded by Project Rede Pantanal (FINEP grant 01.20.0201.00). E.P. acknowledge the Nanyang Technological University (#SUG-NAP 3/19EP) and the Ministry of Education of Singapore (#Tier1 RT06/19, #Tier1 2021-T1-001-056).

Appendix A. Supplementary data

Supplementary data to this article can be found online at https://doi. org/10.1016/j.scitotenv.2022.155386.

Alho, C.J.R., Sabino, J., 2011. A conservation agenda for the Pantanal's biodiversity. Braz. J. Biol. 71, 327–335. https://doi.org/10.1590/S1519-69842011000200012. Alves, J.M.R., Tomé, R., Caldeira, R.M.A., Miranda, P.M.A., 2021. Asymmetric Ocean response to atmospheric forcing in an island wake: a 35-year high-resolution study. Front. Mar. Sci. 8, 56. https://doi.org/10.3389/fmars.2021.624392.

Amaral e Silva, A., Braga, M.Q., Ferreira, J., Santos, V.J., Alves, S.C., Oliveira, J.C., Calijuri, M.L., 2020. Anthropic activities and the Legal Amazon: estimative of impacts on forest

and regional climate for 2030. Remote Sens. Appl.: Soc. Environ. 18 (100304). https://

Archibald. S., Lehmann, C.E.R., Gómez-Dans, J.L., Bradstock, R.A., 2013. Defining pyrome Archibadi, S., Dellinalii, C.E.R., Golinez-Hans, J.L., Bradastock, K.A., 2013. Dellining pyronies and global fire regimes. Proc. Natl. Acad. Sci. 110 (16), 6442 6447. https://doi.org/10.1073/pnas.1211466110.

Assine, M.L., 2005. River avulsions on the Taquari megafan, Pantanal wetland, Brazil. Geo-

Assine, M.L., 2005. River avuisions on the Taquin megarant, Paratinan wetuand, farizit. Geomorphology 70 (3-4), 357-371. https://doi.org/10.1016/j.geomorph.2005.02.013.

Assine, M.L., Merino, E.R., Pupim, F.N., Warren, L.V., Guerreiro, R.L., McGlue, M.M., 2015.
Geology and geomorphology of the Pantanal Basin. In: Bergier, I., Assine, M. (Eds.), Dynamics of the Pantanal Wetland in South America. The Handbook of Environmental
Chemistry. 37. Springer, Cham. https://doi.org/10.1007/698_2015_349.

Bergier, I., Assine, M.L., McGlue, M.M., Alho, C.J.R., Silva, A., Guerreiro, R.L., Carvalho, J.C.,

2018. Amazon rainforest modulation of water security in the pantanal wetland. Sci. Total Environ 619-620 1116-1125 https://doi.org/10.1016/j.scitotenv.2017.11.163

Environ. 619–620, 1116–1125. https://doi.org/10.1016/j.scitotenv.2017.1163.
Berlinck, C.N., Lima, L.H.A., Pereira, A.M.M., Carvallo Jr., E.A.R., Paula, R.C., Thomas, W.M.,
Morato, R.G., 2022. The pantanal is on fire and only a sustainable agenda can save the
largest wetland in the world. Braz. J. Biol. 82. https://doi.org/10.1590/1519-6984.

.,, racstock, R.A., 2020. Unprecedented burn area of Australian mega forest fires. Nat. Clim. Chang. 10, 171–172. https://doi.org/10.1038/s41558-020-0716-1. Boer, M.M., Resco de Dios, V., Bradstock, R.A., 2020. Unprecedented burn area of Australian

0/16-1.
wman, D.M.J.S., Kolden, C.A., Abatzoglou, J.T., Johnston, F.H., van der Werf, G.R., Flannigan, M., 2020. Vegetation fires in the anthropocene. Nat. Rev. Earth Environ. 1 (10), 500-515. https://doi.org/10.1038/s43017-020-0085-3.

Campagnolo, M.L., Libonati, R., Rodrigues, J.A., Pereira, J.M.C., 2021, A comprehensive characterization of MODIS daily burned area mapping accuracy across fire sizes in tropical sas. Remote Sens. Environ. 252 (112115). https://doi.org/10.1016/j.rse.2020.

rdoso, M., Marcuzzo, F., 2010. Mapeamento de três decênios da precipitação pluviométrica

total e sazonal do bioma pantanal. https://rigeo.cprm.gov.br/handle/doc/958. Chuvieco, E., Luzundia-Loiola, J., Pettinari, M.L., et al., 2018. Generation and analysis of a new global burned area product based on MODIS 250 m reflectance bands and thermal anomalies. Earth Syst. Sci. Data 10, 2015–2031. https://doi.org/10.5194/essd-10-

vieco, E., Mouillot, F., van der Werf, G.R., San Miguel, J., Tanase, M., Koutsias, N., Giglio, L., 2019. Historical background and current developments for mapping burned area from satellite earth observation. Remote Sens. Environ. 225, 45–64. https://doi.org/10.1016/ rse.2019.02.013

J.I.S. (2016). J.C., Bishop, C., Dalla Mora Esquerdo, J.C., Kastens, J.H., Brown, J.C., 2016. Dinâmica da agricultura na Bacia do Alto Paraguai, in Anais 6o Simpósio de Geotecnologias no Pantanal, Cuiah, M. T., 22 a 26 de outubro 2016 (Embrapa Informática Agropecuária/INPE), 623623–632.

Damasceno-Junior, G.A., Roque, F.O., Garcia, L.C., Ribeiro, D.B., Tomas, W.M., Scremin-Dias, E., Joseph G. B., A. (1997) A. (2014) A. (2014)

april_2021_wetland_science_practice. Durbin, J., Watson, G.S., 1950. Testing for serial correlation in least squares regression: I. Biometrika 37 (3/4) 409-428 htt c//doi.org/10.2307/2332391

Biometrika 37 (3/4), 409–428. https://doi.org/10.2307/2332391.

Durbin, J., Watson, G.S., 1951. Testing for serial correlation in least squares regression: II. Biometrika 38 (1/2), 159–177. https://doi.org/10.3207/2332325.

Durigan, G., Pilon, N.A., Abreu, R.C., Hoffmann, W.A., Martins, M., Fiorillo, B.F., Vasconcelos, H.L., 2020. No net loss of species diversity after prescribed fires in the Brazillian savanna. Front. For. Glob. Chang. 3, 13. https://doi.org/10.3389/ffgc.2020.00013.

Eloy, L., Bilbao, B.A., Mistry, J., Schmidt, I.B., 2020. From suppression to fire management: Advances and resistances to changes in fire fire policy in the savannas of Brazil and Venezuela. Geograph. J. 185 (1). https://doi.org/10.1111/geoj.12245.

Embrapa, 2020. Clima. https://www.npf.embrapa.br/pesquiss/efb/clima.htm.

Embrapa Solos, 2017. Soil map of Brazil world reference base for soil resource [dataset]. 1: 5.000.000. http://geoinfoc.nps.embrapa.br/pesrs/geonode/%3/soils.brazil.why.wgs84.

Emorapa Solos, 2017. Soli miap of Brazil words reterence base for soli resource (dataset). I: 5.000.000. http://geoinfo.cnps.embrapa.br/alyers/geoinde/%3Asolis_brazil_wrb_wgs84. Fernandes, F.A.F., Fernandes, A.H.B.M., Soares, M.T.S., Pellegrin, L.A., Lima, I.D., 2007. Atualização do mapa de solos da planície pantaneira para o sistema brasileiro de classificação de solos. Comunicado Técnico 61. 2007. Embrapa Pantanal, Corumbá . https://ainfo.cnptia.embrapa.br/digital/bitstream/item/161349/1/COT61.pdf. Fidelis, A., Muller, S.C., Pillar, V.D., Pfadenhauer, J., 2007. Efeito do Fogo na Ecologia de

Proteins, A., Muner, S.C., Pinar, V.D., Pradeninater, J., 2007. Election of Pogo ha Devologia de Populações de Herbácease e Arbustos dos Campos Sulinos. Nota Científica. Revista Brasileira de Biociências, Porto Alegre. 5(S1), pp. 303–305.

Franke, J., Barradas, A.C.S., Borges, M.A., Costa, M.M., Dias, P.A., Hoffmann, A.A., Siegert, F., 2018. Fuel load mapping in the brazilian cerrado in support of integrated fire management. Remote Sens. Environ. 217, 221–232. https://doi.org/10.1016/j.rse.2018.08.018.

Fuller, W.A., 1996. Introduction to Statistical Time Series. 2nd ed. Wiley ISBN: 978-0-471-

55239-U. Garcia, L.C., Szabo, J.K., Roque, F.O., Pereira, A.M., da Cunha, C.N., Damasceno-Junior, G.A., Morato, R.G., Tomas, W.M., Libonati, R., Ribeiro, D.B., 2021. Record-breaking wildfires in the world's largest continuous tropical wetland: integrative fire management is urgently needed for both biodiversity and humans. J. Environ. Manag. 293, 112870. https://doi.org/10.1016/j.jenvman.2021.112870.

nttps://doi.org/10.1016/j.jenvman.2021.112870.

Giglio, L., Justice, C., Boschetti, L., Roy, D., 2015. MCD64A1 MODIS/Terra + Aqua Burned Area Monthly 1.3 Global 500m SIN Grid V006 [dataset]. NASA EOSDIS Land Processes DAAC. https://doi.org/10.5067/MODIS/MCD64A1.006.

Gomes, A.C., Bernardo, N., Alcântara, E., 2017. Accessing the southeastern Brazil 2014 drought severity on the vegetation health by satellite image. Nat. Hazards https://doi.org/10.1007/s11069-017-3029-6.

Guerra, A., Roque, F.O., Garcia, L.C., Ochoa-Quintero, J.M., Sanches de Oliveira, P.T., Dettogni Guariento, R., Rosa, I.M.D., 2020. Drivers and projections of vegetation loss in

- the pantanal and surrounding ecosystems. Land Use Policy 91, 104388. https://doi.org/ 10.1016/i.landusenol.2019.104388
- Honilton, S.K., 2010. Biogeochemical implications of climate change for tropical rivers and floodplains. Hydrobiologia 657, 19–35. https://doi.org/10.1007/s10750-009-0086-1.
 Hardesty, J., Myers, R., Fulks, W., 2005. Fire, ecosystems, and people: a preliminary assess-
- ment of fire as a global conservation issue. No. 4The George Wright Forum. Vol. 22.
- George Wright Society, pp. 78–87. https://www.jstor.org/stable/43597968.
 Helsel, D.R., Hirsch, R.M., 2002. Statistical Methods in Water Resources Techniques of Water Resources Investigations, Book 4, A3. U.S. Geological Survey https://doi.org/10.3133/
- Hoffmann, W.A., Adasme, R., Haridasan, M., Carvalho, M., Geiger, E.L., Pereira, M.A.B., 2009. Free topkill, not mortality, governs the dynamics of alternate stable states at savannaforest boundaries under frequent fire incentral Brazil, Ecology 90, 1326–1337
- Holden, J. Moody, C.S. Edward Turner, T. et al., 2018, Water-level dynamics in natural and uen, J., moody, C.S., Edward i umer, I., et ai., 2018. water-level dynamics in natural and artificial pools in blanket peatlands. Hydrol. Process. 32, 550–561. https://doi.org/10.1002/hyp.11438.
 sain, M., Mahmud, I., 2019. pyMannKendall: a python package for non parametric Mann
- Kendall family of trend tests. J. Open Source Softw. 4 (39), 1556. https://doi.org/10. 21105/joss.01556.
- IBGE, 2012. Manual técnico da vegetação brasileira: sistema fitogeográfico: inventário das ts, 2012. Manual tecnico da vegetação orasmena: sistema inogeogranico. inversamo das formações florestais e campestres: técnicas e manejo de coleções botânicas: procedimentos para mapeamentos. Rio de JaneiroZnd ed. . https://biblioteca.ibge.gov.br/index.php/biblioteca-catalogo?view=detalhes&id=263011.
- br/index.php/biblioteca-catalogo?view=detalhes&id=263011. IBGE, 2019. Mapa de Biomas do Brasil [dataset]. 1:250.000. https://www.ibge.gov.br/ ciencias/informacoes-ambientais/vegetacao.html.
- Ivory, S.J., McGlue, M.M., Spera, S., Silva, A., Bergier, I., 2019. Vegetation, rainfall, and pulsing hydrology in the pantanal, the world's largest tropical wetland. Environ. Res. Lett. 14 (12), 124017.
 Junk, W.J., Cunha, N.C., 2005. Pantanal: a large south American wetland at a crossroads.
- Ecol. Eng. 24 (4), 391–401. https://doi.org/10.1016/j.ecoleng.2004.11.012 Junk, W.J., da Cunha, C.N., Wanteen, K.M., 2006. Biodiversity and its conservation in the Pantanal of Mato Grosso, Brazil. Aquat. Sci 68, 278–309. https://doi.org/10.1007/ s00027-006-0851-4
- s0002/-00-0851-4.

 Koppen, W., 1948. Climatologia: con un estudio de los climas de la tierra (No. QC861 K6).

 Fondo de Cultura Económica, México.

 Kumar, S., Getirana, A., Libonati, R., et al., 2022. Changes in land use enhance the sensitivity
- of tropical ecosystems to fire-climate extremes. Sci. Rep. 12, 964. https://doi.org/10. 1038/s41598-022-05130-0
- Lözaro, W.L., Oliveira-Jónior, E.S., Silva, C.J.D., Castrillon, S.K.I., Muniz, C.C., 2020. Climate change reflected in one of the largest wetland in the world: an overview of the northern pantanal water regime. Acta Limnol. Bras. 32 (104). https://doi.org/10.1590/s2179-75x7619.
- Li. S., Baneriee, T., 2021, Spatial and temporal pattern of wildfires in California from 2000 to
- Libonati, R., Sander, L.A., Peres, L.F., DaCamara, C.C., Garcia, L.C., 2020. Rescue Brazil's burning pantanal wetlands. Nature 588, 217–220. https://doi.org/10.1038/d41586-020-02066.
- Libonati, R., et al., 2022. Assessing the role of compound drought and heatwave events on unprecedented 2020 wildfires in the pantanal. Environ. Res. Lett. 17, 015005. https://doi. /10 1088/1748-9326/ac462e
- Long, T., Zhang, Z., He, G., Jiao, W., Tang, C., Wu, B., Zhang, X., Wang, G., Yin, R., 2019. 30 m resolution global annual burned area mapping based on landsat images and Google earth engine. Remote Sens. 11 (5), 489. https://doi.org/10.3390/rs11050489.
 Machado, R.X., da Costa, E.A., 2018. O turismo de Pesca em Corumbá, na Fronteira oeste do
- brasil. RITUR Rev. Iberoam. Turismo 8 (1), 36-48. brasil. RTIUR Rev. Iberoam. Turismo 8 (1), 36–48.
 Marengo, J.A., Cunha, A.P., Cuartas, L.A., Deusdará Leal, K.R., Broedel, E., Seluchi, M.E.,
 Michelin, C.M., De Praga Baião, C.F., Chuchón Ângulo, E., Almeida, E.K., Kazmierczak,
 M.L., Mateus, N.P.A., Silva, R.C., Bender, F., 2021. Extreme drought in the Brazilian
 pantanal in 2019–2020: characterization, causes, and impacts. Front. Water 3
 (639204), 13. https://doi.org/10.3389/frwa.2021.639204 vol 3.
- Marques, J.F., Alves, M.B., Silveira, C.F., Silva, A.A., Silva, T.A., Santos, V.J., Calijuri, M.J. Ques, Jar, Alves, M.B., Silvela, C.F., Silve, A.A., Silve, I.A., Sallios, V.J., Calljuir, M.J., 2021. Fires dynamics in the pantanal: impacts of anthropogenic activities and climate change. J. Environ. Manag. 299, 113586. https://doi.org/10.1016/j.jenvman.2021.113586.
- Martins, P.I., Belém, L.B.C., Szabo, J.K., Libonati, R., Garcia, L.C., 2022. Prioritising areas for wildfire prevention and post-fire restoration in the brazilian pantanal, Ecol. Eng. 176,
- 106617. https://doi.org/10.1016/j.ecoleng.2021.106517. lezes, L.S., Oliveira, A.M., Santos, F.L.M., Russo, A., Souza, R.A.F., Roque, F., Libonati, R. Menezes, L.S., Oliveira, A.M., Santos, F.L.M., Russo, A., Soluza, R.A.F., Roque, F., Libohadi, K., 2022. Lightning patterns in the pantanal: untangling natural and anthropogenic-induced wildfires. Sci. Total Environ. https://doi.org/10.1016/j.scitotenv.2022.153021.
 Miranda, H.S., 2010. Efeitos do regime de fogo sobre a estrutura de comunidades de Cerrado:
- Projeto Fogo. http://www.ibama.gov.br/sophia/cnia/livros/efeitosdoregimedofogodigital.
- Mota, P.H.S., Rocha, J.S.S., Castro, N.L.M., Marcatti, G.E., França, L.C.J., Schettini, B.L.S. Villanova, P.H., Santos, H.T., Santos, A.R., 2019. Forest fire hazard zoning in Mato Grosso state, Brazil. Land Use Policy 88, 104206. https://doi.org/10.1016/j.landusepol.2019.
- Mourão, G., Tomas, W., Campos, Z., 2010. How much can the number of jabiru stork (Ciconiidae). Nests vary due to change of flood extension in a large neotropical flood-plain? Zoologia 27, 751–756. https://doi.org/10.1590/S1984-46702010000500012.

- Nobre, C.A., 2010. Mudanças climáticas e o brasil-Contextualização. Parcerias Estratégicas 13 (27), 07-18. http://seer.cgee.org.br/index.php/parcerias estrategicas/artic ile/326/320
- Intel 320732D.
 Silva Junior, C.A., 2020. Fire foci related to rainfall and biomes of the state of Mato Grosso do Sul, Brazil. Agric. For. Meteorol. 282–283, 107861. https://doi.org/10.1016/j.agrformet.2019.107861 ISSN 0168-1923.
- Pausas, J.G., Parr, C.L., 2018. Towards an understanding of the evolutionary role of fire in an-
- Faths, J.G., Falt, C.L., 2018. 10was at understanding to the evolutionary for the first arimals. Evol. Ecol. 32 (2), 113–125. https://doi.org/10.1007/s10682-018-9927-6.
 Peel, M.C., Finlayson, B.L., McMahon, T.A., 2007. Updated world map of the Köppen-Geiger climate classification. Hydrol. Earth Syst. Sci. 11, 1633–1644. https://doi.org/10.5194/
- Pivello, V.R., 2011. The use of fire in the cerrado and amazonian rainforests of Brazil: past and
- present. Fire Ecol. 7 (1), 24–39, https://doi.org/10.4996/fireecology.0701024.

 Pivello, V.R., Vieira, I., Christianini, A.V., Ribeiro, D.B., et al., 2021. Understanding Brazil's catastrophic fires: causes, consequences and policy needed to prevent future tragedies.

 Perspect. Ecol. Conserv. https://doi.org/10.1016/j.pecon.2021.06.005.

 Pott, A., Pott, V.J., 2004. Features and conservation of the brazilian pantanal wetland. Wetl. Ecol. Manag. 12, 547–552. https://doi.org/10.1007/s11273-005-1754-1.

 Prestes, N.C.C.S., Massi, K.G., Silva, E.A., et al., 2020. Fire effects on understory Forest regen-
- eration in southern Amazonia. Front. For. Glob. Chang. 3, 10. https://doi.org/10.3389/ ffgc.2020.00010.
- Ingc. 2020.00010.
 Ingc. 2020.00010.
 Ingc. 2020.00011.
 Identify a products represent fire patterns in the Brazilian Savannas biome? An Accuracy Assessment of the MCD64 collections. Int. J. Appl. Earth Obs. Geoinf. 78, 318–331. https://doi.org/10. 1016/i.jag.2019.02.010
- Sabater, J.M., 2019, ERA5-Land monthly averaged data from 1981 to present [dataset], Co-
- Sandett, J.M., 2015. EAC-Land multily averaged untain from 1991 to present (tadaset). Co-penitics Climate Change Service (C3S) Climate Data Store (CDS) https://doi.org/10.24381/cds.68d2b530 Accessed 2021-07-20 from.
 Sansevero, J.B.B., Garbin, M.L., Sánchez-Tapia, A., Valladares, F., Scarano, F.R., 2020. Fire drives abandoned pastures to a savanna-like state in the Brazilian Atlantic Forest.
- Perspect. Ecol. Conserv. 18 (1), 31–36. https://doi.org/10.1016/j.pecon.2019.12.004. Sen, P.K., 1968. 63. J. Am. Stat. Assoc. (324), 1379–1389 https://doi.org/10.2307/2285891 ISSN 0168-1923
- ISSN 0108-1923. nabukuro, Y.E., Arai, E., Duarte, V., Dutra, A.C., Cassol, H.L.G., Sano, E.E., Hoffmann, T.B., 2020. Discriminating land use and land cover classes in Brazil based on the annual PROBA-V 100 m time series. IEEE J. Sel. Top. Appl. Earth Obs. Remote Sens. 13, 3409-3420. https://ieeexplore.ieee.org/stamp/stamp.jsp?arnumber=9103191.
- Somorowska, U., 2016. Changes in drought conditions in Poland over the past 60 years evaluated by the standardized precipitation-evapotranspiration index. Acta Geophys. 64, 2530–2549. https://doi.org/10.1515/acgeo-2016-0110.

 Soriano, B.M.A., Cardoso, E.L., Tomás, W.M., Santos, S.A., Crispim, S.M.A., Pellegrin, L.A., 2020.
- Uso do fogo para o manejo da vegetação no Pantanal; Corumbá: Embrapa Pantanal. 164. ISSN 1981-7223 https://ainfo.cnptia.embrapa.br/digital/bitstream/item/214625/1/Uso-
- logo-inancjo-2020-pl.i.

 Ag. C.M.Z., Shimbo, J., Rosa, M.R., Parente, L.L., Alencar, A., Rudorff, B.F., Azevedo, T.,

 2020. Reconstructing three decades of land use and land cover changes in Brazilian biomes with landsat archive and earth engine [dataset]. Remote Sens. 12 (17), 2735. https://doi.org/10.3390/rs12172735.
- envoorden, J., Meddens, A.J., Martinez, A.J., Foster, L.J., Kissling, W.D., 2019. The potential importance of unburned islands as refugia for the persistence of wildlife spe fire-prone ecosystems. Ecol. Evol. 9 (15), 8800–8812. https://doi.org/10.1002/ece3
- Theil, H., 1992. A rank-invariant method of linear and polynomial regression analysis. In: Raj, B., Koerts, J. (Eds.), Henri Theil's Contributions to Economics and Econometrics. Ad
- B., Noerts, J. (Eds.), Henri Theil's Contributions to Economics and Econometrics. Advanced Studies in Theoretical and Applied Econometrics. vol 23. Springer, Dordrecht. https://doi.org/10.1007/978-94-011-2546-8_20.

 Pelen, D., Schuchmann, K.L., Ramoni-Perazzi, P., Marquez, M., Rojas, W., Quintero, J.I., Marquez, M., 2020. Quo vadis Pantanal? Expected precipitation extremes and drought dynamics from changing sea surface temperature. Plos one 15 (1), e0227437. https:// doi.org/10.1371/journal.pone.0227437.
- Thielen, D., Ramoni-Perazzi, P., Puche, M.L., Márquez, M., Quintero, J.L., Rojas, W., Soto-Thielen, D., Ramoni-Perazzi, P., Puche, M.L., Marquez, M., Quintero, J.I., Rojas, W., Soto-Werschitz, A., Thielen, K., Nunes, A., Libonati, R., 2021. The pantanal under Siege—on the origin, dynamics and forecast of the megadrought severely affecting the largest wetland in the world. Water 13 (21), 3034. https://doi.org/10.3390/w13213034.
 Tomas, W.M., Berlinck, C.N., Chiaravalloti, R.M., et al., 2022. Distance sampling surveys reveal 17 million vertebrates directly killed by the 2020's wildfires in the Pantanal, Brazil. Sci Rep 11, 23547. https://doi.org/10.1038/s41598-021-02844-5.
- UNEP, 2022. Spreading like wildfire: the rising threat of extraordinary landscape fires. Report. https://www.unep.org/resources/report/spreading-wildfire-rising-threat-extraordinary-landscape-fires.
- port. https://www.inep.org/resources/report/spreading-withine-rising-unear-extraordinary-landscape-fires. Uyanto, S.S., 2020. Power comparisons of five Most commonly used autocorrelation tests. Pak. J. Stat. Oper. Res. 16 (1), 119–130. https://doi.org/10.18187/pjsor.v16i1.2691. Witze, A., 2020. The Arctic is burning like never before - and that's bad news for climate
- change. Nature 585, 336–337. https://doi.org/10.1038/d41586-020-02568-y. Yue, S., Wang, C., 2004. The mann-kendall test modified by effective sample size to detect trend in serially correlated hydrological series. Water Resour. Manag. 18, 201–218. https://doi.org/10.1023/B:WARM.0000043140.61082.60.

3 FINAL CONSIDERATIONS

Despite the significant remaining natural vegetation in Pantanal (more than 80%), it does not mean that there are no relevant pressures for land conversion and degradation and susceptibility for major disasters as 2020 wildfires in the biome. Growth of pasture areas, reflecting an intensive anthropogenic activity on basin higher catchment areas followed by cropland settlement (GUERRA et al., 2020) could be also seen in our results in lower lands. On the other hand, humid areas and grassland yearly areas are more susceptible to weather conditions of dry/ warmer versus humid periods as also discussed by Marengo et al. (2021) and shrubland area decreasing.

The months with highest values of ABA for Pantanal were found for August and September when lowest rainfall mean occurs and mean river levels starts to decline on southwest portion. But, accordingly to Menezes et al. (2022), lightening dischargers, natural fire ignitors, are concentrated from December to February. During dry season, ignitors are considered human related and agricultural practices when cattle ranchers use fire to renew grazing are applied.

Beside 2020, the years of 2005 and 2001 also burned major areas in grassland class considering specific months. In addition, with 2020, 2002 and 2019 had also prolonged burning seasons in Pantanal, from July to November depending on the year, all of them reaching more than 19000 km² of ABA. Proportionally to the respective class area (RBA), forest, humid areas, and pasture appear in contrast to other classes in 2020 burns, being highly affected while in 2005 the RBA highlight is for grassland, pasture and savannah. Relatively, forests, grasslands and croplands were classes with higher burned areas especially during dry season months (August and September) and pastures, with lower registers than the others LULC. In lower scale we saw that since 2014, cropland burnings are growing exponentially during LD season.

These differences reinforce that when accounting BAs at regional level, we must understand and compare values from multiple perspectives, especially when burns can have different drivers and implications to fauna, flora and climate depending on LULC and periods of the year. A tropical forest burn result in more GHG emissions and vegetation recovery time than grassland burns, for example while for grasslands and savannas, burning is an important ecological factor (HARDESTY et al. 2005). So, fires can be detrimental for some parts of Pantanal (HOFFMANN et al., 2009)

Most of Pantanal was burnt at least once in the last twenty years. Across the 243 analyzed months, the maximum burn frequency reached 19 times that of fire recurrence in the same pixel, specially over current (2019 map) grassland class, in south Pantanal. Smaller burned areas were frequent for all LULC classes. As for larger burned areas (more than 8.000 km²), they were observed in forest and grassland at a lower frequency. These results suggest that burned spots with small areas in Pantanal is consistent for the last 20 years.

But still, our results of burned areas for the last 20 years across different LULC in Pantanal, showed that the 2020 fires were the most important event even though the biome has been continuously burning along the past two decades. This is likely due to severe droughts in the region caused by a decrease in precipitation and river levels (MARENGO et al. 2021, LÁZARO et al. 2020, OLIVEIRA-JÚNIOR et al. 2020). Extreme droughts in 2020 and 2005 for all LULC classes coincided with great fire events reported in other studies (CARDOSO AND MARCUZZO, 2010; MARENGO et al. 2021). Coupled with 2019 and 2020's climate and biomass conditions, wildfires in Pantanal reached huge proportions.

2020's fire season was exceptional, but conditions that led to these blazes, such as increase of ignition sources, biomass availability, drier and warmer atmosphere, are becoming increasingly common (LIBONATI et al., 2020). On the Pantanal portion of Mato Grosso do Sul State, Oliveira-Junior (2020) had already identified a growth trend of fire foci with fire dataset ranging from 2000 to 2015.

We hypothesized a growing trend in burned areas in all LULC in Pantanal, especially in grasslands, savannas and pastures. We have found an increase in burned areas, but in cropland land use class. In general, cropland expansion does not imply in burnings once it settles in older pasture areas, an already converted LULC class from natural to anthropic stage (CHUVIECO et al., 2021). The land use changes involved with deforestation of native vegetation traditionally involve fire, as when for opening new pasture areas (ARCHIBALD et al., 2013). deforestation and the increase in large agribusiness properties in Pantanal could be linked to these recent fire events (MOTA et al., 2019; MARQUES et al., 2021). The higher BAs proportions found on native vegetation compared to anthropic managed areas also highlight a possible of lack of fire control and adequate management.

According to Damasceno-Junior et al. (2021), fire helps to reduce biomass in grasslands. This represents high fire frequency but less intensity in each event (ELOY et al. 2018). Nevertheless, it is important to note that the intensity of the LD season fires is higher as they release more heat and have higher flames and combustor factor (DOS SANTOS et al.

2021). Fires are thus, more destructive in the LD season than early dry season. As a result, fire ignited in the grassland can spread to the savannas and forest due open areas, wind and dry vegetation. Almost 17% of Pantanal has been deforested using fire, mainly the savanna areas (ALHO, 2008).

Relying directly on fire data, most studies in the Pantanal have focused on a single discipline, for example, plant ecology (LIBONATI et al., 2020). Integrated knowledge about how fire distribution over Pantanal different phytophysiognomies and historical comparisons between land use and land cover were scarce until this present study. Different global BA products complement each other (CHUVIECO et al. 2018, LONG et al. 2019) once all of them has its limitations. So, we encourage the investigation with field campaigns for a better understanding of these edges especially for Pantanal and comparations between BA products with their multiple parameters, as fire intensity and burn polygon.

REFERENCES

- ALHO, C. J. R.; SABINO, J. A conservation agenda for the Pantanal's biodiversity. **Brazilian Journal of Biology**, v. 71, p. 327-335, 2011. Available on: https://doi.org/10.1590/S1519-69842011000200012 Acesso em: 14 ago. 2022.
- ARCHIBALD, S.; LEHMANN, C. E. R.; GÓMEZ-DANS, J. L.; BRADSTOCK, R. A. Defining pyromes and global syndromes of fire regimes. **Proceedings of the National Academy of Sciences**, v. 110, n. 16, p. 6442-6447, 2013. DOI: 10.1073/pnas.1211466110
- BOWMAN, D. M. J. S.; KOLDEN, C. A.; ABATZOGLOU, J. T.; JOHNSTON, F. H.; VAN DER WERF, G. R.; FLANNIGAN, M. Vegetation fires in the Anthropocene. **Nature Reviews Earth & Environment**, v. 1, n. 10, p. 500-515, 2020. Available on: https://doi.org/10.1038/s43017-020-0085-3 . Acesso em: 14 ago. 2022.
- CARDOSO, M. R. D.; MARCUZZO, F. F. N. Mapeamento de três decênios da precipitação pluviométrica total e sazonal do bioma Pantanal. 2010. Available on: https://rigeo.cprm.gov.br/handle/doc/958 Acesso em: 14 ago. 2022.
- CHUVIECO, E.; LUZUNDIA-LOIOLA, J.; PETTINARI, M. L. et al. Generation and analysis of a new global burned area product based on MODIS 250 m reflectance bands and thermal anomalies. **Earth System Science Data**, v. 10, n. 4, p. 2015-2031, 2018. Available on: https://doi.org/10.5194/essd-10-2015-2018 Acesso em: 14 ago. 2022.
- DAMASCENO-JUNIOR, G. A.; ROQUE, F. O.; GARCIA, L. C., RIBEIRO, D. B.; TOMAS, W. M., SCREMIN-DIAS, E. et al. Lessons to be Learned from the Wildfire Catastrophe of 2020 in the Pantanal Wetland. **Wetland Science & Practice**, v. 38, p. 107-15, 2021. Available on: https://issuu.com/societyofwetlandscientists/docs/april_2021_wetland_science___practice Acesso em: 14 ago. 2022.
- DURIGAN, G.; PILON, N. A.; ABREU, R. C.; HOFFMANN, W. A.; MARTINS, M.; FIORILLO, B. F.; VASCONCELOS, H. L. No net loss of species diversity after prescribed fires in the Brazilian savanna. **Frontiers in Forests and Global Change**, v. 3, p. 13, 2020. DOI: 10.3389/ffgc.2020.00013
- FIDELIS, A.; MULLER, S. C.; PILLAR, V. D.; PFADENHAUER, J. Efeito do fogo na ecologia de populações de herbáceas e arbustos dos campos sulinos. **Revista Brasileira de Biociências**, v. 5, n. S1, p. 303-305, 2007.
- GARCIA, L. C.; SZABO, J. K.; ROQUE, F. O.; PEREIRA, A. M.; DA CUNHA, C. N.; DAMASCENO-JUNIOR, G. A et al. Record-breaking wildfires in the world's largest continuous tropical wetland: integrative fire management is urgently needed for both biodiversity and humans. **Journal of environmental management**, v. 293, p. 112870, 2021. Available on: https://doi.org/10.1016/j.jenvman.2021.112870 Acesso em: 14 ago. 2022.
- GUERRA, A.; ROQUE, F. O.; GARCIA, L. C.; OCHOA-QUINTERO, J. M.; SANCHES DE OLIVEIRA, P. T.; DETTOGNI GUARIENTO, R.; ROSA, I. M. D. Drivers and projections of vegetation loss in the Pantanal and surrounding ecosystems. **Land Use Policy**, v. 91, p. 104388, 2020. Available on: https://doi.org/10.1016/j.landusepol.2019.104388. Acesso em: 14 ago. 2022.
- HARDESTY, J.; MYERS, R.; FULKS, W. Fire, ecosystems, and people: a preliminary assessment of fire as a global conservation issue. In: **The George Wright Forum**. George Wright Society, v. 22, n. 4, p. 78-87, 2005. Available on: https://www.istor.org/stable/43597968. Acesso em: 14 ago. 2022.
- HOFFMANN, W. A.; ADASME, R.; HARIDASAN, M.; CARVALHO, M., GEIGER, E. L., et al. Tree topkill, not mortality, governs the dynamics of savanna–forest boundaries under frequent fire in central Brazil. **Ecology**, v. 90, n. 5, p. 1326-1337, 2009.
- JUNK, W. J.; DA CUNHA, C. N.; WANTZEN, K. M et al. Biodiversity and its conservation in the Pantanal of Mato Grosso, Brazil. **Aquatic Sciences**, v. 68, n. 3, p. 278-309, 2006. Available on: https://doi.org/10.1007/s00027-006-0851-4 Acesso em: 14 ago. 2022.

- KARNAUSKAS, K. B.; LUNDQUIST, J. K.; ZHANG, L. Southward shift of the global wind energy resource under high carbon dioxide emissions. **Nature Geoscience**, v. 11, n. 1, p. 38-43, 2018. Available on: https://doi.org/10.1038/s41561-017-0029-9 Acesso em: 14 ago. 2022.
- LÁZARO, W. L.; OLIVEIRA-JÚNIOR, E. S.; SILVA, C. J. D.; CASTRILLON, S. K. I.; MUNIZ, C. C. Climate change reflected in one of the largest wetlands in the world: an overview of the Northern Pantanal water regime. **Acta Limnologica Brasiliensia**, v. 32, 2020. DOI: 10.1590/s2179-975x7619
- LIBONATI, R.; SANDER, L. A.; PERES, L. F.; DACAMARA, C. C.; GARCIA, L. C. Rescue Brazil's burning Pantanal wetlands. **Nature**, v. 588, p. 217–220. 2020. Available on: https://doi.org/10.1038/d41586-020-03464-1. Acesso em: 14 ago. 2022.
- LIBONATI, R. et al. Twenty-first century droughts have not increasingly exacerbated fire season severity in the Brazilian Amazon. **Scientific reports**, v. 11, n. 1, p. 1-13, 2021. Available on: https://doi.org/10.1038/s41598-021-82158-8 Acesso em: 14 ago. 2022.
- LONG, T. et al. 30 m resolution global annual burned area mapping based on Landsat Images and Google Earth Engine. **Remote Sensing**, v. 11, n. 5, p. 489, 2019. Available on: https://doi.org/10.3390/rs11050489 Acesso em: 14 ago. 2022.
- MARENGO, J. A. et al. Extreme drought in the Brazilian Pantanal in 2019–2020: characterization, causes, and impacts. **Frontiers in Water**, v. 3, p. 639204, 2021. DOI: 10.3389/frwa.2021.639204 v.3
- MARQUES, J. F. et al. Fires dynamics in the Pantanal: Impacts of anthropogenic activities and climate change. **Journal of Environmental Management**, v. 299, p. 113586, 2021. Available on: https://doi.org/10.1016/j.jenvman.2021.113586. Acesso em: 14 ago. 2022.
- MENEZES, L. S. et al. Lightning patterns in the Pantanal: Untangling natural and anthropogenic-induced wildfires. **Science of The Total Environment**, v. 820, p. 153021, 2022. Available on: http://dx.doi.org/10.1016/j.scitotenv.2022.153021 Acesso em: 14 ago. 2022.
- MOTA, P. H. S. et al. Forest fire hazard zoning in Mato Grosso state, Brazil. Land use policy, v. 88, p. 104206, 2019. DOI: 10.1016/j.landusepol.2019.104206
- OLIVEIRA, G. et al. Smoke pollution's impacts in Amazonia. **Science**, v. 369, n. 6504, p. 634-635, 2020. Available on: https://doi.org/10.1126/science.abd5942. Acesso em: 14 ago. 2022.
- OLIVEIRA-JUNIOR, J. F. et al. Fire foci related to rainfall and biomes of the state of Mato Grosso do Sul, Brazil. **Agricultural and Forest Meteorology**, v. 282, p. 107861, 2020. Available on: https://doi.org/10.1016/j.agrformet.2019.107861. Acesso em: 14 ago. 2022.
- PAUSAS, J. G.; PARR, C. L. Towards an understanding of the evolutionary role of fire in animals. **Evolutionary Ecology**, v. 32, n. 2, p. 113-125, 2018. Available on: https://doi.org/10.1007/s10682-018-9927-6 Acesso em: 14 ago. 2022.
- PEREIRA, P. et al. Post-fire soil management. **Current Opinion in Environmental Science & Health**, v. 5, p. 26-32, 2018. Available on: https://doi.org/10.1016/j.coesh.2018.04.002 Acesso em: 14 ago. 2022.
- PIVELLO, V. R.; VIEIRA, I.; CHRISTIANINI, A. V.; RIBEIRO, D. B. Understanding Brazil's catastrophic fires: Causes, consequences and policy needed to prevent future tragedies. **Perspectives in Ecology and Conservation**, v. 19, n. 3, p. 233-255, 2021. Available on: https://doi.org/10.1016/j.pecon.2021.06.005 Acesso em: 14 ago. 2022.
- SOMMERS, W. T.; COLOF, S. G.; CONRAD, S. G. Fire History and Climate Change. Report Submitted to the Joint Fire Science Program for Project 01-2-01-09. 2011. Cap. 3. Available on: https://www.firescience.gov/JFSP_fire_history.cfm Acesso em: 14 ago. 2022.

STEENVOORDEN, J. et al. The potential importance of unburned islands as refugia for the persistence of wildlife species in fire-prone ecosystems. **Ecology and evolution**, v. 9, n. 15, p. 8800-8812, 2019. Available on: https://doi.org/10.1002/ece3.5432 Acesso em: 14 ago. 2022.

THIELEN, D. et al. Quo vadis Pantanal? Expected precipitation extremes and drought dynamics from changing sea surface temperature. **Plos one**, v. 15, n. 1, p. e0227437, 2020. Available on: https://doi.org/10.1371/journal.pone.0227437 . Acesso em: 14 ago. 2022.

Tomas, W. M., Berlinck, C. N., Chiaravalloti, R. M. et al. Distance sampling surveys reveal 17 million vertebrates directly killed by the 2020's wildfires in the Pantanal, Brazil. **Scientific reports**, v. 11, n. 1, p. 1-8, 2021. Available on: https://doi.org/10.1038/s41598-021-02844-5 Acesso em: 14 ago. 2022.