

Nota Científica

Utilization of shells of the snail *Achatina fulica* Bowdich, 1822 (Mollusca, Gastropoda) by the hermit crab *Clibanarius vittatus* (Bosc, 1802) (Decapoda, Anomura) in the São Vicente Estuary, São Paulo, Brazil

**Bruno S. Sant'Anna, Cilene M. Zangrande & Alvaro L.D. Reigada
Research Group in Crustacean Biology (Crusta), Campus do Litoral Paulista
Universidade Estadual Paulista, Praça Infante D. Henrique s/n
11330-900 São Vicente (SP), Brazil**

ABSTRACT. Hermit crabs depend on mollusc shells for housing. In this study, an unusual resource is reported for a hermit crab that usually inhabits marine gastropod shells. During a field study conducted from May 2001 to April 2003 in an estuarine area in São Vicente, state of São Paulo, Brazil, 21 individuals of *Clibanarius vittatus* (Bosc, 1802) were found inhabiting the shells of the terrestrial gastropod *Achatina fulica* Bowdich, 1822. *A. fulica* occurs in nearshore grass patches, where occasional contact with sea water kills them, and their shells then become available to the hermit crabs.

Key words: *Clibanarius*, Gastropoda, Anomura, hermit crab, Brazil.

Utilización de conchas de caracoles terrestres *Achatina fulica* Bowdich, 1822 (Mollusca, Gastropoda) por el cangrejo ermitaño *Clibanarius vittatus* (Bosc, 1802) (Decapoda, Anomura) en el estuario de São Vicente, São Paulo, Brasil

RESUMEN. El cangrejo ermitaño depende de conchas de moluscos para hacer de ella su casa. Este estudio ha descubierto una manera inusual de como vive este cangrejo, que generalmente habita la concha de gastrópodos marinos. El estudio fue realizado en la estación marítima de São Vicente, provincia de São Paulo, de mayo de 2001 hasta abril de 2003. Durante ese periodo, un total de 21 cangrejos *Clibanarius vittatus* (Bosc, 1802) fueron encontrados viviendo en las conchas del caracol terrestre *Achatina fulica* Bowdich, 1822. *A. fulica* ocurre en áreas de césped cercanos a la costa, donde el contacto ocasional con el agua de mar provoca su muerte y sus conchas quedan disponibles para el cangrejo ermitaño.

Palabras clave: *Clibanarius*, Gastropoda, Anomura, cangrejo ermitaño, Brasil.

The occupation pattern of gastropod shells by hermit crabs is complex (Fernandes-Góes, 1997), but accessibility to empty shells is considered a key factor (Negreiros-Franozo *et al.*, 1991; Pinheiro *et al.*, 1993; Turra & Leite, 2001). Shell availability may be provided by a high mortality of gastropods residing nearby (Fernandes-Góes, 2000).

Achatina fulica Bowdich, 1822 is a pulmonate terrestrial gastropod, first introduced into Brazil for farming (Teles *et al.*, 1997), but later found breeding in the wild. In other countries, the introduction of *A. fulica* has led to the devastation of several crops (Vasconcellos & Pile, 2001). Besides its effect as a plague, *A. fulica* is also the intermediate host of

Angiostrongylus cantonensis, (Chen, 1935), a nematode that causes meningitis. In this study, the utilization of shells of *A. fulica* by the hermit crab *Clibanarius vittatus* (Bosc, 1822) is reported.

Hermit crabs were collected by hand, from May 2001 to April 2003, in an estuarine region in São Vicente (23°58'21"S; 46°23'35"W), São Paulo State, Brazil, during low tide when the crabs are exposed on the sediment. All hermit crabs and their shells were identified according to Melo (1999) and Rios (1994), and sexed from the gonopore position: in males the gonopores are open in coxae of the fifth pereopods, and in females in the third pereopods. The intersex has both male and female gonopores open, as observed by Turra (2004). When necessary, gastropod shells were sent to specialists for identification.

The 2,344 hermit crabs captured during the study period were using 13 gastropod species. Of this total, 21 used shells of *A. fulica*: these were 19 males and 2 intersexes, ranging from 7.0 to 13.1 mm shield length. We note that shells of *A. fulica* were the largest resource of shell utilized for *C. vittatus* (41.5 to 88.3 mm shield length) in this environment.

C. vittatus has been recorded using different shell types of marine gastropods, such as *Melongena corona* (Gmelin, 1791) (Lowery & Nelson, 1988) and *Thais (Stramonita) haemastoma* (Linnaeus, 1767) (Negreiros-Fransozo *et al.*, 1991; Reigada & Santos, 1997).

Sexual dimorphism in hermit crabs is very common. In *C. vittatus*, males reach a larger size than females (Reigada & Santos, 1997; Turra & Leite, 2000). Size-related sexual dimorphism is also observed in other hermit species, such as *Paguristes tortugae* Stimpson, 1759 (Negreiros-Fransozo & Fransozo, 1992) and *Clibanarius antillensis* Stimpson, 1759 (Turra & Leite, 1999). In most crustaceans, this difference in size is associated with the reproductive functions, where females have a larger energetic cost in the production of eggs and in the care of the young, whereas males direct their energy toward growth.

Large females of the terrestrial hermit crabs *Coenobita cavipes* Stimpson, 1858 and *Coenobita purpureus* Stimpson, 1858 were also found occupying shells of *A. fulica* in Okinawa, Japan (Nakasone, 2001). The author suggested that energetic expenditure in adult individuals using these large light shells would be low compared to

conspecifics housed in marine shells. However, according to Meireles *et al.* (2003), the competition among hermit crabs in the area as well as the few resources for occupation available to larger specimens of *Dardanus insignis* (Saussure, 1858) may be other stressing elements.

In the present study, all shells of *A. fulica* were used by males and a few intersexes, probably because they reach a larger size. By using these shells, energetic expenditure for locomotion would decrease and resources may be directed for intra and interspecific competition. The use of shells of *A. fulica* is due to its presence in grass patches along the shore, where contact with sea water kills them, making their shells available for *C. vittatus*.

ACKNOWLEDGEMENTS

For laboratory support during the development of this work, we thank the Universidade Estadual de São Paulo (Unesp), Campus do Litoral Paulista. We are also grateful to geologist Paulo Mattioli and to biologist Carlo Magenta Cunha for identifying this species of gastropod, to Dr. Augusto A.V. Flores for translating the manuscript into English, and to Andressa Hattori for translating the Abstract into Spanish. Dr. Janet W. Reid checked the english text of the final version. We also thank the referee, for valuable recommendations.

REFERENCES

- Fernandes-Góes, L.C. 1997.** Distribuição e biologia populacional de *Dardanus insignis* (Saussure, 1858) (Crustacea: Decapoda: Anomura) na região de Ubatuba, SP. Dissertação de Mestrado, Universidade Estadual Paulista Campus de Botucatu, São Paulo, Brasil, 150 pp.
- Fernandes-Góes, L.C. 2000.** Diversidade e bioecologia das comunidades de anomuros (Crustacea, Decapoda) do substrato não consolidado da região de Ubatuba, SP. Tese de Doutorado, Universidade Estadual Paulista, Campus de Botucatu, São Paulo, Brasil, 133 pp.
- Lowery, W.A. & W.G. Nelson. 1988.** Population ecology of the hermit crab *Clibanarius vittatus* (Decapoda, Diogenidae) at Sebastian Inlet, Florida. *J. Crust. Biol.*, 8(4): 548-556.

- Meireles, A.L., R. Biagi & F.L.M. Mantelatto. 2003.** Hermit crabs in evidence: usual gastropod shell occupation. *Nauplius*, 11(1): 63-66.
- Mello, G.A.S. 1999.** Manual de identificação dos Crustacea Decapoda do litoral brasileiro: Anomura, Thalassinidea, Palinuridea, Astacidea. Editora Plêide, São Paulo, 551 pp.
- Nakasone, Y. 2001.** Reproductive biology of three land hermit crabs (Decapoda: Anomura: Coenobitidae) in Okinawa, Japan. *Pac. Sci.*, 55(2): 157-169.
- Negreiros-Fransozo, M.L., A. Fransozo & N.J. Hebling. 1991.** Estrutura populacional e determinação do tamanho da concha em 4 espécies de ermitões (Crustacea, Decapoda, Anomura) do litoral paulista. *Biotemas*, 4(2): 135-148.
- Negreiros-Fransozo, M.L. & A. Fransozo. 1992.** Estrutura populacional e relação com a concha em *Paguristes tortugae* Schimitt, 1933 (Crustacea, Diogenidae), no litoral norte do estado de São Paulo, Brasil. *Naturalia*, 17: 31-42.
- Pinheiro, M.A.A., A. Fransozo & M.L. Negreiros-Fransozo. 1993.** Seleção e relação com a concha em *Isocheles sawayai* (Forest & Saint Laurent) 1967 (Crustacea, Anomura, Diogenidae). *Arq. Biol. Technol.*, 36(4): 745-752.
- Reigada, A.L.D. & S. Santos. 1997.** Biologia e relação com a concha em *Clibanarius vittatus* (Bosc, 1802) (Crustacea, Diogenidae) em São Vicente, S.P., Brasil. *Arq. Biol. Technol.*, 40(4): 941-952.
- Rios, E.C. 1994.** Seashells of Brazil. Instituto Acqua, Museu Oceanográfico do Rio Grande e Universidade do Rio Grande, 368 pp.
- Teles, H.M.S., J.F Vaz, L.R Fontes & M.F. Domingos. 1997.** Registro de *Achatina fulica* Bowdich, 1822 (Mollusca, Gastropoda) no Brasil: caramujo hospedeiro intermediário da angiostrongilíase. *Rev. Saúde Pública*, 31(3): 310-12.
- Turra, A. 2004.** Intersexuality in hermit crabs: reproductive role and fate of gonopores in intersex individuals. *J. Mar. Biol. Ass. U.K.*, 84: 754-759.
- Turra, A. & F.P.P. Leite. 1999.** Population structure and fecundity of the hermit crab *Clibanarius antillensis* Stimpson 1862 (Anomura, Diogenidae) in southeastern Brazil. *Bull. Mar. Sci.*, 64(2): 281-289.
- Turra, A. & F.P.P. Leite. 2000.** Population biology and growth of three sympatric species of intertidal hermit crabs in south-eastern Brazil. *J. Mar. Biol. Ass. U.K.*, 80: 1061-1069.
- Turra, A. & F.P.P. Leite. 2001.** Shell utilization patterns of a tropical rocky intertidal hermit crab assemblage: I. The case of Grande Beach. *J. Crust. Biol.*, 21(2): 393-406.
- Vasconcellos, M.C. & E. Pile. 2001.** Ocorrência de *Achatina fulica* no Vale do Paraíba, Estado do Rio de Janeiro, Brasil. *Rev. Saúde Pública*, 35(6): 582-584.

Recibido: 22 febrero 2005; Aceptado: 5 octubre 2005