

Measurement of the Semileptonic Branching Ratio of B_s^0 to an Orbitally Excited D_s^{**} State: $\text{Br}(B_s^0 \rightarrow D_{s1}^- (2536) \mu^+ \nu X)$

V. M. Abazov,³⁶ B. Abbott,⁷⁶ M. Abolins,⁶⁶ B. S. Acharya,²⁹ M. Adams,⁵² T. Adams,⁵⁰ E. Aguilo,⁶ S. H. Ahn,³¹ M. Ahsan,⁶⁰ G. D. Alexeev,³⁶ G. Alkhazov,⁴⁰ A. Alton,⁶⁵ G. Alverson,⁶⁴ G. A. Alves,² M. Anastasoiaie,³⁵ L. S. Ancu,³⁵ T. Andeen,⁵⁴ S. Anderson,⁴⁶ B. Andrieu,¹⁷ M. S. Anzels,⁵⁴ Y. Arnaud,¹⁴ M. Arov,⁶¹ M. Arthaud,¹⁸ A. Askew,⁵⁰ B. Åsman,⁴¹ A. C. S. Assis Jesus,³ O. Atramentov,⁵⁰ C. Autermann,²¹ C. Ay,²⁴ F. Badaud,¹³ A. Baden,⁶² L. Bagby,⁵³ B. Baldin,⁵¹ D. V. Bandurin,⁶⁰ S. Banerjee,²⁹ P. Banerjee,²⁹ E. Barberis,⁶⁴ A.-F. Barfuss,¹⁵ P. Bargassa,⁸¹ P. Baringer,⁵⁹ J. Barreto,² J. F. Bartlett,⁵¹ U. Bassler,¹⁸ D. Bauer,⁴⁴ S. Beale,⁶ A. Bean,⁵⁹ M. Begalli,³ M. Begel,⁷² C. Belanger-Champagne,⁴¹ L. Bellantoni,⁵¹ A. Bellavance,⁵¹ J. A. Benitez,⁶⁶ S. B. Beri,²⁷ G. Bernardi,¹⁷ R. Bernhard,²³ I. Bertram,⁴³ M. Besançon,¹⁸ R. Beuselinck,⁴⁴ V. A. Bezzubov,³⁹ P. C. Bhat,⁵¹ V. Bhatnagar,²⁷ C. Biscarat,²⁰ G. Blazey,⁵³ F. Blekman,⁴⁴ S. Blessing,⁵⁰ D. Bloch,¹⁹ K. Bloom,⁶⁸ A. Boehnlein,⁵¹ D. Boline,⁶³ T. A. Bolton,⁶⁰ G. Borissov,⁴³ T. Bose,⁷⁸ A. Brandt,⁷⁹ R. Brock,⁶⁶ G. Brooijmans,⁷¹ A. Bross,⁵¹ D. Brown,⁸² N. J. Buchanan,⁵⁰ D. Buchholz,⁵⁴ M. Buehler,⁸² V. Buescher,²² S. Bunichev,³⁸ S. Burdin,⁴³ S. Burke,⁴⁶ T. H. Burnett,⁸³ C. P. Buszello,⁴⁴ J. M. Butler,⁶³ P. Calfayan,²⁵ S. Calvet,¹⁶ J. Cammin,⁷² W. Carvalho,³ B. C. K. Casey,⁵¹ N. M. Cason,⁵⁶ H. Castilla-Valdez,³³ S. Chakrabarti,¹⁸ D. Chakraborty,⁵³ K. M. Chan,⁵⁶ K. Chan,⁶ A. Chandra,⁴⁹ F. Charles,^{19,*} E. Cheu,⁴⁶ F. Chevallier,¹⁴ D. K. Cho,⁶³ S. Choi,³² B. Choudhary,²⁸ L. Christofek,⁷⁸ T. Christoudias,⁴⁴ S. Cihangir,⁵¹ D. Claes,⁶⁸ Y. Coadou,⁶ M. Cooke,⁸¹ W. E. Cooper,⁵¹ M. Corcoran,⁸¹ F. Couderc,¹⁸ M.-C. Cousinou,¹⁵ S. Crépe-Renaudin,¹⁴ D. Cutts,⁷⁸ M. Ćwiok,³⁰ H. da Motta,² A. Das,⁴⁶ G. Davies,⁴⁴ K. De,⁷⁹ S. J. de Jong,³⁵ E. De La Cruz-Burelo,⁶⁵ C. De Oliveira Martins,³ J. D. Degenhardt,⁶⁵ F. Déliot,¹⁸ M. Demarteau,⁵¹ R. Demina,⁷² D. Denisov,⁵¹ S. P. Denisov,³⁹ S. Desai,⁵¹ H. T. Diehl,⁵¹ M. Diesburg,⁵¹ A. Dominguez,⁶⁸ H. Dong,⁷³ L. V. Dudko,³⁸ L. Duflot,¹⁶ S. R. Dugad,²⁹ D. Duggan,⁵⁰ A. Duperrin,¹⁵ J. Dyer,⁶⁶ A. Dyshkant,⁵³ M. Eads,⁶⁸ D. Edmunds,⁶⁶ J. Ellison,⁴⁹ V. D. Elvira,⁵¹ Y. Enari,⁷⁸ S. Eno,⁶² P. Ermolov,³⁸ H. Evans,⁵⁵ A. Evdokimov,⁷⁴ V. N. Evdokimov,³⁹ A. V. Ferapontov,⁶⁰ T. Ferbel,⁷² F. Fiedler,²⁴ F. Filthaut,³⁵ W. Fisher,⁵¹ H. E. Fisk,⁵¹ M. Ford,⁴⁵ M. Fortner,⁵³ H. Fox,²³ S. Fu,⁵¹ S. Fuess,⁵¹ T. Gadfort,⁷¹ C. F. Galea,³⁵ E. Gallas,⁵¹ E. Galyaev,⁵⁶ C. Garcia,⁷² A. Garcia-Bellido,⁸³ V. Gavrilov,³⁷ P. Gay,¹³ W. Geist,¹⁹ D. Gelé,¹⁹ C. E. Gerber,⁵² Y. Gershtein,⁵⁰ D. Gillberg,⁶ G. Ginther,⁷² N. Gollub,⁴¹ B. Gómez,⁸ A. Goussiou,⁵⁶ P. D. Grannis,⁷³ H. Greenlee,⁵¹ Z. D. Greenwood,⁶¹ E. M. Gregores,⁴ G. Grenier,²⁰ Ph. Gris,¹³ J.-F. Grivaz,¹⁶ A. Grohsjean,²⁵ S. Gründahl,⁵¹ M. W. Grünewald,³⁰ J. Guo,⁷³ F. Guo,⁷³ P. Gutierrez,⁷⁶ G. Gutierrez,⁵¹ A. Haas,⁷¹ N. J. Hadley,⁶² P. Haefner,²⁵ S. Hagopian,⁵⁰ J. Haley,⁶⁹ I. Hall,⁶⁶ R. E. Hall,⁴⁸ L. Han,⁷ P. Hansson,⁴¹ K. Harder,⁴⁵ A. Harel,⁷² R. Harrington,⁶⁴ J. M. Hauptman,⁵⁸ R. Hauser,⁶⁶ J. Hays,⁴⁴ T. Hebbeker,²¹ D. Hedin,⁵³ J. G. Hegeman,³⁴ J. M. Heinmiller,⁵² A. P. Heinson,⁴⁹ U. Heintz,⁶³ C. Hensel,⁵⁹ K. Herner,⁷³ G. Hesketh,⁶⁴ M. D. Hildreth,⁵⁶ R. Hirosky,⁸² J. D. Hobbs,⁷³ B. Hoeneisen,¹² H. Hoeth,²⁶ M. Hohlfeld,²² S. J. Hong,³¹ S. Hossain,⁷⁶ P. Houben,³⁴ Y. Hu,⁷³ Z. Hubacek,¹⁰ V. Hynek,⁹ I. Iashvili,⁷⁰ R. Illingworth,⁵¹ A. S. Ito,⁵¹ S. Jabeen,⁶³ M. Jaffré,¹⁶ S. Jain,⁷⁶ K. Jakobs,²³ C. Jarvis,⁶² R. Jesik,⁴⁴ K. Johns,⁴⁶ C. Johnson,⁷¹ M. Johnson,⁵¹ A. Jonckheere,⁵¹ P. Jonsson,⁴⁴ A. Juste,⁵¹ E. Kajfasz,¹⁵ A. M. Kalinin,³⁶ J. R. Kalk,⁶⁶ J. M. Kalk,⁶¹ S. Kappler,²¹ D. Karmanov,³⁸ P. A. Kasper,⁵¹ I. Katsanos,⁷¹ D. Kau,⁵⁰ R. Kaur,²⁷ V. Kaushik,⁷⁹ R. Kehoe,⁸⁰ S. Kermiche,¹⁵ N. Khalatyan,⁵¹ A. Khanov,⁷⁷ A. Kharchilava,⁷⁰ Y. M. Kharzheev,³⁶ D. Khatidze,⁷¹ T. J. Kim,³¹ M. H. Kirby,⁵⁴ M. Kirsch,²¹ B. Klima,⁵¹ J. M. Kohli,²⁷ J.-P. Konrath,²³ V. M. Korablev,³⁹ A. V. Kozelov,³⁹ D. Krop,⁵⁵ T. Kuhl,²⁴ A. Kumar,⁷⁰ S. Kunori,⁶² A. Kupco,¹¹ T. Kurča,²⁰ J. Kvita,⁹ F. Lacroix,¹³ D. Lam,⁵⁶ S. Lammers,⁷¹ G. Landsberg,⁷⁸ P. Lebrun,²⁰ W. M. Lee,⁵¹ A. Leflat,³⁸ F. Lehner,⁴² J. Lellouch,¹⁷ J. Leveque,⁴⁶ J. Li,⁷⁹ Q. Z. Li,⁵¹ L. Li,⁴⁹ S. M. Lietti,⁵ J. G. R. Lima,⁵³ D. Lincoln,⁵¹ J. Linnemann,⁶⁶ V. V. Lipaev,³⁹ R. Lipton,⁵¹ Y. Liu,⁷ Z. Liu,⁶ A. Lobodenko,⁴⁰ M. Lokajicek,¹¹ P. Love,⁴³ H. J. Lubatti,⁸³ R. Luna,³ A. L. Lyon,⁵¹ A. K. A. Maciel,² D. Mackin,⁸¹ R. J. Madaras,⁴⁷ P. Mättig,²⁶ C. Magass,²¹ A. Magerkurth,⁶⁵ P. K. Mal,⁵⁶ H. B. Malbouisson,³ S. Malik,⁶⁸ V. L. Malyshev,³⁶ H. S. Mao,⁵¹ Y. Maravin,⁶⁰ B. Martin,¹⁴ R. McCarthy,⁷³ A. Melnitchouk,⁶⁷ L. Mendoza,⁸ P. G. Mercadante,⁵ M. Merkin,³⁸ K. W. Merritt,⁵¹ J. Meyer,²² A. Meyer,²¹ T. Millet,²⁰ J. Mitrevski,⁷¹ J. Molina,³ R. K. Mommsen,⁴⁵ N. K. Mondal,²⁹ R. W. Moore,⁶ T. Moulík,⁵⁹ G. S. Muanza,²⁰ M. Mulders,⁵¹ M. Mulhearn,⁷¹ O. Mundal,²² L. Mundim,³ E. Nagy,¹⁵ M. Naimuddin,⁵¹ M. Narain,⁷⁸ N. A. Naumann,³⁵ H. A. Neal,⁶⁵ J. P. Negret,⁸ P. Neustroev,⁴⁰ H. Nilsen,²³ H. Nogima,³ S. F. Novaes,⁵ T. Nunnemann,²⁵ V. O'Dell,⁵¹ D. C. O'Neil,⁶ G. Obrant,⁴⁰ C. Ochando,¹⁶ D. Onoprienko,⁶⁰ N. Oshima,⁵¹ J. Osta,⁵⁶ R. Otec,¹⁰ G. J. Otero y Garzón,⁵¹ M. Owen,⁴⁵ P. Padley,⁸¹ M. Pangilinan,⁷⁸ N. Parashar,⁵⁷ S.-J. Park,⁷² S. K. Park,³¹ J. Parsons,⁷¹ R. Partridge,⁷⁸ N. Parua,⁵⁵ A. Patwa,⁷⁴ G. Pawloski,⁸¹ B. Penning,²³ M. Perfilov,³⁸ K. Peters,⁴⁵ Y. Peters,²⁶ P. Pétróff,¹⁶ M. Petteni,⁴⁴ R. Piegaia,¹ J. Piper,⁶⁶ M.-A. Pleier,²² P. L. M. Podesta-Lerma,³³ V. M. Podstavkov,⁵¹

Y. Pogorelov,⁵⁶ M.-E. Pol,² P. Polozov,³⁷ B. G. Pope,⁶⁶ A. V. Popov,³⁹ C. Potter,⁶ W. L. Prado da Silva,³ H. B. Prosper,⁵⁰ S. Protopopescu,⁷⁴ J. Qian,⁶⁵ A. Quadt,²² B. Quinn,⁶⁷ A. Rakitine,⁴³ M. S. Rangel,² K. Ranjan,²⁸ P. N. Ratoff,⁴³ P. Renkel,⁸⁰ S. Reucroft,⁶⁴ P. Rich,⁴⁵ J. Rieger,⁵⁵ M. Rijssenbeek,⁷³ I. Ripp-Baudot,¹⁹ F. Rizatdinova,⁷⁷ S. Robinson,⁴⁴ R. F. Rodrigues,³ M. Rominsky,⁷⁶ C. Royon,¹⁸ P. Rubinov,⁵¹ R. Ruchti,⁵⁶ G. Safronov,³⁷ G. Sajot,¹⁴ A. Sánchez-Hernández,³³ M. P. Sanders,¹⁷ A. Santoro,³ G. Savage,⁵¹ L. Sawyer,⁶¹ T. Scanlon,⁴⁴ D. Schaile,²⁵ R. D. Schamberger,⁷³ Y. Scheglov,⁴⁰ H. Schellman,⁵⁴ T. Schliephake,²⁶ C. Schwanenberger,⁴⁵ A. Schwartzman,⁶⁹ R. Schwienhorst,⁶⁶ J. Sekaric,⁵⁰ H. Severini,⁷⁶ E. Shabalina,⁵² M. Shamim,⁶⁰ V. Shary,¹⁸ A. A. Shchukin,³⁹ R. K. Shivpuri,²⁸ V. Siccaldi,¹⁹ V. Simak,¹⁰ V. Sirotenko,⁵¹ P. Skubic,⁷⁶ P. Slattery,⁷² D. Smirnov,⁵⁶ J. Snow,⁷⁵ G. R. Snow,⁶⁸ S. Snyder,⁷⁴ S. Söldner-Rembold,⁴⁵ L. Sonnenschein,¹⁷ A. Sopczak,⁴³ M. Sosebee,⁷⁹ K. Soustruznik,⁹ B. Spurlock,⁷⁹ J. Stark,¹⁴ J. Steele,⁶¹ V. Stolin,³⁷ D. A. Stoyanova,³⁹ J. Strandberg,⁶⁵ S. Strandberg,⁴¹ M. A. Strang,⁷⁰ M. Strauss,⁷⁶ E. Strauss,⁷³ R. Ströhmer,²⁵ D. Strom,⁵⁴ L. Stutte,⁵¹ S. Sumowidagdo,⁵⁰ P. Svoisky,⁵⁶ A. Sznajder,³ M. Talby,¹⁵ P. Tamburello,⁴⁶ A. Tanasijczuk,¹ W. Taylor,⁶ J. Temple,⁴⁶ B. Tiller,²⁵ F. Tissandier,¹³ M. Titov,¹⁸ V. V. Tokmenin,³⁶ T. Toole,⁶² I. Torchiani,²³ T. Trefzger,²⁴ D. Tsybychev,⁷³ B. Tuchming,¹⁸ C. Tully,⁶⁹ P. M. Tuts,⁷¹ R. Unalan,⁶⁶ S. Uvarov,⁴⁰ L. Uvarov,⁴⁰ S. Uzunyan,⁵³ B. Vachon,⁶ P. J. van den Berg,³⁴ R. Van Kooten,⁵⁵ W. M. van Leeuwen,³⁴ N. Varelas,⁵² E. W. Varnes,⁴⁶ I. A. Vasilyev,³⁹ M. Vaupel,²⁶ P. Verdier,²⁰ L. S. Vertogradov,³⁶ M. Verzocchi,⁵¹ F. Villeneuve-Seguié,⁴⁴ P. Vint,⁴⁴ P. Vokac,¹⁰ E. Von Toerne,⁶⁰ M. Voutilainen,⁶⁸ R. Wagner,⁶⁹ H. D. Wahl,⁵⁰ L. Wang,⁶² M. H. L. S Wang,⁵¹ J. Warchol,⁵⁶ G. Watts,⁸³ M. Wayne,⁵⁶ M. Weber,⁵¹ G. Weber,²⁴ L. Welty-Rieger,⁵⁵ A. Wenger,⁴² N. Wermes,²² M. Wetstein,⁶² A. White,⁷⁹ D. Wicke,²⁶ G. W. Wilson,⁵⁹ S. J. Wimpenny,⁴⁹ M. Wobisch,⁶¹ D. R. Wood,⁶⁴ T. R. Wyatt,⁴⁵ Y. Xie,⁷⁸ S. Yacoub,⁵⁴ R. Yamada,⁵¹ M. Yan,⁶² T. Yasuda,⁵¹ Y. A. Yatsunenko,³⁶ K. Yip,⁷⁴ H. D. Yoo,⁷⁸ S. W. Youn,⁵⁴ J. Yu,⁷⁹ A. Zatserklyaniy,⁵³ C. Zeitnitz,²⁶ T. Zhao,⁸³ B. Zhou,⁶⁵ J. Zhu,⁷³ M. Zielinski,⁷² D. Zieminska,⁵⁵ A. Zieminski,^{55,*} L. Zivkovic,⁷¹ V. Zutshi,⁵³ and E. G. Zverev³⁸

(D0 Collaboration)

¹Universidad de Buenos Aires, Buenos Aires, Argentina²LAFEX, Centro Brasileiro de Pesquisas Físicas, Rio de Janeiro, Brazil³Universidade do Estado do Rio de Janeiro, Rio de Janeiro, Brazil⁴Universidade Federal do ABC, Santo André, Brazil⁵Instituto de Física Teórica, Universidade Estadual Paulista, São Paulo, Brazil⁶University of Alberta, Edmonton, Alberta, Canada,

Simon Fraser University, Burnaby, British Columbia, Canada,

York University, Toronto, Ontario, Canada,

and McGill University, Montreal, Quebec, Canada

⁷University of Science and Technology of China, Hefei, People's Republic of China⁸Universidad de los Andes, Bogotá, Colombia⁹Center for Particle Physics, Charles University, Prague, Czech Republic¹⁰Czech Technical University, Prague, Czech Republic¹¹Center for Particle Physics, Institute of Physics, Academy of Sciences of the Czech Republic, Prague, Czech Republic¹²Universidad San Francisco de Quito, Quito, Ecuador¹³LPC, Univ Blaise Pascal, CNRS/IN2P3, Clermont, France¹⁴LPSC, Université Joseph Fourier Grenoble 1, CNRS/IN2P3, Institut National Polytechnique de Grenoble, France¹⁵CPPM, IN2P3/CNRS, Université de la Méditerranée, Marseille, France¹⁶LAL, Univ Paris-Sud, IN2P3/CNRS, Orsay, France¹⁷LPNHE, IN2P3/CNRS, Universités Paris VI and VII, Paris, France¹⁸DAPNIA/Service de Physique des Particules, CEA, Saclay, France¹⁹IPHC, Université Louis Pasteur et Université de Haute Alsace, CNRS/IN2P3, Strasbourg, France²⁰IPNL, Université Lyon 1, CNRS/IN2P3, Villeurbanne, France and Université de Lyon, Lyon, France²¹III. Physikalisches Institut A, RWTH Aachen, Aachen, Germany²²Physikalisches Institut, Universität Bonn, Bonn, Germany²³Physikalisches Institut, Universität Freiburg, Freiburg, Germany²⁴Institut für Physik, Universität Mainz, Mainz, Germany²⁵Ludwig-Maximilians-Universität München, München, Germany²⁶Fachbereich Physik, University of Wuppertal, Wuppertal, Germany²⁷Panjab University, Chandigarh, India²⁸Delhi University, Delhi, India²⁹Tata Institute of Fundamental Research, Mumbai, India

- ³⁰University College Dublin, Dublin, Ireland
³¹Korea Detector Laboratory, Korea University, Seoul, Korea
³²SungKyunKwan University, Suwon, Korea
³³CINVESTAV, Mexico City, Mexico
³⁴FOM-Institute NIKHEF and University of Amsterdam/NIKHEF, Amsterdam, The Netherlands
³⁵Radboud University Nijmegen/NIKHEF, Nijmegen, The Netherlands
³⁶Joint Institute for Nuclear Research, Dubna, Russia
³⁷Institute for Theoretical and Experimental Physics, Moscow, Russia
³⁸Moscow State University, Moscow, Russia
³⁹Institute for High Energy Physics, Protvino, Russia
⁴⁰Petersburg Nuclear Physics Institute, St. Petersburg, Russia
⁴¹Lund University, Lund, Sweden, Royal Institute of Technology and Stockholm University, Stockholm, Sweden, and Uppsala University, Uppsala, Sweden
⁴²Physik Institut der Universität Zürich, Zürich, Switzerland
⁴³Lancaster University, Lancaster, United Kingdom
⁴⁴Imperial College, London, United Kingdom
⁴⁵University of Manchester, Manchester, United Kingdom
⁴⁶University of Arizona, Tucson, Arizona 85721, USA
⁴⁷Lawrence Berkeley National Laboratory and University of California, Berkeley, California 94720, USA
⁴⁸California State University, Fresno, California 93740, USA
⁴⁹University of California, Riverside, California 92521, USA
⁵⁰Florida State University, Tallahassee, Florida 32306, USA
⁵¹Fermi National Accelerator Laboratory, Batavia, Illinois 60510, USA
⁵²University of Illinois at Chicago, Chicago, Illinois 60607, USA
⁵³Northern Illinois University, DeKalb, Illinois 60115, USA
⁵⁴Northwestern University, Evanston, Illinois 60208, USA
⁵⁵Indiana University, Bloomington, Indiana 47405, USA
⁵⁶University of Notre Dame, Notre Dame, Indiana 46556, USA
⁵⁷Purdue University Calumet, Hammond, Indiana 46323, USA
⁵⁸Iowa State University, Ames, Iowa 50011, USA
⁵⁹University of Kansas, Lawrence, Kansas 66045, USA
⁶⁰Kansas State University, Manhattan, Kansas 66506, USA
⁶¹Louisiana Tech University, Ruston, Louisiana 71272, USA
⁶²University of Maryland, College Park, Maryland 20742, USA
⁶³Boston University, Boston, Massachusetts 02215, USA
⁶⁴Northeastern University, Boston, Massachusetts 02115, USA
⁶⁵University of Michigan, Ann Arbor, Michigan 48109, USA
⁶⁶Michigan State University, East Lansing, Michigan 48824, USA
⁶⁷University of Mississippi, University, Mississippi 38677, USA
⁶⁸University of Nebraska, Lincoln, Nebraska 68588, USA
⁶⁹Princeton University, Princeton, New Jersey 08544, USA
⁷⁰State University of New York, Buffalo, New York 14260, USA
⁷¹Columbia University, New York, New York 10027, USA
⁷²University of Rochester, Rochester, New York 14627, USA
⁷³State University of New York, Stony Brook, New York 11794, USA
⁷⁴Brookhaven National Laboratory, Upton, New York 11973, USA
⁷⁵Langston University, Langston, Oklahoma 73050, USA
⁷⁶University of Oklahoma, Norman, Oklahoma 73019, USA
⁷⁷Oklahoma State University, Stillwater, Oklahoma 74078, USA
⁷⁸Brown University, Providence, Rhode Island 02912, USA
⁷⁹University of Texas, Arlington, Texas 76019, USA
⁸⁰Southern Methodist University, Dallas, Texas 75275, USA
⁸¹Rice University, Houston, Texas 77005, USA
⁸²University of Virginia, Charlottesville, Virginia 22901, USA
⁸³University of Washington, Seattle, Washington 98195, USA
(Received 26 December 2007; published 3 February 2009)

In a data sample of approximately 1.3 fb^{-1} collected with the D0 detector between 2002 and 2006, the orbitally excited charm state $D_{s1}^{\pm}(2536)$ has been observed with a measured mass of $2535.7 \pm 0.6(\text{stat}) \pm 0.5(\text{syst}) \text{ MeV}/c^2$ via the decay mode $B_s^0 \rightarrow D_{s1}^-(2536)\mu^+\nu_{\mu}X$. A first measurement is made of the branching ratio product $\text{Br}(\bar{b} \rightarrow D_{s1}^-(2536)\mu^+\nu_{\mu}X) \times \text{Br}(D_{s1}^- \rightarrow D^{*-}K_S^0)$. Assuming that $D_{s1}^-(2536)$

production in semileptonic decay is entirely from B_s^0 , an extraction of the semileptonic branching ratio $\text{Br}(B_s^0 \rightarrow D_{s1}^-(2536)\mu^+\nu_\mu X)$ is made.

DOI: 10.1103/PhysRevLett.102.051801

PACS numbers: 13.25.Hw, 14.40.Lb

Semileptonic B_s^0 decays into orbitally excited P -wave strange-charm mesons (D_s^{**}) are expected to make up a significant fraction of B_s^0 semileptonic decays and are therefore important when comparing inclusive and exclusive decay rates, extracting CKM matrix elements, and using semileptonic decays in B_s^0 mixing analyses. For B meson semileptonic decays to heavier excited charm states, more of the available phase space is near zero recoil, increasing the importance of corrections in heavy-quark effective theory (HQET) [1], effectively tested here.

D_s^{**} mesons (also denoted D_{sJ}) are composed of a heavy charm quark and a lighter strange quark in an $L = 1$ state of orbital momentum. In the heavy-quark limit, the spin s_Q of the heavy quark and the total angular momentum, $j_q = s_q + L$ of the light degrees of freedom (quark and gluons), are separately conserved and the latter has possible values of $j_q = \frac{1}{2}$ or $\frac{3}{2}$. The surprisingly light masses of the $j_q = \frac{1}{2}$ states: $D_{s0}^*(2317)$ and $D_{s1}(2460)$ [2], plus the observation of new D_{sJ} states [3], deepens the need for a better understanding of these D_s^{**} systems since they may be quark molecular states, a new and very different arrangement of quarks.

In our decay of interest, the $j_q = \frac{3}{2}$ angular momentum can combine with the heavy-quark spin to form the $J^P = 1^+$ (D_{s1}) state which must decay through a D wave to conserve $j_q = \frac{3}{2}$. The $D_{s1}^\pm(2536)$ is expected to decay

dominantly into a D^* and K meson to conserve angular momentum.

In this Letter we present the first measurement of semileptonic B_s^0 decay into the narrow $D_{s1}^\pm(2536)$ state. This state is just above the $D^*K_S^0$ mass threshold and has been observed previously [4]. Events compatible with the decay chain $\bar{b} \rightarrow D_{s1}^-(2536)\mu^+\nu_\mu X$, $D_{s1}^-(2536) \rightarrow D^{*-}K_S^0$, $D^{*-} \rightarrow \bar{D}^0\pi^-$, $K_S^0 \rightarrow \pi^+\pi^-$, $\bar{D}^0 \rightarrow K^+\pi^-$ are reconstructed. Charge conjugate modes and reactions are always implied in this Letter.

Assuming that $D_{s1}^\pm(2536)$ production in a semileptonic decay is entirely from B_s^0 , the branching ratio $\text{Br}(B_s^0 \rightarrow D_{s1}^\pm(2536)\mu^+\nu_\mu X)$ can be determined by normalizing to the known value of the branching fraction $\text{Br}(\bar{b} \rightarrow D^{*-}\mu^+\nu_\mu X) = (2.75 \pm 0.19)\%$ [5] to avoid uncertainties in the b -quark production rate. This semileptonic branching ratio includes any decay channel or sequence of channels resulting in a D^* and a lepton (muon in our case), and all b hadrons, and therefore includes the relative production of each b hadron species starting from a \bar{b} quark. Since the final state of interest, $D_{s1}^-(2536) \rightarrow D^{*-}K_S^0$, is reconstructed from a D^* and a K_S^0 , the selection is broken up into two sections: one to reconstruct the D^* with an associated muon, coming dominantly from B meson decays resulting in a number of candidates, $N_{D^*\mu}$, and then the addition and subsequent formation of a vertex of a K_S^0 with the D^* and muon, resulting in $N_{D_{s1}}$ candidates. To find the branching ratio, the following formula is used:

$$f(\bar{b} \rightarrow B_s^0)\text{Br}(B_s^0 \rightarrow D_{s1}^-\mu^+\nu_\mu X)\text{Br}(D_{s1}^- \rightarrow D^{*-}K_S^0) = \text{Br}(\bar{b} \rightarrow D^{*-}\mu^+\nu_\mu X) \frac{N_{D_{s1}}}{N_{D^*\mu}} \frac{\epsilon(\bar{b} \rightarrow D^*\mu)}{\epsilon(B_s^0 \rightarrow D_{s1}\mu \rightarrow D^*\mu)} \frac{1}{\epsilon_{K_S^0}}. \quad (1)$$

The input $f(\bar{b} \rightarrow B_s^0)$ [5] is the fraction of decays where a b quark will hadronize to a B_s^0 hadron. $\epsilon_{K_S^0}$ is the efficiency in the signal decay channel to reconstruct and make a vertex with a K_S^0 to form a $D_{s1}(2536)$, given that a D^* and a muon have already been reconstructed. Later we will identify the ratio of efficiencies as $R_{D^*}^{\text{gen}} = \epsilon(B_s^0 \rightarrow D_{s1}\mu \rightarrow D^*\mu)/\epsilon(\bar{b} \rightarrow D^*\mu)$.

The D0 detector [6] and following analysis [7] are described in more detail elsewhere. The main elements relevant to this analysis are the silicon microstrip tracker (SMT), central fiber tracker (CFT), and muon detector systems.

This measurement uses a large data sample, corresponding to approximately 1.3 fb^{-1} of integrated luminosity collected by the D0 detector between April 2002 and March 2006. Events were reconstructed using the standard D0 software suite. To avoid lifetime biases compared to the

MC simulation, the small fraction of events were removed that entered the sample only via triggers that included requirements on impact parameters of tracks.

To evaluate signal mass resolution and efficiencies, Monte Carlo (MC) simulated samples were generated for signal and background. The standard D0 simulation and event reconstruction chain was used. Events were generated with the PYTHIA generator [8] and decay chains of heavy hadrons were simulated with the EVTGEN decay package [9]. The detector response was modeled by GEANT [10]. Two background MC samples were also generated: a $c\bar{c}$ sample, and an inclusive b -quark sample containing all b hadron species with forced semileptonic decays to a muon. In both cases, all events containing both a D^* and a muon were retained.

B mesons were first selected using their semileptonic decays, $B \rightarrow D^{*-}\mu^+X$. At this point in the selection, the

$D^* + \mu$ sample is dominated by $B_d^0 \rightarrow D^{*-} \mu^+ \nu_\mu X$ decays. For this analysis, muons were required to have hits in more than one muon layer, to have an associated track in the central tracking system, and to have transverse momentum $p_T^\mu > 2$ GeV/c, pseudorapidity $|\eta^\mu| < 2$, and total momentum $p^\mu > 3$ GeV/c. Two oppositely charged tracks with $p_T > 0.7$ GeV/c and $|\eta| < 2$ were required to form a common \bar{D}^0 vertex which were then combined with a muon candidate to form a common decay point following the procedure described in Ref. [11]. For each $\bar{D}^0 \mu^+$ candidate, an additional soft pion was searched for with charge opposite to the charge of the muon and $p_T > 0.18$ GeV/c. The K^- and π^+ from the decay of the D^0 were both required to have more than five CFT hits. To reduce the contribution from prompt $c\bar{c}$ production, a requirement was made on the transverse decay length, L_{xy} , significance of the $D^* \mu$ vertex of $L_{xy}/\sigma(L_{xy}) > 1$. After these cuts, the total number of D^* candidates in the mass difference, $M(D^*) - M(D^0)$, peak of Fig. 1 is $N_{D^* \mu} = 87506 \pm 496$ (stat).

$D_{s1}^\pm(2536)$ candidates were formed by combining a D^* candidate with a K_S^0 . D^* candidates were first selected by requiring the mass difference $M(D^*) - M(D^0)$ to be in the range 0.142–0.149 GeV/c². The two tracks from the decay of the K_S^0 were required to have opposite charge and to have more than five hits in the CFT detector. The p_T of the K_S^0 was required to be greater than 1 GeV/c to reduce the contribution of background K_S^0 mesons from fragmentation. A vertex was then formed using the reconstructed K_S^0 and the D^* candidate of the event. The decay length of the K_S^0 was required to be greater than 0.5 cm. To compute the $D_{s1}^\pm(2536)$ invariant mass, a mass constraint was applied using the known $D^{*\pm}$ mass [5] instead of the measured invariant mass of the $K\pi\pi$ system. Finally, the invariant mass of the reconstructed $D_{s1}^\pm(2536)$ and muon was required to be less than the mass of the B_s^0 meson [5].




FIG. 1 (color online). The mass difference $M(D^*) - M(D^0)$ for events with $1.8 < M(D^0) < 1.95$ GeV/c² and an associated muon. The number $N_{D^* \mu}$ was defined as the number of signal events in the mass difference range of 0.142–0.149 GeV/c².

The signal model employed for the fit to the $D^* K_S^0$ invariant mass spectrum was a relativistic Breit-Wigner convoluted with a Gaussian function, with the resonance width fixed to the value $1.03 \pm 0.05(\text{stat}) \pm 0.12(\text{syst})$ MeV/c² measured by the *BABAR* Collaboration [12] and a Gaussian width determined to be 2.8 MeV/c² from MC simulation of the signal. The MC width value was scaled up by a factor of 1.10 ± 0.10 to account for differences between data and MC resolution estimates. The unbinned likelihood fit used an exponential function plus a first-order polynomial to model the background with a threshold cutoff of $M(D^*) + M(K_S^0)$. The fit, shown in Fig. 2, gives a central value for the mass peak of $2535.7 \pm 0.7(\text{stat})$ MeV/c², a yield of $N_{D_{s1}} = 45.9 \pm 9.1(\text{stat})$ events, and a significance of 6.1σ for the background to fluctuate up to or above the observed number of signal events.

The efficiencies used in Eq. (1) are estimated using the MC simulation, after implementing suitable correction factors to ensure proper modeling of the underlying b -hadron p_T spectrum, as well as trigger effects. An event-by-event weight, applied as a function of the generated p_T of the B_s , was determined by comparing the generated $p_T(B)$ in MC with the p_T distribution of fully reconstructed $B^+ \rightarrow J/\psi K^+$ candidates in data collected primarily with a dimuon trigger [13]. Most events for this analysis were recorded using single muon triggers, and an additional weight was applied as a function of $p_T(\mu)$ to further improve the simulation of trigger effects. Reweighted MC events were used in the determination of efficiencies described below, and indicated uncertainties are due to MC statistics.

Using the MC sample of inclusive $\bar{b} \rightarrow D^* \mu X$ events, specific major decay modes were identified. Efficiencies for each of these decay modes to pass the $D^* \mu$ selection, including the efficiency to reconstruct the soft pion from the D^* , were then determined. The predicted fraction F_i of each channel contributing to the $D^* \mu$ sample before fur-




FIG. 2 (color online). Invariant mass of $D^* K_S^0$ with an associated muon. Shown is the result of the fit of the $D^* K_S^0$ mass with the function described in the text.

ther cuts was found following a procedure similar to that given in Ref. [14]. The efficiency ϵ_i for each channel was found and a weighted sum was calculated, giving an estimated total efficiency for reconstruction of $\epsilon(\bar{b} \rightarrow D^* \mu) = (5.88 \pm 0.80)\%$, where the uncertainty is dominated by the MC statistics used to find ϵ_i , and uncertainties on external inputs [5] used to estimate F_i . Applying the same cuts for reconstructing the $D^* \mu$ for the signal channel, the efficiency $\epsilon(B_s^0 \rightarrow D_{s1} \mu \rightarrow D^* \mu) = (3.20 \pm 0.02)\%$, results in a ratio of efficiencies of $R_{D^*}^{\text{gen}} = 0.547 \pm 0.075$.

The signal MC sample was used to determine the efficiency to reconstruct $D_{s1}^-(2536) \rightarrow D^{*-} K_S^0$ given a reconstructed $D^* \mu$ as a starting point. This efficiency is hence effectively that of reconstructing a $K_S^0 \rightarrow \pi^+ \pi^-$ and forming a vertex with the $D^* \mu$, and includes the branching ratio $\text{Br}(K_S^0 \rightarrow \pi^+ \pi^-)$ [5] for ease of use in calculating the branching ratio product. The reconstruction efficiency was found to be $\epsilon_{K_S^0} = (10.3 \pm 0.4)\%$ where the uncertainty is due to MC statistics.

The process $c\bar{c} \rightarrow D^{*-} \mu^+ \nu_\mu X$ can contribute to $N_{D^* \mu}$ since a D^* meson can come from the hadronization of the \bar{c} quark, and the muon can come from the semileptonic decay of the hadron containing the c quark. To determine the number of events in our signal reconstructed from a prompt D^* , a comparison was made of the decay length significance distribution observed in the data with the same distribution predicted by MC for $b \rightarrow D^* \mu X$ and any excess at shorter significances was interpreted as $c\bar{c}$ contribution. For the decay length significance cut used in the analysis, $L_{xy}/\sigma(L_{xy}) > 1$, the fraction of $N_{D^* \mu}$ from $c\bar{c}$ production was estimated to be $(3.9 \pm 2.5)\%$. A check using a prompt $c\bar{c}$ MC sample results in a consistent estimate. The value of $N_{D^* \mu}$ was corrected downward accordingly.

The contribution from $c\bar{c}$ production to $N_{D_{s1}}$ where one charm quark hadronizes directly to a $D_{s1}(2536)$ and the other decays directly to a muon was estimated to be negligible using relative production ratios and spin-counting arguments [15].

Systematic uncertainties for the branching ratio product are summarized in Table I and discussed below. The uncertainty in the normalizing branching ratio [5] $\text{Br}(\bar{b} \rightarrow D^* \mu X)$ was taken as a systematic uncertainty. For determining $N_{D^* \mu}$, the signal and background model parameters

$$f(\bar{b} \rightarrow B_s^0) \text{Br}(B_s^0 \rightarrow D_{s1}^- \mu^+ \nu_\mu X) \text{Br}(D_{s1}^- \rightarrow D^{*-} K_S^0) = [2.66 \pm 0.52(\text{stat}) \pm 0.45(\text{syst})] \times 10^{-4}.$$

To assess the systematic uncertainty on the mass measurement, the same variations of the $D_{s1}(2536)$ mass signal model, as well as background functional form, were applied as described above. The mass values used for the mass constraints on the decay products were varied within their PDG uncertainties and were also set to the D0 central fit values. Ensemble tests indicated that the statistical error is correct. From the observed variations, a total systematic

TABLE I. Estimated systematic uncertainties.

Source	Systematic uncertainty
$\text{Br}(\bar{b} \rightarrow D^* \mu X)$	6.9%
$N_{D^* \mu}$	2.9%
$N_{D_{s1}}$	5.5%
$\epsilon_{K_S^0}$	11.0%
$R_{D^*}^{\text{gen}}$	8.6%
Total	16.8%

were varied in a correlated fashion and a systematic uncertainty was assigned. The estimated $c\bar{c}$ production contribution was varied by the indicated uncertainty. In the determination of $N_{D_{s1}}$, the functional forms of the signal and background models were varied in a number of ways to determine the sensitivity of the candidate yield. In addition, the scaling of the widths was varied by $\pm 10\%$ to check the sensitivity to uncertainty in mass resolution.

By comparing the $p_T(\mu)$ distribution for the signal using the default ISGW2 decay model [16] to the HQET semileptonic decay model [9], a weighting factor was found and applied to the fully simulated signal MC events, and the efficiency determined again. The difference observed was assigned as a contribution to the systematic uncertainty of $\epsilon_{K_S^0}$ and $R_{D^*}^{\text{gen}}$.

When estimating $\epsilon_{K_S^0}$, the uncertainty due to modeling of the b hadron p_T spectrum was derived by using an alternate weighting technique. The cuts on the p_T and decay length of the K_S^0 were varied and a systematic uncertainty on the efficiency due to this source was also assigned. Discrepancies in track reconstruction efficiencies between data and MC in low- p_T tracks were accounted for by assigning a systematic uncertainty to each of the pion tracks in the K_S^0 reconstruction [17,18].

The uncertainty in $R_{D^*}^{\text{gen}}$ is due to a combination of MC statistics and uncertainties in PDG branching ratio values and production fractions, $f(\bar{b} \rightarrow b \text{ hadron})$. The uncorrelated systematic uncertainty is given in Table I.

The estimated systematic uncertainties were added in quadrature to obtain a total estimated systematic uncertainty on the branching ratio product of 16.8%. The branching ratio product was determined to be:

mass uncertainty of $0.5 \text{ MeV}/c^2$ was taken, for a mass measurement of:

$$m(D_{s1}) = 2535.7 \pm 0.6(\text{stat}) \pm 0.5(\text{syst}) \text{ MeV}/c^2.$$

This measured mass value is in good agreement with the PDG average value of $2535.34 \pm 0.31 \text{ MeV}/c^2$ [5].

To allow comparison of this measurement to theoretical predictions, the semileptonic branching ratio alone as

TABLE II. Experimental measurement compared with various theoretical predictions.

Source	$\text{Br}(B_s^0 \rightarrow D_{s1}^-(2536)\mu^+\nu_\mu X)$
This result	$[1.03 \pm 0.20(\text{stat}) \pm 0.17(\text{syst}) \pm 0.14(\text{prod.frac.})]\%$
Theoretical Predictions	$\text{Br}(B_s^0 \rightarrow D_{s1}^-(2536)\mu^+\nu_\mu)$
ISGW2 [1]	$(0.53 \pm 0.27)\%$
Relativistic Quark Model & $1/m_Q$ corrections [19]	$(1.06 \pm 0.16)\%$
Nonrel. HQET and ISGW [20]	0.195%

shown in Table II is extracted by taking the hadronization fraction into B_s^0 as $f(\bar{b} \rightarrow B_s^0) = 0.103 \pm 0.014$ [5] and also assuming that $\text{Br}(D_{s1}(2536) \rightarrow D^*K_S^0) = 0.25$ [9]. This is the first experimental measurement of this semi-leptonic branching ratio and is compared to a number of theoretical predictions [1,19,20] of the exclusive rate in Table II. The systematic uncertainty on this quantity is as described earlier, and the error labeled “(prod. frac.)” is due to the current uncertainty on $f(\bar{b} \rightarrow B_s^0)$. The first two theoretical predictions include relativistic and $1/m_Q$ corrections, while the third does not. The result is found to be consistent within uncertainties with the first two theoretical predictions, and demonstrates the need for such corrections.

In summary, using 1.3 fb^{-1} of integrated luminosity collected with the D0 detector, a first measurement of the semileptonic B_s^0 decay into the narrow $D_{s1}^\pm(2536)$ state has been made and compared with theory. In addition, the mass of the $D_{s1}^\pm(2536)$ was measured and found to be in good agreement with the PDG value.

We thank the staffs at Fermilab and collaborating institutions, and acknowledge support from the DOE and NSF (USA); CEA and CNRS/IN2P3 (France); FASI, Rosatom and RFBR (Russia); CAPES, CNPq, FAPERJ, FAPESP, and FUNDUNESP (Brazil); DAE and DST (India); Colciencias (Colombia); CONACyT (Mexico); KRF and KOSEF (Korea); CONICET and UBACyT (Argentina); FOM (The Netherlands); Science and Technology Facilities Council (United Kingdom); MSMT and GACR (Czech Republic); CRC Program, CFI, NSERC and WestGrid Project (Canada); BMBF and DFG (Germany); SFI (Ireland); The Swedish Research Council (Sweden); CAS and CNSF (China); Alexander von Humboldt Foundation; and the Marie Curie Program.

*Deceased.

- [1] D. Scora and N. Isgur, Phys. Rev. D **52**, 2783 (1995), using the results of Ref. [20] for input.
- [2] B. Aubert *et al.* (BABAR Collaboration), Phys. Rev. Lett. **90**, 242001 (2003); D. Besson *et al.* (CLEO Collaboration), Phys. Rev. D **68**, 032002 (2003); P. Krokovny *et al.* (Belle Collaboration), Phys. Rev. Lett. **91**, 262002 (2003).
- [3] Y. Kubota *et al.* (CLEO Collaboration) Phys. Rev. Lett. **72**, 1972 (1994); A. V. Evdokimov *et al.* (SELEX Collaboration), *ibid.* **93**, 242001 (2004); B. Aubert *et al.* (BABAR Collaboration), *ibid.* **97**, 222001 (2006); J. Brodzicka *et al.* (Belle Collaboration), *ibid.* **100**, 092001 (2008).
- [4] A. H. Heister *et al.* (ALEPH Collaboration), Phys. Lett. B **526**, 34 (2002); A. H. Heister *et al.*, (OPAL Collaboration), Z. Phys. C **76**, 425 (1997); J. Alexander *et al.* (CLEO Collaboration), Phys. Lett. B **303**, 377 (1993); H. Albrecht *et al.* (ARGUS Collaboration), *ibid.* **297**, 425 (1992); B. Aubert *et al.* (BABAR Collaboration), Phys. Rev. D **74**, 032007 (2006).
- [5] W.-M. Yao *et al.*, J. Phys. G **33**, 1 (2006).
- [6] V. Abazov *et al.* (D0 Collaboration), Nucl. Instrum. Methods Phys. Res., Sect. A **565**, 463 (2006).
- [7] J. Rieger, Report No. FERMILAB-THESIS-2007-46, 2007.
- [8] T. Sjöstrand *et al.*, Comput. Phys. Commun. **135**, 238 (2001).
- [9] D. J. Lange, Nucl. Instrum. Methods Phys. Res., Sect. A **462**, 152 (2001). For the signal, the ISGW2 semileptonic decay model [18] is used, and the VVS_PWAVE option for the spin-1 D_{s1} decay.
- [10] R. Brun *et al.*, CERN Report No. DD/EE/84-1, 1984.
- [11] V. Abazov *et al.* (D0 Collaboration), Phys. Rev. Lett. **94**, 182001 (2005).
- [12] B. Aubert *et al.* (BABAR Collaboration), arXiv:hep-ex/0607084.
- [13] V. Abazov *et al.* (D0 Collaboration), Phys. Rev. Lett. **94**, 071802 (2005).
- [14] V. Abazov *et al.* (D0 Collaboration), Phys. Rev. D **74**, 112002 (2006).
- [15] D. Acosta *et al.* (CDF Collaboration), Phys. Rev. Lett. **91**, 241804 (2003); A. F. Falk and M. E. Peskin, Phys. Rev. D **49**, 3320 (1994).
- [16] N. Isgur, D. Scora, B. Grinstein, and M. B. Wise, Phys. Rev. D **39**, 799 (1989).
- [17] V. Abazov *et al.* (D0 Collaboration), Phys. Rev. Lett. **94**, 232001 (2005).
- [18] V. Abazov *et al.* (D0 Collaboration), Phys. Rev. Lett. **95**, 171803 (2005).
- [19] D. Ebert, R. N. Faustov, and V. O. Galkin, Phys. Rev. D **61**, 014016 (1999).
- [20] H. B. Mayorga, A. Moreno Briceno, and J. H. Munoz, J. Phys. G **29**, 2059 (2003).