



OPEN ACCESS

EDITED BY

Sonia Jangra,
The Rockefeller University, United States

REVIEWED BY

Ekaterini Simoes Goudouris,
Federal University of Rio de Janeiro, Brazil

*CORRESPONDENCE

Manuela Berto Pucca
✉ manuela.pucca@unesp.br

RECEIVED 29 June 2025

ACCEPTED 15 September 2025

PUBLISHED 30 September 2025

CORRECTED 06 May 2026

CITATION

Filardi ETM, Carbonell RCC, Pavan FR,
Cerni FA and Pucca MB (2025) One hundred
years of BCG: the journey of tuberculosis
vaccination in Brazil.
Front. Immunol. 16:1655969.
doi: 10.3389/fimmu.2025.1655969

COPYRIGHT

© 2025 Filardi, Carbonell, Pavan, Cerni and
Pucca. This is an open-access article
distributed under the terms of the [Creative
Commons Attribution License \(CC BY\)](#). The
use, distribution or reproduction in other
forums is permitted, provided the original
author(s) and the copyright owner(s) are
credited and that the original publication in
this journal is cited, in accordance with
accepted academic practice. No use,
distribution or reproduction is permitted
which does not comply with these terms.

One hundred years of BCG: the journey of tuberculosis vaccination in Brazil

Eloise Trostdorf Monteiro Filardi¹,
Roberto Carlos Cruz Carbonell^{1,2}, Fernando Rogério Pavan¹,
Felipe Augusto Cerni² and Manuela Berto Pucca^{1*}

¹Graduate Program in Bioscience and Biotechnology Applied to Pharmacy, School of
Pharmaceutical Sciences, São Paulo State University (UNESP), Araraquara, São Paulo, Brazil, ²Medical
School, Federal University of Roraima, Boa Vista, Brazil

KEYWORDS

BCG, bacillus Calmette-Guérin, vaccine, *Mycobacterium tuberculosis*, immunization

Vaccines represent one of the most significant achievements in medical science, constituting safe preparations or substances developed with the primary purpose of stimulating the immune system (1, 2). Their central function is to protect the human body against a myriad of infectious diseases that can be prevented through immunization. From an educational perspective, vaccines can be described as biological preparations that “train” the immune system to recognize and mount an effective defense against specific pathogens (e.g., viruses or bacteria) prior to natural exposure, thereby preventing the onset of disease (3).

The relevance of vaccines transcends individual protection, extending to a considerable socioeconomic impact. When employed as a public health strategy, vaccines are widely considered one of the best investments available, primarily due to their exceptional cost-effectiveness (4). The benefits of vaccination extend far beyond the immediate prevention of disease in individuals. By enabling large-scale disease control, vaccines significantly reduce the burden on healthcare systems, leading to fewer hospital admissions, diminished reliance on high-cost medical interventions, and reduced demand for long-term care services. Moreover, individuals protected through immunization are generally healthier and more capable of contributing to the workforce, thereby enhancing economic productivity at both community and national levels (5, 6).

The history of vaccination is a narrative of scientific triumphs and transformative impact on global health. Over the past five decades, global immunization efforts have resulted in an extraordinary achievement. It is estimated that at least 154 million lives have been saved, the vast majority of them (approximately 101 million) being children (7). A particularly eloquent testament to this success is the impact of measles vaccination, which alone accounted for about 60% of those lives saved, equivalent to approximately 94 million children protected since 1974 (8, 9).

This year marks a significant milestone, a century of the Bacillus Calmette-Guérin (BCG) vaccination in Brazil, a pivotal tool in the fight against tuberculosis (TB), a disease responsible for thousands of deaths worldwide. Officially introduced in Brazil in 1925, the BCG vaccine became a symbol of prevention and public health advancement against one of the most lethal infectious diseases globally (10, 11). However, the vaccine's origin dates to

1921 (ie., four years before), when French scientists Albert Calmette and Camille Guérin, working at the Pasteur Institute in Paris, developed the immunizing agent after years of dedicated research. The scientists attenuated a strain of *Mycobacterium bovis*, a close relative of *M. tuberculosis*, the pathogen responsible for human tuberculosis. Over the course of 13 years, they subjected the bacterium to 231 serial passages in subcultures containing bile, gradually reducing its virulence while retaining its immunogenic capacity. The outcome was a live attenuated vaccine capable of inducing protective immunity without causing disease, first administered to humans in 1921 and subsequently adopted worldwide (12).

In Brazil, before the introduction of BCG vaccine (1925) and, later, the development of effective therapeutics from the mid-20th century onwards, the tuberculosis scenario in Brazil was critical. The disease, often called the “white plague,” was one of the leading causes of death, surrounded by fear and social stigma (13). In the absence of specific treatments, therapeutic approaches were largely palliative and of limited efficacy. In that time, the main strategy for managing patients consisted of isolation in sanatoria, institutions typically built in locations considered healthy climates (mountains or coast), where prolonged rest, enhanced nutrition, and exposure to fresh air, known as “climatotherapy,” were prescribed (14). Although these represented an effort to care for patients and attempt to contain disease spread, sanatoria were insufficient to meet demand and inaccessible to most of the affected population, especially the poor. The effectiveness of these measures in curing the disease was questionable, serving more as palliation and a form of segregation (15).

The introduction of the BCG vaccine in Brazil occurred relatively soon after its development in France, demonstrating a proactive response from the Brazilian scientific and medical community to the serious tuberculosis problem. The protagonist of this introduction was the physician and researcher Arlindo de Assis, who brought the BCG strain from Paris to Brazil in 1925 (Figure 1A). Linked to the Brazilian League against Tuberculosis, Assis initiated the production and application of the vaccine in the country, initially via the oral route, a method he advocated for many years (16). Subsequent production and research efforts involved renowned institutions such as the Oswaldo Cruz Institute (Fiocruz) (17), consolidating the vaccine’s presence in the national scenario from the late 1920s and early 1930s. Brazil’s early engagement with BCG technology, facilitated by figures like Arlindo de Assis and institutions like Fiocruz, signaled recognition of the magnitude of the TB problem and an openness to incorporating new scientific interventions, laying the foundation for the future integration of BCG into national immunization programs. Despite significant progress in the fight against TB, the disease remains a major global health challenge. The variable efficacy of BCG vaccine, particularly in adults, and the increasing prevalence of *M. tuberculosis* strains resistant to first-line drugs are critical concerns (18). These biomedical challenges are further compounded by persistent social determinants of health, including overcrowded living conditions, malnutrition, and

limited access to timely diagnosis and effective treatment in underserved populations (19, 20).

Another important historical milestone in Brazil occurred in 1975, when tuberculosis was officially included among the diseases subject to compulsory notification. This was established by Federal Law No. 6,259, enacted on October 30, 1975, which organized the national system of epidemiological surveillance. The law was later regulated by Decree No. 78,231 of August 12, 1976 (21, 22). From that point on, all diagnosed cases of tuberculosis had to be reported to health authorities. This measure played a crucial role in strengthening public health efforts, allowing for more accurate monitoring of the disease, improved resource allocation, and the implementation of more effective tuberculosis control strategies nationwide (23–25).

Although TB is a preventable and curable disease, it remains one of the leading causes of death from infectious diseases worldwide, second only to COVID-19 during certain periods. In 2023, it is estimated that approximately 1.25 million people died from TB globally (26). In Brazil, 5,845 TB-related deaths were reported in 2022, corresponding to a mortality rate of 2.72 per 100,000 inhabitants, the highest recorded in over two decades of surveillance (27, 28). The high lethality of TB is often associated with factors such as delayed diagnosis, treatment abandonment, social vulnerability, and coinfections, particularly TB-HIV, which significantly worsens patient prognosis (29). These indicators highlight the urgent need for integrated strategies focused on surveillance, early diagnosis, treatment adherence, and public policies aimed at reducing social inequalities.

A temporal analysis of new reported tuberculosis (TB) cases in Brazil from 2013 to 2023, together with national BCG vaccination coverage, is shown in Figure 1B. The data demonstrate that, despite a decline in vaccination coverage during the COVID-19 pandemic, there was no immediate proportional increase in TB notifications. However, a marked rise in reported cases is observed in 2022, likely related to the resumption of surveillance activities and improved case detection.

BCG vaccination coverage in Brazil remained consistently high until 2018, with rates above 95% (Figure 1). From 2019 onward, a gradual decline was observed, becoming more pronounced between 2020 and 2021, when coverage dropped below 80%, despite its mandatory status under Federal Law. This decrease is likely associated with disruptions in health services, logistical challenges, and vaccine hesitancy exacerbated by the COVID-19 pandemic, as previously discussed by Filardi et al. (37). Importantly, despite the drop in coverage, no immediate proportional increase in TB cases was evident during this period. This highlights the critical need to restore and maintain high BCG vaccination coverage and to strengthen epidemiological surveillance systems to ensure timely case detection and control. Encouragingly, coverage rose to 83.5% in 2023, and preliminary data for 2024 indicate a further increase to 92.8%, suggesting a recovery toward pre-pandemic vaccination levels (30).

Vaccination remains a crucial tool in the global fight against TB, particularly given the disease’s persistent lethality and impact on




FIGURE 1
Historical and epidemiological context of BCG vaccination and tuberculosis in Brazil. (A) Timeline illustrating key milestones in BCG implementation in Brazil. (B) Temporal trends in new reported tuberculosis (TB) cases and national BCG vaccination coverage (2013–2023).

vulnerable populations. The BCG vaccine, although not fully effective in preventing pulmonary TB in adults, plays a fundamental role in protecting children against the most severe and life-threatening forms of the disease, such as tuberculous meningitis and miliary TB. By reducing the incidence of these severe manifestations, BCG contributes directly to lowering TB-associated mortality and alleviating the overall burden on public

health systems. Mass immunization programs, therefore, are essential to prevent avoidable deaths and to control the progression of TB, especially in high-risk communities (31, 32).

Although BCG vaccination coverage is not universal and has shown a concerning decline in several regions over recent years, it remains the most effective public health intervention for preventing severe forms of TB in children, particularly extrapulmonary

manifestations. Robust evidence from observational studies and meta-analyses supports the vaccine's efficacy, with protection ranging from 70% to 80% against these critical forms. A systematic review by the Cochrane Collaboration (2014) demonstrated a 73% reduction in the risk of tuberculous meningitis and a 77% reduction in the incidence of miliary TB among vaccinated individuals. These findings reinforce the importance of maintaining high BCG vaccination coverage as a cornerstone of global TB control strategies, particularly in endemic regions. As illustrated in the graphs for Brazil, while BCG coverage remained above 95% until 2015, a subsequent decline may jeopardize these public health gains and is potentially associated with the observed fluctuations in TB mortality. Together, these data emphasize the urgent need to strengthen immunization efforts to ensure sustained protection against severe TB and prevent a resurgence in disease-related mortality (33).

Looking ahead, scientific innovation stands as a crucial driver in the global effort to eliminate TB. Promising candidates for next-generation vaccines are in advanced stages of development, aiming to provide broader and more durable protection across age groups and geographic settings (34). In parallel, advances in molecular diagnostics and digital health technologies are enhancing early detection, while targeted public health campaigns are raising awareness and promoting adherence to treatment regimens. Together, these integrated strategies are essential for accelerating progress toward TB control and, ultimately, eradication (35, 36).

As Brazil marks a century of tuberculosis vaccination (1925–2025), this milestone serves not only as a testament to the progress achieved but also as a call to address the persistent challenges that remain. Sustained investment in public health infrastructure, research, and equitable access to healthcare, combined with ongoing scientific innovation, will be essential to ensuring that, in the coming century, tuberculosis is no longer a threat to public health in Brazil or across the globe.

Author contributions

EF: Conceptualization, Writing – review & editing, Writing – original draft, Investigation. RC: Investigation, Writing – review & editing. FP: Investigation, Writing – review & editing. FC: Writing – review & editing, Investigation. MP: Investigation, Writing – original draft, Writing – review & editing, Resources, Conceptualization, Supervision.

References

1. *Vaccines and immunization: What is vaccination?*. Available online at: <https://www.who.int/news-room/questions-and-answers/item/vaccines-and-immunization-what-is-vaccination> (Accessed May 8, 2025).
2. National Research Council (US) Division of Health Promotion and Disease Prevention. *Vaccine Supply and Innovation* Washington (DC): National Academies

Funding

The author(s) declare financial support was received for the research and/or publication of this article. This study was financed in part by the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior - Brasil (CAPES) - Finance Code 001. We thank Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP; scholarship to ETMF. No. 2024/13258–0).

Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

The author(s) declared that they were an editorial board member of *Frontiers*, at the time of submission. This had no impact on the peer review process and the final decision.

Correction note

A correction has been made to this article. Details can be found at: [10.3389/fimmu.2026.1858910](https://doi.org/10.3389/fimmu.2026.1858910).

Generative AI statement

The author(s) declare that no Generative AI was used in the creation of this manuscript.

Any alternative text (alt text) provided alongside figures in this article has been generated by *Frontiers* with the support of artificial intelligence and reasonable efforts have been made to ensure accuracy, including review by the authors wherever possible. If you identify any issues, please contact us.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

- Press (US). (1985) 2, *Vaccines: Past, Present, and Future*. Available online at: <https://www.ncbi.nlm.nih.gov/books/NBK216821>.

3. Ghattas M, Dwivedi G, Lavertu M, Alameh M-G. Vaccine technologies and platforms for infectious diseases: current progress, challenges, and opportunities. *Vaccines (Basel)*. (2021) 9:1490. doi: [10.3390/vaccines9121490](https://doi.org/10.3390/vaccines9121490)

4. Nandi A, Shet A. Why vaccines matter: understanding the broader health, economic, and child development benefits of routine vaccination. *Hum Vaccin Immunother.* (2020) 16:1900–4. doi: 10.1080/21645515.2019.1708669
5. Rodrigues CMC, Plotkin SA. Impact of vaccines; health, economic and social perspectives. *Front Microbiol.* (2020) 11:1526. doi: 10.3389/fmicb.2020.01526
6. Quilici S, Smith R, Signorelli C. Role of vaccination in economic growth. *J Mark Access Health Policy.* (2015) 3:10. doi: 10.3402/jmahp.v3.27044
7. Global immunization efforts have saved at least 154 million lives over the past 50 years. Available online at: <https://www.who.int/news/item/24-04-2024-global-immunization-efforts-have-saved-at-least-154-million-lives-over-the-past-50-years> (Accessed May 8, 2025).
8. Minta AA. Progress toward measles elimination — Worldwide, 2000–2023. *MMWR Morb Mortal Wkly Rep.* (2024) 73:1036–42. doi: 10.15585/mmwr.mm7345a4
9. UNICEF. (2025). Measles cases are spiking globally. UNICEF. Available online at: <https://www.unicef.org/stories/measles-cases-spiking-globally> (Accessed April 23, 2025).
10. Institut Pasteur. BCG vaccine: the first tuberculosis vaccination took place a century ago. Institut Pasteur (2021). Available online at: <https://www.pasteur.fr/en/home/research-journal/news/bcg-vaccine-first-tuberculosis-vaccination-took-place-century-ago> (Accessed September 23, 2025).
11. BCG. Ministério da Saúde. Available online at: <https://www.gov.br/saude/pt-br/assuntos/saude-de-a-a-z/b/bcg/bcg> (Accessed May 12, 2025).
12. Vacina BCG: a primeira vacinação contra tuberculose ocorreu há um século. Available online at: https://www.pasteur-fr.translate.google.com/en/home/research-journal/news/bcg-vaccine-first-tuberculosis-vaccination-took-place-century-ago?_x_tr_sl=en&_x_tr_tl=pt&_x_tr_hl=pt&_x_tr_pto=sge (Accessed May 12, 2025).
13. Bertolli Filho C. *Social history of tuberculosis and the tuberculous: 1900–1950*. Rio de Janeiro: FIOCRUZ Press. (2001). Available online at: <https://books.scielo.org/id/4> (Accessed September 23, 2025).
14. Diniz JPDF. A Climatoterapia e o tratamento sanatorial. A busca pelo tratamento da tuberculose em Belo Horizonte – MG (1920–1950). *Contraponto.* (2021) 10:404.
15. Hijjar MA, Gerhardt G, Teixeira GM, Procópio MJ. Retrospecto do controle da tuberculose no Brasil. *Rev Saúde Pública.* (2007) 41:50–7. doi: 10.1590/S0034-89102007000800008
16. Agência Brasil. BCG: 100 years – how the vaccine changed the history of tuberculosis control. Agência Brasil (2021). Available online at: <https://agenciabrasil.ebc.com.br/saude/noticia/2021-07/bcg-100-anos-como-vacina-mudou-historia-no-combate-tuberculose> (Accessed September 23, 2025).
17. Rosenberg J. Tuberculose - Aspectos históricos, realidades, seu romantismo e transculturação. *Boletim Pneumologia Sanitária.* (1999) 7:5–29. doi: 10.5123/S0103-460X1999000200002
18. Bloom BR, Atun R, Cohen T, Dye C, Fraser H, Gomez GB, et al. Tuberculosis. In: Holmes KK, Bertozzi S, Bloom BR, Jha P, editors. *Major Infectious Diseases*. The International Bank for Reconstruction and Development/The World Bank, Washington (DC) (2017). Available online at: <http://www.ncbi.nlm.nih.gov/books/NBK525174/>.
19. . The urgency of investing in health systems in Latin America and the Caribbean. (United Nations, Paho). (2024). Available online at: <https://www.paho.org/en/documents/urgency-investing-health-systems-latin-america-and-caribbean-reduce-inequality-and>
20. Gulumbe BH, Abdulrahman A, Ahmad SK, Lawan KA, Danlami MB. WHO report signals tuberculosis resurgence: Addressing systemic failures and revamping control strategies. *Decoding Infect Transm.* (2025) 3:100044. doi: 10.1016/j.dcit.2025.100044
21. Teixeira M da G, Penna GO, Risi JB, Penna ML, Alvim MF, de MJC, et al. Seleção das doenças de notificação compulsória: critérios e recomendações para as três esferas de governo. *Informe Epidemiol do Sus.* (1998) 7:7–28. doi: 10.5123/S0104-16731998000100002
22. BRASIL. *Dispõe sobre a organização das ações de Vigilância Epidemiológica, o Programa Nacional de Imunizações, estabelece a obrigatoriedade de notificação compulsória de doenças e dá outras providências.* (Brazil: National Immunization Program (PNI)) (1975).
23. Notificação Compulsória. Ministério da Saúde. Available online at: <https://www.gov.br/saude/pt-br/composicao/svsa/notificacao-compulsoria/notificacao-compulsoria> (Accessed May 20, 2025).
24. Bartholomay P, Oliveira GPd, Pinheiro RS, Vasconcelos AMN. Melhoria da qualidade das informações sobre tuberculose a partir do relacionamento entre bases de dados. *Cad Saúde Pública.* (2014) 30:2459–70. doi: 10.1590/0102-311X00116313
25. Bartholomay P, Oliveira GPde, Pinheiro RS, Vasconcelos AMN. Melhoria da qualidade das informações sobre tuberculose a partir do relacionamento entre bases de dados. *Cadernos De Saúde Pública.* (2014) 30(11):2459–70. doi: 10.1590/0102-311X00116313
26. Miranda G. (2024). Tuberculosis surpasses COVID-19 and becomes the leading cause of death from infectious disease worldwide. Medscape. Available online at: <https://portugues.medscape.com/verartigo/6511893?form=fpf> (Accessed September 23, 2025).
27. Associação Paulista de Medicina. Eighty thousand people fall ill from tuberculosis annually in Brazil. APM (2024). Available online at: <https://www.apm.org.br/oitenta-mil-pessoas-adoecem-por-tuberculose-anualmente-no-brasil/> (Accessed September 23, 2025).
28. CNN Brasil. Almost 16 people died from tuberculosis per day in Brazil in 2022. CNN Brasil (2024). Available online at: <https://www.cnnbrasil.com.br/saude/quase-16-pessoas-morreram-de-tuberculose-por-dia-no-brasil-em-2022/> (Accessed September 23, 2025).
29. Leite OH. Tuberculosis deaths in Brazil reach a record number in nearly two decades. *Jornal da USP* (2023). Available online at: <https://jornal.usp.br/atualidades/mortes-por-tuberculose-no-brasil-atingem-numero-recorde-em-quase-duas-decadas/> (Accessed September 23, 2025).
30. Vacinação do Calendário Nacional. Ministério da Saúde do Brasil(2024). Available online at: https://infoms.saude.gov.br/extensions/SEIDIGI_DEMAS_VACINACAO_CALENDARIO_NACIONAL_MENU_COBERTURA/SEIDIGI_DEMAS_VACINACAO_CALENDARIO_NACIONAL_MENU_COBERTURA.html (Accessed June 28, 2025).
31. BUTANTAN. BCG vaccine: Immunization should be administered within the first hours of life and is the best strategy to protect children from severe forms of tuberculosis. Instituto Butantan. (2024). Available online at: <https://butantan.gov.br/noticias/vacina-bcg-imunizante-deve-ser-aplicado-nas-primeiras-horas-de-vida-e-e-melhor-estrategia-para-protger-criancas-das-formas-graves-da-tuberculose> (Accessed September 23, 2025).
32. Pereira SM, Dantas OMS, Ximenes R, Barreto ML. Vacina BCG contra tuberculose: efeito protetor e políticas de vacinação. *Rev Saúde Pública.* (2007) 41:59–66. doi: 10.1590/S0034-89102007000800009
33. Trunz BB, Fine P, Dye C. Effect of BCG vaccination on childhood tuberculous meningitis and military tuberculosis worldwide: a meta-analysis and assessment of cost-effectiveness. *Lancet.* (2006) 367:1173–80. doi: 10.1016/S0140-6736(06)68507-3
34. Junqueira-Kipnis AP, Leite LC de C, Croda J, Chimara E, Carvalho ACC, Arcêncio RA. Advances in the development of new vaccines for tuberculosis and Brazil's role in the effort forward the end TB strategy. *Mem Inst Oswaldo Cruz.* (2024) 119:e240093. doi: 10.1590/0074-02760240093
35. Ahmed MM, Okesanya OJ, Olaleke NO, Adigun OA, Adebayo UO, Oso TA, et al. Integrating digital health innovations to achieve universal health coverage: promoting health outcomes and quality through global public health equity. *Healthcare.* (2025) 13:1060. doi: 10.3390/healthcare13091060
36. National Academies of Sciences, Engineering, and Medicine, Health and Medicine Division, Board on Global Health, and Forum on Microbial Threats, Nicholson A, Liao J, Biffl C. Detection. In: *Innovations for Tackling Tuberculosis in the Time of COVID-19: Proceedings of a Workshop*. National Academies Press (US) (2022). Available online at: <https://www.ncbi.nlm.nih.gov/books/NBK587312/>.
37. Filardi ETM, Pucca MB, Araujo Junior JP, Costa PID. Pandemic paradox: the impact of the COVID-19 on the global and Brazilian tuberculosis epidemics. *Front Public Health.* (2024) 12. doi: 10.3389/fpubh.2024.1399860