

Research Article

Variation of the Equator due to a Highly Inclined and Eccentric Disturber

Clair do Nascimento and Tadashi Yokoyama

Universidade Estadual Paulista, IGCE-DEMAC Caixa Postal 178, CEP 13.500-970 Rio Claro, SP, Brazil

Correspondence should be addressed to Tadashi Yokoyama, tadashi@ms.rc.unesp.br

Received 18 June 2009; Accepted 19 October 2009

Recommended by Dane Quinn

The New Horizons project is currently in progress, but until recently years ago, a mission to Neptune-Triton System predicted in the mid term 2008–2013 was one of the priorities of NASA's Solar System Exploration theme. Any way, it is important to increase our knowledge on the dynamics of the inner satellites of Neptune, since according to some authors, this was a key question in the mission to Neptune-Triton system. In a previous work, we presented the expansion of the disturbing function for the dynamics of this system. Here we derive the averaged classical equations of the precession of the equator for this problem. The highly inclined and retrograde orbit of Triton makes this problem very unusual. Therefore, the classical truncations in the inclinations are not acceptable, so that the precession equations must be obtained in closed form for the inclination. With a significant mass and due to its distance from Neptune, which is continuously decreasing, Triton should exert in the future, important precession on Neptune's equator. The effects of this precession on the inner satellites are shown, including some resonant cases predicted in the future. Although Triton's orbit is almost circular, no expansion in the eccentricity is needed.

Copyright © 2009 C. do Nascimento and T. Yokoyama. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction

Some years ago, the exploration of the Neptune system in the next ten years was posed as an important priority of NASA's Solar System Exploration theme [1]. Although the New Horizons project is now the top priority for the far reaches of the solar system, most probably, to design a mission to explore the Neptune-Triton system will be just a matter of time. The authors [1] raised some important questions about the dynamics (stability) of the arcs and inner satellites of Neptune. In order to give support to the mission, we must increase our knowledge on the dynamics of this system. In this work we derive the classical equations of the precession for the equator of Neptune. Basically, this equator is disturbed by the presence of Triton since it has a remarkable mass: $m_T/M_N \approx 2.09 \times 10^{-4}$. (m_T , M_N are




Figure 1: XYZ: inertial system, xyz: moving system fixed on the equator.

the masses of Triton and Neptune, resp.). Moreover, Triton's inclination with respect to Neptune's equator is quite unusual ($\approx 157.35^\circ$). In a short time span, this precession should not be so important on the dynamics of the inner satellites. However, for a longer time (more than 600 years), this effect must be investigated. This work is organized in the following way: in Section 2 the averaged precessional equations for Neptune's equator considering arbitrary inclination obtained. Some authors [2] hypothesize that in the past, the eccentricity of Triton (e_T), might have been very high. Therefore, the precessional equations are also obtained for arbitrary e_T . In Section 3, we discuss the definition of the invariable plane and the existence of a corresponding equilibrium solution when we assume circular orbit for Triton. Considering the current dynamics of Neptune-Triton system, several authors have improved the ephemerides and data of the small satellites (e.g., Owen et al. [3]). Here, some numerical simulations showing the effects of the precession, particularly for the future 2 : 1 and 3 : 1 resonances involving Triton and Proteus, are presented. Section 4 is devoted to the conclusions.

2. The $\tilde{I}$ and $\tilde{\Omega}$ equations

Let XYZ be an arbitrary reference system (inertial) and a moving system xyz fixed on the equator of Neptune (see Figure 1). We define $\tilde{I}$ = inclination of the equator and $\tilde{\Omega}$ = node of the equator [4]. Due to the presence of a massive satellite (Triton), the equator of Neptune will be disturbed. Let $\mathbf{T}$ be the associated torque due to Triton and let W_T be the corresponding oblateness potential of Neptune:

$$W_T = -\frac{GM_N m_T J_2 R_N^2}{r_T^3} \left(-\frac{1}{2} + \frac{3\sin^2 \beta}{2} \right), \quad (2.1)$$

where G is the gravitation constant, M_N and m_T are the masses of Neptune and Triton, respectively. The oblateness constant is J_2 , β is the Triton's latitude and r_T is the radial distance

Neptune-Triton in the xyz system. The equatorial radius of Neptune is indicated by R_N . Then:

$$\mathbf{T} = (L, M, N) = -\mathbf{r}_T \times \nabla_T W_T, \quad (2.2)$$

where ∇_T is the usual gradient operator with respect to Triton coordinates and L , M , and N are the components of the torque T , that is:

$$\begin{aligned} L &= \left(\frac{\partial W_T}{\partial y_T} \right) z_T - \left(\frac{\partial W_T}{\partial z_T} \right) y_T, \\ M &= -\left(\frac{\partial W_T}{\partial x_T} \right) z_T + \left(\frac{\partial W_T}{\partial z_T} \right) x_T, \\ N &= \left(\frac{\partial W_T}{\partial x_T} \right) y_T - \left(\frac{\partial W_T}{\partial y_T} \right) x_T, \end{aligned} \quad (2.3)$$

where x_T , y_T and z_T are the cartesian coordinates of Triton with respect to the moving system xyz . On account of the rotational symmetry, we can show that $N = 0$. To this end, since β is the latitude we have: $\sin(\beta) = z_T / r_T$. Then a direct calculation shows that W_T depends on Triton coordinates only through r_T and z_T / r_T , therefore:

$$\begin{aligned} W_T &= W_T \left(r_T, \frac{z_T}{r_T} \right), \quad r_T = (x_T^2 + y_T^2 + z_T^2)^{1/2}, \\ \frac{\partial W_T}{\partial x_T} &= \frac{\partial W_T}{\partial r_T} \frac{x_T}{r_T}, \quad \frac{\partial W_T}{\partial y_T} = \frac{\partial W_T}{\partial r_T} \frac{y_T}{r_T}. \end{aligned} \quad (2.4)$$

Substituting these last two relations in N given in (2.3), we get $N = 0$. Now, let $\mathbf{w} = (p_N, q_N, w_N)$ be the rotation vector of Neptune. According to Euler's dynamical equations:

$$\begin{aligned} \dot{p}_N + \frac{C-A}{A} w_N q_N &= \frac{L}{A}, \\ \dot{q}_N - \frac{C-A}{A} w_N p_N &= \frac{M}{A}, \\ \dot{w}_N &= \frac{N}{C} = 0, \end{aligned} \quad (2.5)$$

where $A (= B)$ and C are the moments of inertia of the planet. From the geometric equations, $\mathbf{w}$ can be expressed in terms of the inertial XYZ system:

$$\begin{aligned} p_N &= \sin(\tilde{I}) \sin(\chi) \ddot{\Omega} + \cos(\chi) \ddot{I}, \\ q_N &= \sin(\tilde{I}) \cos(\chi) \ddot{\Omega} - \sin(\chi) \ddot{I}, \\ w_N &= \cos(\tilde{I}) \ddot{\Omega} + \dot{\chi}. \end{aligned} \quad (2.6)$$

Taking the time derivative of p_N ,

$$\begin{aligned} \dot{p}_N = & \cos(\tilde{I})\dot{\tilde{I}}(\sin(\chi)\ddot{\tilde{\Omega}}) + \sin(\tilde{I})\left(\dot{\chi}\cos(\chi)\ddot{\tilde{\Omega}} + \sin(\chi)\ddot{\tilde{\Omega}}\right) \\ & - \sin(\chi)\dot{\chi}\dot{\tilde{I}} + \cos(\chi)\ddot{\tilde{I}}. \end{aligned} \quad (2.7)$$

Assuming that the polar component of the rotation of the planet is dominant, the following quantities: $\tilde{I}\ddot{\tilde{\Omega}}$, $\dot{\tilde{I}}\ddot{\tilde{\Omega}}$ e $\ddot{\tilde{\Omega}}^2$ can be neglected when compared to $\dot{\chi}$ [5, 6]. Then,

$$\dot{p}_N = \sin(\tilde{I})\dot{\chi}\cos(\chi)\ddot{\tilde{\Omega}} - \sin(\chi)\dot{\chi}\dot{\tilde{I}}. \quad (2.8)$$

Substituting $\dot{p}_N$ e q_N in the first Euler's equation,

$$\begin{aligned} \frac{L}{A} = & \left(\ddot{\tilde{\Omega}}\dot{\chi}\cos(\chi)\sin(\tilde{I}) - \dot{\tilde{I}}\dot{\chi}\sin(\chi)\right) \\ & + \frac{C}{A}w_N\left(\ddot{\tilde{\Omega}}\sin(\tilde{I})\cos(\chi) - \sin(\chi)\ddot{\tilde{I}}\right) \\ & - w_N\left(\ddot{\tilde{\Omega}}\sin(\tilde{I})\cos(\chi) - \sin(\chi)\ddot{\tilde{I}}\right). \end{aligned} \quad (2.9)$$

Substituting $w_N = \cos(\tilde{I})\ddot{\tilde{\Omega}} + \dot{\chi}$ and neglecting terms factored by $\ddot{\tilde{\Omega}}^2$ and $\dot{\tilde{I}}\ddot{\tilde{\Omega}}$, we have

$$\frac{C}{A}w_N\left(\sin(\tilde{I})\cos(\chi)\ddot{\tilde{\Omega}} - \sin(\chi)\ddot{\tilde{I}}\right) = \frac{L}{A}. \quad (2.10)$$

Similarly, putting $\dot{q}_N$ and p_N into the second Euler dynamical equation, we obtain:

$$-\frac{C}{A}w_N\left(\sin(\tilde{I})\sin(\chi)\ddot{\tilde{\Omega}} + \cos(\chi)\ddot{\tilde{I}}\right) = \frac{M}{A}. \quad (2.11)$$

Now we can eliminate L and M in the above equations. Defining $W_0 = -Gm_TM_NJ_2R_N^2/2$, we have

$$\begin{aligned} W_T = & W_0\left(\frac{3z_T^2}{r_T^5} - \frac{1}{r_T^3}\right), \\ \frac{\partial W_T}{\partial y_T} = & W_0\left[\frac{\partial}{\partial r_T}\left(\frac{3z_T^2}{r_T^5} - \frac{1}{r_T^3}\right)\right]\frac{y_T}{r_T}, \\ \frac{\partial W_T}{\partial z_T} = & W_0\left[\frac{\partial}{\partial r_T}\left(\frac{3z_T^2}{r_T^5} - \frac{1}{r_T^3}\right)\right]\frac{z_T}{r_T} + W_z, \end{aligned} \quad (2.12)$$

and so,

$$L = \frac{\partial W_T}{\partial y_T} z_T - \frac{\partial W_T}{\partial z_T} y_T = -W_z y_T, \quad (2.13)$$

where $W_z = 6W_0 z_T / r_T^5$. Similarly, $M = W_z x_T$. From (2.10) and (2.11),

$$\begin{aligned} \ddot{\tilde{\Omega}} \sin(\tilde{I}) &= -\frac{W_z}{Cw_N} (y_T \cos(\chi) + x_T \sin(\chi)), \\ \ddot{\tilde{I}} &= \frac{W_z}{Cw_N} (-x_T \cos(\chi) + y_T \sin(\chi)). \end{aligned} \quad (2.14)$$

If a sequence of three rotations, $\tilde{\Omega}$, $\tilde{I}$, is χ are applied on the system XYZ , it will coincide with xyz . Therefore we can write x_T , y_T , and z_T in terms of the well-known elliptic orbital elements of Triton.

$$\begin{aligned} x_T &= r_T \cos(f_T + w_T) \cos \Omega_T - r_T \sin(f_T + w_T) \sin \Omega_T \cos I_T, \\ y_T &= r_T \cos(f_T + w_T) \sin \Omega_T + r_T \sin(f_T + w_T) \cos \Omega_T \cos I_T, \\ z_T &= r_T \sin(f_T + w_T) \sin I_T, \end{aligned} \quad (2.15)$$

where r_T , f_T , w_T , Ω_T , w_T , and I_T refer to classical orbital elements of Triton referred to (X, Y, Z) inertial system.

Finally, averaging the resulting equations with respect to the mean anomaly of Triton, the precessional equations, without any restriction on the inclination I_T or on the eccentricity e_T , are

$$\begin{aligned} \ddot{\tilde{I}} &= \frac{F_0}{4a_T^3(1-e_T^2)^{3/2}} \left[\sin(\tilde{I}) \sin(I_T)^2 \sin(2\tilde{\Omega} - 2\Omega_T) + \cos(\tilde{I}) \sin(2I_T) \sin(\tilde{\Omega} - \Omega_T) \right], \\ \ddot{\tilde{\Omega}} &= \frac{F_0}{4a_T^3(1-e_T^2)^{3/2}} \left[\left(1 - 3\cos(I_T)^2 \right) \cos(\tilde{I}) + \frac{2\sin(I_T) \cos(I_T) \cos(2\tilde{I}) \cos(\tilde{\Omega} - \Omega_T)}{\sin(\tilde{I})} \right. \\ &\quad \left. + \sin(I_T)^2 \cos(\tilde{I}) \cos(2\tilde{\Omega} - 2\Omega_T) \right], \end{aligned} \quad (2.16)$$

where

$$F_0 = \frac{3Gm_T M_N R_N^2 J_2}{Cw_N}. \quad (2.17)$$

Notice that, as expected in this approximation, these two equations are decoupled from χ equation.




Figure 2: $\mathbf{t}_1$ is perpendicular to the orbital plane of Triton, $\mathbf{n}_1$ is perpendicular to Neptune's equator, and $\mathbf{R}$ is the resulting vector which defines the invariable plane.

3. The Invariable Plane

In this section, we define the reference plane. Let $\mathbf{t}_1$ and $\mathbf{n}_1$ be the orbital angular momentum of Triton and rotation angular momentum of Neptune, respectively. Then $\mathbf{R} = \mathbf{t}_1 + \mathbf{n}_1$ is a constant vector. Therefore, the plane perpendicular to the direction of $\mathbf{R}$ is invariable (see Figure 2), and it will be adopted as our reference plane.

The dynamics of the system involving $\tilde{I}$, $\tilde{\Omega}$, Ω_T , and I_T shows an interesting solution in this invariable plane. To see that, let us assume Triton in a circular Keplerian orbit. In this case, we have $t_1 = m_T n_T a_T^2$, where n_T is the mean motion, and m_T is the Triton's mass. On the other hand, $n_1 = C w_N$, where C is the polar moment of inertia of Neptune and w_N is its angular velocity of rotation (spin rate). Now, from Figure 2, we have

$$\sin(\pi - I_T) = \frac{S}{m_T n_T a_T^2}, \quad \sin \tilde{I} = S C w_N \quad (3.1)$$

that is, the inclination $\tilde{I}$ of the invariable plane is given by:

$$\sin \tilde{I} = \frac{m_T n_T a_T^2 \sin I_T}{C w_N}. \quad (3.2)$$

In order to have the complete set of equations that govern the motion of the equator, we have to consider (2.16) and also the corresponding equations for I_T and Ω_T . They are

$$\begin{aligned} \frac{dI_T}{dt} &= -\frac{1}{n_T a_T^2 \sin(I_T)} \frac{\partial R_{J_2}}{\partial \Omega_T}, \\ \frac{d\Omega_T}{dt} &= \frac{1}{n_T a_T^2 \sin(I_T)} \frac{\partial R_{J_2}}{\partial I_T}, \end{aligned} \quad (3.3)$$

where R_{J_2} is the disturbing function due to the oblateness of Neptune which is written with respect to the following invariable plane:

$$R_{J_2} = S_0 \left[\frac{1}{2} \left(1 - \frac{3}{2} \sin^2 \tilde{I} \right) \left(1 - \frac{3}{2} \sin^2 I_T \right) + \frac{3}{8} \sin(2\tilde{I}) \sin(2I_T) \cos(\Omega_T - \tilde{\Omega}) + \frac{3}{8} \sin^2 \tilde{I} \sin^2 I_T \cos(2\Omega_T - 2\tilde{\Omega}) \right], \quad (3.4)$$

where

$$S_0 = \frac{GM_N R_N^2 J_2}{[a_T^3 (1 - e_T^2)^{3/2}]}. \quad (3.5)$$

For $\tilde{\Omega} - \Omega_T = \pi$, we easily see that

$$\begin{aligned} \dot{I}_T &= \tilde{I} = 0, \\ \dot{\tilde{\Omega}} &= -\frac{S_2}{\sin(\tilde{I})} \sin(2\tilde{I} + 2I_T), \\ \dot{\Omega}_T &= -\frac{3}{4} \frac{GM_N R_N^2 J_2}{n_T a_T^5 \sin I_T} \sin(2\tilde{I} + 2I_T), \end{aligned} \quad (3.6)$$

where

$$S_2 = \frac{3Gm_T M_N R_N^2 J_2}{4Cw_N a_T^3}. \quad (3.7)$$

Therefore,

$$\dot{\tilde{\Omega}} - \dot{\Omega}_T = \frac{3GM_N R_N^2 J_2}{4a_T^3} \sin(2\tilde{I} + 2I_T) \left[\frac{1}{n_T a_T^2 \sin I_T} - \frac{m_T}{Cw_N \sin(\tilde{I})} \right], \quad (3.8)$$

that is, the term inside the brackets is automatically zero provided the reference plane is the invariable plane (see (3.2)). In this case; $\tilde{\Omega} - \Omega_T = 0$, I_T and $\tilde{I}$ are constants; $\tilde{\Omega}$ and Ω_T are synchronous. This means that the precession rate of the Neptune's equator is the same of the Triton's node on the invariable plane (≈ 688 years). It is also easy to show that $\tilde{\Omega} - \Omega_T = \pi$ defines a stable periodic solution of the system (simple calculations show that such solution does not exist for $\tilde{\Omega} - \Omega_T = 0$). This means that it is very convenient to choose the invariable plane as the reference for our purposes.




Figure 3: (a), (c): the lighter points were obtained taking into account the $\tilde{I}$ and $\tilde{\Omega}$ equations in the integration, while in the thick points these contributions were neglected. (b), (d): time variation of the eccentricity (b) and inclination (d) for two orbits. Initially their paths are similar, but when their semimajor axes enter the 2 : 1 resonance region, a jump appears in the eccentricity (b), first in the orbit which contains the effect of the precession. The same occurs for the inclination (d), that is, the precession accelerates the onset of the jump of the inclination.

4. Some Effects on the Inner Satellites

Initially we tested our (2.16) for the case of Earth and Mars. As expected, the results agree quite well with very known values (≈ 26000 and ≈ 175000 years for Earth and Mars, resp.). For the Neptune's equator, considering $a_T = 14.063R_N$, our numerical experiments show ≈ 688 years for the period of precession, referred to the invariable plane Neptune-Triton. This period is rather sensitive for the adopted value of the semimajor axis of Triton. In this section we test the effect of the precession of Neptune's equator on an inner satellite of this planet. In Figure 3, we always integrate two orbits. The lighter points in the left panels, correspond to the orbits when we include the precessional effect, while the thick points come from the orbit

when we neglect (2.16). The inclinations are referred to the invariable plane. We consider a test particle (Proteus) disturbed by Triton.

In the integration of the equations of the motion, we used our averaged disturbing function R as described Yokoyama et al. [7] which is very efficient for this case. The initial conditions for Proteus are $a = 4.751$, $e = 0.0005$, $w = 0^\circ$, $\Omega = 90^\circ$, $I = 1^\circ$ Figure 3(a) and $I = 10^\circ$ Figure 3(c), and $\lambda = 90^\circ$, while for Triton we consider $a = 14.325$, $e = 0.000016$, $w = 0^\circ$, $\Omega = 270^\circ$, $I = 157^\circ$, and $\lambda = 270^\circ$. The effect of the precession is very clear: Figure 3(a), shows the variation of Proteus' inclination which remains confined in a very narrow interval when precession equations are considered. In Figure 3(c), in opposition to the above panel, we take Proteus at high inclination (10°). Then, although the periods are different, the inclinations of both orbits undergo a large variation. Moreover, there is quite clear a periodic oscillation of about 670 years, which is related to the precession of the node (Ω_T). Panels on the right, refer to the variation of the eccentricity Figure 3(b) and inclination Figure 3(d) of Proteus in the case of future 2 : 1 resonance. As described by Yokoyama [8] and Yokoyama et al. [7], due to the tides, Triton is falling down to the planet, so that, its orbital period will be in a 3 : 1 or 2 : 1 orbital resonance with Proteus in the future. For these two panels, instead of the averaged equations, we integrated the exact equations and we applied a slow dissipation on the semimajor axis of Triton in order to simulate the tidal effect. As pointed in Yokoyama et al. [7] the eccentricity suffers a strong variation and Proteus can be ejected. The effect of the precession is to anticipate the ejection. However, this is not a general rule, since some tests performed in the 3 : 1 resonance showed different results, that is, the precession can delay the ejection. More tests must be carried on in order to clarify these details.

5. Conclusion

In this work, we derived the classical averaged equations for the precession of the Neptune's equator, extending for the case of arbitrary inclination and eccentricity of the disturber. Due to the high retrograde inclination of Triton, these equations must be obtained in closed form for this inclination. If the invariable plane is chosen as the reference plane, the averaged equations of the precession in the case of $e_T = 0$, satisfy an equilibrium solution where $\tilde{\Omega}$ and Ω_T remain synchronous while $\tilde{I}$ and I_T are constants. The numerical experiments show that, when the precessional equations are considered, the inclination of an inner satellite of Neptune (Proteus) remains confined in a very narrow range, that is, the variation of the inclination with respect to the invariable plane is very small. However for high initial inclination ($\geq 10^\circ$) of the inner satellite, this confinement is not preserved anymore, that is, even considering the precessional effect, the amplitude of the variation of the inclination is significant. Additional simulations show that the precessional effects are also important in the future 2 : 1 and 3 : 1 resonances, since the jump in inclination or eccentricity can be anticipated or delayed.

Acknowledgments

The authors thank FAPESP, CNPQ, and FUNDUNESP.

References

- [1] H. B. Hammel, K. H. Baines, J. D. Cuzzi, et al., "Exploration of the Neptune system," in *The Future of Solar System Exploration, 2003–2013: Community Contributions to the NRC Solar System Exploration Decadal Survey*, M. V. Sykes, Ed., vol. 272 of *ASP Conference Series*, pp. 297–323, 2002.
- [2] P. Goldreich, N. Murray, P. Y. Longaretti, and D. Banfield, "Neptune's story," *Science*, vol. 245, pp. 500–504, 1989.
- [3] W. M. Owen Jr., R. M. Vaughan, and S. P. Synnott, "Orbits of the six new satellites of Neptune," *Astronomical Journal*, vol. 101, no. 4, pp. 1511–1515, 1991.
- [4] E. K. Woolard, *Astronomical Papers of American Ephemeris*, vol. 15, part 1, 1953.
- [5] R. Kurth, *Introduction to the Mechanics of the Solar System*, Pergamon Press, London, UK, 1959.
- [6] D. Lazzaro, "Semi-Analytical Theory for the Motion of Neptune's Satellites," Doctor thesis, Universidade de São Paulo, São Paulo, Brazil, 1987.
- [7] T. Yokoyama, C. Nascimento, and M. T. Santos, "Inner satellites of Neptune—I: the mutual disturbing function," *Advances in Space Research*, vol. 36, pp. 569–577, 2005.
- [8] T. Yokoyama, "Possible effects of secular resonances in Phobos and Triton," *Planetary and Space Science*, vol. 50, no. 1, pp. 63–77, 2002.

