

Molecular dynamics simulations illuminate the role of sequence context in the ELF3-PrD-based temperature sensing mechanism in plants

Richard J. Lindsay^{1,2}, Rafael Giordano Viegas^{3,4}, Vitor B. P. Leite³, Philip A. Wigge^{5,6}, Sonya M. Hanson^{1,2*}

*For correspondence:

shanson@flatironinstitute.org (SMH)

¹Center for Computational Biology, Flatiron Institute, New York, NY 10010, USA; ²Center for Computational Mathematics, Flatiron Institute, New York, NY 10010, USA; ³Department of Physics, São Paulo State University (UNESP), Institute of Biosciences, Humanities and Exact Sciences, São José do Rio Preto, SP, 15054-000, Brazil; ⁴Federal Institute of Education, Science and Technology of São Paulo (IFSP), Catanduva, São Paulo 15.808-305, Brazil; ⁵Leibniz-Institut für Gemüse-und Zierpflanzenbau, Großbeeren, Germany; ⁶Institute of Biochemistry and Biology, University of Potsdam, Potsdam, Germany

Abstract

The evening complex (EC) is a tripartite DNA repressor and a core component of the circadian clock that provides a mechanism for temperature-responsive growth and development of many plants. ELF3, a component of the EC, is a disordered scaffolding protein that blocks transcription of growth genes at low temperature. At increased temperature EC DNA binding is disrupted and ELF3 is sequestered in a reversible nuclear condensate, allowing transcription and growth to proceed. The condensation is driven by a low complexity prion-like domain (PrD), and the sensitivity of the temperature response is modulated by the length of a variable polyQ tract, with a longer polyQ tract corresponding to enhanced condensate formation and hypocotyl growth at increased temperature. Here, a series of computational studies provides evidence that polyQ tracts promote formation of temperature-sensitive helices in flanking residues with potential impacts for EC stability under increasing temperature. REST2 simulations uncover a heat-induced population of condensation-prone conformations that results from the exposure of 'sticky' aromatic residues by temperature-responsive breaking of long-range contacts. Coarse-grained Martini simulations reveal both polyQ tract length and sequence context modulate the temperature dependence of cluster formation. Understanding the molecular mechanism underlying the ELF3-PrD temperature response in plants has implications for technologies including modular temperature-response elements for heat-responsive protein design and agricultural advances to enable optimization of crop yields and allow plants to thrive in increasingly inhospitable environments.

Introduction

Temperature sensing mechanisms are critical to the survival of all life, but for plants, which are mostly stationary and yet flourish in a vast array of climates and seasonal variations, temperature sensors have a key, yet still poorly understood, role. While plants have diverse schemes to survive and thrive at a broad range of temperatures, such as vernalization, heat stress, and cold stress (Kerbler and Wigge, 2023), one mechanism that is more specific to plants is thermomorphogenesis, or when plants adjust their morphology in response to temperature (Quint et al., 2016).

Several thermosensory biomolecules have been identified, like the temperature-responsive uncoiling of chromatin, bending of DNA to increase gene transcription, and the melting of hairpin loops in RNA to enhance protein expression (Paik and Huq, 2019; Lin et al., 2020; Sengupta and Garity, 2013). One specific component known to be involved in thermomorphogenesis is the Evening Complex (EC), whose function depends on protein-protein and protein-DNA interactions, and has been well-studied in *Arabidopsis thaliana*. The EC is conserved in land plants (Liu et al., 2001) and is a tripartite protein complex that provides a tunable temperature-sensing mechanism with implications for growth rate, flowering, and potentially senescence and circadian rhythm (Zagotta et al., 1992; Anwer et al., 2014; Nusinow et al., 2011; Thines and Harmon, 2010; Box et al., 2015; Jung et al., 2020). Understanding the molecular basis for the temperature sensing mechanism of the Evening Complex has implications on a warming planet where even a 1°C rise in temperature would affect crop yield (Zhao et al., 2017).

The EC is a transcription repressor with three known protein components – ELF3, ELF4, and LUX ARRythmo – where LUX is a transcription factor (Nusinow et al., 2011; Zhang et al., 2019), ELF4 is a small helical protein responsible for stabilizing the complex, and ELF3 is a large mostly disordered protein that sequesters into reversible nuclear condensates at high temperatures (Fig. 1A) resulting in the release of the rest of the repressor complex, allowing the expression of growth genes (Jung et al., 2020). While ELF3 is a 695 residue long intrinsically disordered protein (IDP) the C-terminal prion-like domain (PrD), first recognized by Jung et al. (2020), is necessary and sufficient for condensate formation (Fig. 1C) and makes up a quarter of the ELF3 sequence (residues 492 to 664 out of 695 total residues) (Jung et al., 2020). Prion-like domains are identified by matching sequence patterns found initially in amyloid-forming prions, namely via an enrichment in polar amino acids like glutamine and asparagine (Lancaster et al., 2014), but this has proved a successful way of identifying low-complexity domains that are indicative of disordered protein regions (Romero et al., 2001). The PrD in ELF3 (hereafter referred to as the ELF3-PrD) is critically involved in the temperature sensitivity of the EC, as the ELF3 of the Mediterranean grass species *Brachypodium distachyon* has no PrD and does not undergo phase change or accelerated growth at higher temperatures (Jung et al., 2020).

A central feature of ELF3-PrD in *Arabidopsis thaliana* is the presence of a poly-glutamine (polyQ) tract of variable length found to differ in *Arabidopsis* populations by geographic location (Anwer et al., 2014). A similar polyQ tract expansion is responsible for the formation of pathogenic huntingtin protein aggregates in Huntington's disease patients. However the tract lengths observed in Huntington's disease are longer (>35 residues) than those of ELF3, which contains polyQ tracts of 0 to 29 glutamine residues. (Reiner et al., 2011). Experiments by Jung et al. found the length of this polyQ tract modulates the sensitivity of the temperature response, observing that longer polyQ enhances condensate formation and hypocotyl growth rate at high temperature (Jung et al., 2020). The mechanistic insights gained from studies of ELF3 condensate formation and temperature responsiveness could shed light on general principles governing polyQ-mediated interactions in native cellular contexts.

That evolution has co-opted biological condensates for temperature sensing in *Arabidopsis* aligns with the intrinsic temperature-dependence of condensate formation as well as the nature of IDPs as key sensors of cellular physicochemistry in general (Moses et al., 2023). However understanding the driving forces for IDP condensate formation at high (LCST) vs. low (UCST) tempera-

tures is still just beginning to be understood. Here, condensate formation at higher temperatures, like ELF3-PrD, correspond to those with a lower critical solution temperature (LCST) and those that form condensates at lower temperatures have an upper critical solution temperature (UCST) – these abbreviations will be used from now on. The understanding of LCST vs. UCST behavior has improved recently, for example via sequence heuristics proposed in (Quiroz and Chilkoti, 2015; Ruff et al., 2018), but the understanding physicochemical drivers of native LCSTs is still a work in progress. Sequence heuristics in general are a useful way to understand IDP functions (Maristany et al., 2023), for example they can accurately identify classes of IDP by simply looking at fraction of positive and negative charges in the sequence (Fig. 1D) (Das and Pappu, 2013). However, understanding the interplay between the polyQ regions of ELF3-PrD and the other sequence heuristics available is non-trivial. Previous studies have shown that polyQ tracts often enhance helical propensity in adjacent regions (Totzeck et al., 2017), which is of particular interest because short regions of transient helicity called Short Linear Motifs (SLiMs) are often responsible for formation of intra- and inter-protein interactions necessary for biological function (Davey et al., 2012).

The dynamic nature of IDPs has historically complicated their structural characterization (Bari and Prakashchand, 2021; Schramm et al., 2019). Experimental approaches can provide information about ensemble averages, but elucidating information about specific states or clusters of states remains a challenge (Nag et al., 2022). Using computational approaches to bridge this gap and access conformational ensembles, the molecular characterization of condensates is becoming increasingly accessible. While atomistic molecular dynamics (MD) simulations offer the most precise access to these ensembles, the computational cost of sampling the full conformational space of an IDP, not to mention a condensate, is steep. Recent methods for sampling IDPs by decomposing them into their component peptides and sampling these smaller systems individually before stitching them back together has been a fruitful approach in this space (Stelzl et al., 2022; Pietrek et al., 2020; Lindsay et al., 2021). However, this approach is limited, for example, in that it does not explicitly sample long distance interactions. The development of specific forcefields to accurately sample disordered proteins (Best et al., 2014; Song et al., 2017; Robustelli et al., 2018; Wu et al., 2018; Abascal and Vega, 2005) and enhanced sampling methods, like replica exchange with solute tempering (REST2) (Wang et al., 2011a), have also improved the tractability of atomistic MD simulations of IDPs. However, efficiently computing properties of IDPs, especially on the condensate level, necessitates the use of coarse-grained models, either at the Martini level (Thomassen et al., 2022) or even more coarse grained at the single-bead-per-residue level (Regy et al., 2021; Tesei et al., 2021). Further levels of coarse graining have pushed the possible timescales and system sizes even further with lattice-models (Choi et al., 2019) and even machine-learning algorithms like ALBATROSS and CALVADOS (Lotthammer et al., 2024; Tesei and Lindorff-Larsen, 2022; von Bulow et al., 2024).

In this work, we employ computational techniques to understand how temperature impacts the structural character of the ELF3-PrD ensemble, how polyQ length and sequence context impacts these characteristics, and the impact of each of these factors on dynamics underlying ELF3 condensation. Using a fast and efficient chain-growth algorithm, we produce ensembles of ELF3-PrD at a variety of polyQ length and temperature conditions (Fig. 1B,1E). We discover temperature-responsive helices in polyQ-adjacent regions and identify a residue, F527, involved in the temperature-sensing mechanism of the PrD. Next, we employ all-atom replica exchange with solute tempering (REST2) simulations to characterize temperature-sensitive conformation changes that promote condensate formation (Fig. 1F). In addition to the wildtype ELF3-PrD, we examine the F527A mutant to characterize the key role played by the F527 residue in the temperature-response mechanism and a 0Q mutant with the variable polyQ tract removed to better understand the role of this polyQ tract in modulating the structural ensemble with increasing temperature. Finally, we perform coarse-grained Martini simulations to more directly study how ELF3-PrD condensation dynamics depend on temperature and polyQ length. Simulating 100 ELF3-PrD monomers in a small box allows us to investigate specific residues that play key roles in driving condensation, includ-

139 ing a major role observed for aromatic residues in driving protein-protein interactions underlying
140 condensate formation, while also revealing the impact of sequence context and the expansion or
141 deletion of the variable polyQ tract on the temperature range and sensitivity of the temperature-
142 responsive phase transition (Fig. 1G). Through this computational lens we are able to better under-
143 stand the role and molecular mechanism of polyQ-adjacent helices and aromatic residues in the
144 temperature-sensitive growth response of *Arabidopsis thaliana* mediated by the ELF3-PrD.




Figure 1. An overview of the Evening Complex (EC), key features of ELF3, and the computational approaches taken to elucidate the molecular mechanisms of temperature sensing of the ELF3-PrD. A.

An illustration of the temperature-responsive mechanism of the evening complex. The left panel shows the intact EC repressing a growth gene target. Second to left illustrates the complex dissociating from DNA at high temperature. The next section describes the prion-like domain required for condensation and the poly-glutamine region responsible for tuning the sensitivity of the temperature response. The right-most cartoon illustrates the enhanced growth response characteristic of expanded poly-glutamine tracts. **B.** Helical propensity of hierarchical chain growth (HCG) ensembles by residue at four temperatures. The 7Q (wild-type) system is shown in the main panel with a region of interest of 7Q and 0Q shown in the inset. The light blue regions mark the locations of the poly-glutamine tracts and the yellow region highlights an aromatic helix of mechanistic importance. **C.** A structure of wild-type ELF3-PrD taken from a REST2 simulation. **D.** A charge fraction plot commonly used to classify IDPs based on sequence. The red dot indicates ELF3-PrD falls in region 1, classifying it as a weak polyampholite characterized by a globular or 'tadpole-like' shape. **E.** An illustration of the hierarchical chain growth method used to produce ELF3-PrD ensembles at a variety of polyQ lengths and temperature conditions. **F.** A cartoon representation of the REST2 enhanced sampling simulation method. **G.** A snapshot from a coarse-grained Martini simulation used here to investigate temperature-dependent condensation propensity of ELF3-PrD variants.

Results

Sequence-based insights on ELF3-PrD properties

Sequence heuristics were already initially useful to identify the existence of the ELF3-PrD (Jun et al., 2020; Lancaster et al., 2014), and further investigation of other heuristics can provide insight into its structural and dynamical properties. One simple first step is to look at the ratio of positive to negative charges in any IDP, which in the case of ELF3-PrD suggests a globular or ‘tadpole-like’ shape (Fig. 1D) (Das and Pappu, 2013). Another simple sequence-level heuristic is the Ω_{aro} , which indicates ‘patchiness’ of the aromatic residue distribution. For ELF3-PrD this value is high at 0.897, indicating the aromatics are distributed in clusters rather than evenly through the sequence (see Fig. S7 for a visualization of the aromatics within the ELF3-PrD sequence), which is in contrast to other well-known PrD’s like FUS that have a bias toward uniformly distributed aromatics (Martin et al., 2020). A further, sequence-based view is that of (Quiroz and Chilkoti, 2015), who demonstrated that repeats of Pro- X_n -Gly were sufficient to provide a generic disordered scaffold that can be tuned to exhibit UCST or LCST characteristic phase separation in designed polymers. Inclusion of polar residues and zwitterionic pairs of charged residues in the X position or adjacent to the Pro- X_n -Gly motifs, where n is ≤ 4 , resulted in UCST character while LCST behavior results from the inclusion of nonpolar residues or lysine at these positions (see Fig. S7 for these motifs highlighted within the ELF3-PrD sequence). The X_n component of these motifs is split between non-polar and polar residues with those proximal to aromatic residues being mostly non-polar, suggesting these regions may promote LCST phase separation.

Two sequence-based web utilities to quickly probe the structural features of IDPs are ALBATROSS and CALVADOS. ALBATROSS (Lotthammer et al., 2024) is a machine learning-based approach trained on coarse-grained simulations of synthetic disordered IDPs using Mpipi (Joseph et al., 2021b), a coarse-grained force field which emphasizes π - π interactions like those dominant in the dynamics of low-complexity domains (Vernon et al., 2018). CALVADOS (Tesei and Lindorff-Larsen, 2022) predicts structural properties based on an input sequence. However, CALVADOS does not utilize a deep learning algorithm, but instead runs a coarse-grained simulation in real time using a single bead-per-residue approach in which hydropathy scales inform amino acid interaction strengths. The Flory scaling exponent is calculated by ALBATROSS to be between 0.36 and 0.394 for each variant, slightly above the theoretical value for a collapsed globule, 0.33. This is in agreement with the globular or ‘tadpole-like’ shape suggested by an earlier heuristic. The radius of gyration (R_g) reported by ALBATROSS is 3.259 nm for 0Q, 7Q, and 19Q, with only the F527A mutant exhibiting a different value of 3.31, suggesting an insensitivity of ALBATROSS to variations in sequence. More structural predictions obtained from ALBATROSS for ELF3-PrD can be found in Table S1. Unlike ALBATROSS, CALVADOS was able to give estimates for values of interest for IDPs at different temperatures, however it estimated the WT Flory scaling exponent, ν , as 0.49, more suggestive of an ideal polymer with little self-interaction and a more extended shape, which is not in line with other methods. CALVADOS predicts a slightly larger R_g value for the WT than ALBATROSS, with the 290K R_g predicted as 3.41 nm. The WT ensemble is predicted to become more expanded at 340K with an R_g of 3.73, as is the 13Q variant, shifting from 3.38 nm to 3.73 nm, the 19Q variant, moving from 3.7 nm to 3.8 nm, and the F527A mutant, expanding from an R_g of 3.31 nm to 3.89 nm. Only the 0Q system is not predicted to undergo significant expansion upon heating from 290K to 340K. A comparison of ELF3-PrD R_g values estimated with six different methods can be found in Fig. S8. A complete accounting of CALVADOS predictions for ELF3-PrD can be found in Table S2. While these quick approaches provide initial insights from just the amino acid sequence of any IDP, we found they sacrificed accuracy and precision in their ability to probe the range of environmental conditions relevant to characterize the effect of temperature and polyQ tract length on the conformational landscape of ELF3-PrD.

Chain growth ensembles exhibit helicity in PolyQ-flanking residues

In order to understand how polyQ length and temperature change the behavior of ELF3-PrD, we examined the structural differences between ensembles with varying polyQ tract lengths constructed at different representative temperatures. Using the hierarchical chain growth (HCG) method by (Pietrek *et al.*, 2020), we constructed ensembles of polyQ of 0, 7, 13 and 19 glutamine residues in length each at temperatures of 290K, 300K, 320K and 415K to produce a total of 16 ensembles with 32,000 conformations each, a total of 512,000 structures. For each residue we summed the conformers with α , 3_{10} or π -helix character to obtain a total helical propensity (Fig. 1B). We found ELF3-PrD to be composed of short transient helices, with most having helical propensity of only around 5%. Though most of the ELF3-PrD exhibits only light helical character, the helices adjacent to the variable-length polyQ tract had significantly higher propensities, up to 30% for the N-terminal residues and 10% for the C-terminal residues. These regions, which we refer to as putative SLiMs, display some helicity with or without polyQ as seen in the 0Q system, but the polyQ tract seems to increase the portion of the ensemble with helical character (Totzeck *et al.*, 2017). An additional feature of the polyQ-adjacent SLiMs is a degree of temperature-sensitivity not seen elsewhere in the PrD. In the 7Q system, the N-terminal polyQ adjacent SLiM falls from 30% at 290K to 20% at 300K and finally 10% propensity at 415K, while the same region in the 0Q system shows no appreciable reduction in propensity until 415K. The increase in temperature-responsiveness of these SLiMs in 7Q could be a by-product of the higher possible helical propensity enabled by the polyQ tract. Ensembles with polyQ tracts of 13 and 19 residues showed nearly identical results to the 7Q system (Fig. S9). This could be due to a limitation of the 5-residue fragments simulated to build libraries for the HCG method. Overall, it is clear the presence of a polyQ tract has a significant influence on the protein's local structural character and temperature responsiveness.

E-PCA (Lindsay *et al.*, 2018) is a contact analysis method that isolates the top sources of conformational variance (or dynamics when used on MD trajectories) by identifying residue pairs that form and break in a concerted manner. We apply it here to our HCG-generated ELF3-PrD ensembles to identify regions of correlated contacts which may be of mechanistic importance. For each system, the biggest signal consistently comes from three SLiMs directly N-terminally adjacent to the polyQ tract, suggesting correlated contact formation or breaking of contacts in this region explains most of the conformational variance, at least partially due to helix formation. At 290K, PC1 of all polyQ-containing systems (Fig. S10A, top-left panel, blue) contains a bimodal signal where each peak corresponds to interactions of residues within a polyQ-adjacent SLiM, suggesting some level of communication between these regions. The dominant mode of the 0Q system, on the other hand, exhibits a single peak, indicating communication between these SLiMs plays a less significant role in the contact dynamics of this system. At the maximum temperature, 415K, the contact dynamics of all systems converge, regardless of polyQ presence or length, as evidenced by the nearly identical PC1 values at this temperature (Fig. S10A, top-right panel, red). This peak at the high-temperature condition indicates formation of contacts that comprise the third polyQ N-terminal SLiM. However, one prominent negative value is consistently observed indicating interaction between F527 and Y530 strongly opposes contact formation within this SLiM. The persistence of this signal across ensembles and its location in the region identified as the primary source of conformational variability suggests a potential role for this contact pair in the general mechanism of ELF3-PrD, warranting further study. Further analysis of these chain growth ensembles can be found in the Supplementary Information.

All-atom REST2 simulations reveal temperature-responsive conformation change in ELF3-PrD

While the static conformers generated by the HCG approach provided a relatively quick and resource-inexpensive look at the effects of varying the polyQ tract length and temperature on the local structure of the ELF3-PrD, we turned to a more intensive MD simulation approach to gain a more complete understanding of dynamics underlying ELF3 function. Based on trends seen in the initial

HCG results, we decided to run three separate all-atom MD simulations using the REST2 method, including the wild-type ELF3-PrD to obtain a better understanding of the general molecular mechanism underlying temperature-responsiveness, the F527A mutant to understand the contribution of F527 to condensate formation and temperature responsiveness, and the 0Q system to isolate the role of the variable polyQ tract in ELF3-PrD dynamics. As with the HCG ensembles, helical propensity was calculated for all three PrD variants at four separate effective temperatures, 290K, 300K, 320K, and 405K (Fig. 2A, B and C respectively). Measurements of helical propensity around the variable-length polyQ tract from these trajectories are similar in trend to the HCG ensemble but with significantly larger propensity values, in the range of 30-65% instead of the 10-30%, for the N-terminal adjacent helix (Fig. 2A). Additionally, some level of helical enhancement was observed adjacent to all three polyQ tracts of ELF3-PrD rather than only the first as in the HCG case (*Totzeck et al., 2017*). The second polyQ tract (WT residues 568 to 572) exhibits enhancement in propensity and temperature-responsiveness in its N-terminal SLiM, and the third and last polyQ tract (WT residues 581 to 586) sees minor enhancement in its C-terminal SLiM, increasing to 10% or about double the baseline propensity. This enhancement of helical propensity indicates long-range interactions absent in the HCG ensembles further promote transient structure in the polyQ-flanking residues. The concerted nature of the enhancement of helicity and temperature-responsiveness may be more than a coincidence as enhancement of potential helical propensity could increase the ability to sense temperatures by providing a greater range of signals with more differentiable outcomes. The helical enhancement of polyQ-adjacent regions demonstrates how structural outcomes depend on sequence context, as degree and location of enhancement vary among the three tracts. The impact of sequence is not just local, as is apparent from the structural analysis of the F527A system. Relative to the WT, the F527A mutation attenuates helices, including those around each polyQ tract, despite being 15+ residues from the nearest tract.

While the 0Q system is reported to be significantly less sensitive to changes in temperature, some temperature-responsiveness does remain intact in experiments. The analysis of helical propensity for this trajectory reveals drastic structural deviation due to the absence of the polyQ (Fig. 5A), pointing to fundamental differences in the temperature-sensing mechanism compared with WT. In the absence of the variable polyQ tract, a large helix appears between residues 554 and 566 (Fig. 2C). This helix exhibits impressive stability at 290K (77% helicity) and rapidly declines linearly as temperature increases, bottoming out at 35% at 405K. This 0Q-native helix includes the first (second in WT) polyQ tract with the bulk of the helix forming N-terminal to polyQ. Aside from formation of this new helix, a distal aromatic helix, discussed further in the next section, is abolished in the 0Q system and helices that were flanking each side of the removed variable polyQ are attenuated. This result indicates the 0Q system may respond to temperature change by a mechanism vastly different from that of the WT, and the location of this new helix is further evidence that polyQ tracts promote helix formation in adjacent residues. Regardless, the interplay of variables like tract length (*Totzeck et al., 2017*), sequence context (*Ramazzotti et al., 2012*), tract location, and proximity to other polyQ tracts could all affect protein character and warrant further study.




Figure 2. REST2 simulations reveal local and global conformation changes in response to temperature.

A. Helical propensity of WT REST2 simulation at a range of temperatures. The variable polyQ tract is shaded in blue and the two polyQ tracts of consistent length are shaded in purple. A legend denoting regions of interest is located at the top. **B.** Helical propensity of F527A REST2 simulation at a range of temperatures. The red line indicates the position of residue 527. **C.** Helical propensity of 0Q REST2 simulation at four temperatures. The gray shaded region indicates the removal of the variable polyQ tract. **D.** Top: Solvent-accessible surface area of WT and F527A systems taken from REST2 trajectories. **E.** The radius of gyration of the first 100 residues (black) and last 73 residues (grey) of ELF3-PrD. Top panel depicts the WT and the bottom F527A, both at 290K.

282 Aromatics play a major role in temperature-sensitive conformational change

283 The REST2 simulations painted a more complete picture of the ELF3-PrD dynamics and revealed
284 some mechanistically important features absent in the HCG ensembles, partly due to the inclusion
285 of long-range interactions. The most prominent was a 10-residue helix of heavy aromatic character
286 (residues 494 to 504, "H_{aro}") which was formed between 40-52% of the time in the REST2 WT simu-
287 lations (Fig. 2A) but only present at baseline levels in the HCG ensembles. The F527A mutant sees a
288 significant reduction in the size of this helix (Fig. 2B), suggesting long-range interactions with F527
289 may stabilize H_{aro}. Indeed, it seems this interaction may have consequences for the global confor-
290 mation of ELF3-PrD. In the WT, both the regions around F527 and H_{aro} are largely buried, as shown
291 by the low solvent-accessible surface area (SASA) in Fig. 2D, and their relative immobility as exhib-
292 ited by RMSF values shown in Fig. 2E. A methionine residue of H_{aro}, M504, promotes compaction of
293 this region through interaction of the sulfur atom with the ring of multiple local aromatic (tyrosine)
294 residues (Fig. S11), a common but often overlooked motif (Valley et al., 2012). Upon mutation
295 of F527 to alanine, both the F527 and H_{aro} regions become relatively solvent-exposed and flexi-
296 ble. This interaction influences the global shape of the protein by pushing the N-terminal region
297 towards a globular conformation while the C-terminal region remains relatively extended. This is
298 demonstrated by comparing the R_g of the first 100 and last 73 residues for the WT and mutant
299 systems, as in Fig. 2F. The first 100 residues of the WT (top panel) have a sharp peak indicating a
300 compactness not observed in the mutant (bottom panel) while the last 73 residues sample a wide
301 range of R_g values in each case, indicating a more expanded and flexible nature. The mean R_g of
302 the whole PrD for the WT 290K ensemble is 2.75 nm, significantly more compact than predicted

by heuristic models ALBATROSS and CALVADOS which reported values of 3.26 nm and 3.41 nm, respectively, as shown in Fig. S8.

Upon discovering the global structural influence of the long-range interaction between F527 and H_{aro} , we set out to determine its role in temperature-responsiveness. The minimum distances between atoms of F527 and H_{aro} were examined under different temperature conditions, and we found a linear increase in F527 and H_{aro} distance (and decrease in interaction) with increasing temperature, as demonstrated by the histograms in Fig. 3A. To get a better sense for the role of this temperature-responsive property in the conformational landscape of the protein, we plotted the F527- H_{aro} distance vs the radius of gyration in Fig. 3B. At 290K, the WT is largely homogeneous, mostly constrained to a single population of tightly interacting F527- H_{aro} and compact structures. However, with increasing temperature the ensemble shifts towards a second population that is slightly expanded with the F527- H_{aro} contacts broken. Representative structures of the low-temperature and emergent high-temperature populations are depicted in Fig. 3A on the left and right, respectively. The F527A mutant reinforces the importance of F527 in the WT conformational landscape as F527A samples a diverse range of populations in this space. However, the most persistent F527A population has significant overlap with the WT high temperature population, suggesting the F527A mutant may mimic the phenotype of WT ELF3-PrD at high temperatures. In Fig. 3C, we've employed the ELViM method (*Viegas et al., 2024*) to project the WT and F527A conformations onto a common space, where each conformer is a dot colored by the minimum distance between F527 and H_{aro} value. The projections on the left of Fig. 3C represent conformers from all four temperature conditions while those to the right represent specific temperature conditions. These projections further demonstrate the degree to which the F527A mutant alters the conformations sampled by ELF3-PrD and illustrates how increasing the temperature from 290K to 320K promotes access of the WT to new regions of conformational space. Additional visualizations and analysis of the ELF3-PrD conformation space can be found in the SI. Interestingly, in the 0Q system, the F527- H_{aro} interaction loses this temperature-responsiveness (Fig. S12) at all but the highest temperatures where we actually see an increase in the interaction propensity. This along with the 0Q-specific temperature-responsive helix suggest a different temperature-sensing mechanism in absence of the variable polyQ tract.

The temperature-dependent breaking of the F527/ H_{aro} interaction is particularly interesting due to its potential relevance to ELF3-PrD condensate formation. A significant increase in total SASA is observed in the high-temperature population with a major source being aromatic residues (Fig. S13). As previously mentioned, H_{aro} and F527 are regions of enhanced aromatic residue density, as seen from the dashed red lines in Fig. 3D. Breaking the interaction between F527/ H_{aro} frees up 5 aromatic residues from these regions (Y496, Y499, Y500, Y503 & F527) and as well as four aromatic residues located between F527 and H_{aro} which can become buried during their interaction. Indeed, our high-temperature population exhibits enhanced solvent accessible surface area for these residues and for the majority of the 18 aromatic amino acids present in the ELF3-PrD with those residues located between H_{aro} and F527 especially affected (Fig. 3D). Given the well-established role of aromatic residues in driving interactions in low complexity disordered protein regions (*Martin and Mittag, 2018a*), it is reasonable to expect an increase in condensate formation propensity when these residues are accessible by other ELF3-PrD monomers. Indeed, previous work has established a link between single-chain dimensions and phase-separation propensity as a function of aromatic character (*Dignon et al., 2018; Martin et al., 2020*).




Figure 3. Temperature-responsive breaking of aromatic interactions exposes sticky aromatic residues to the solvent. **A.** The minimum distance between residue F527 and H_{aro} is plotted by temperature from the WT REST2 simulation. Representative low and high temperature structures are inside of a dashed blue or red box, respectively. F527 is in red and H_{aro} is in yellow. **B.** Free energy landscapes of WT (left) and F527A (right) with the X-axis defined as the minimum distance between residue 527 and H_{aro} and the y-axis defined as the radius of gyration. Each ensemble is represented at 290K and 320K. The blue box is the most populated region of conformation space at low temperature and the red box demarcates the emergent high temperature cluster. **C.** ELViM plots illustrating the conformation space of the full ensemble (left), the 290K ensemble (middle) and the 320K ensemble (right) where the color of each dot represents the minimum distance between F527 and H_{aro} . Below are example clusters of conformers with the left representing a commonly populated area at low temperature and to the right a cluster only observed at increased temperature. **D.** Fold increase in SASA between the identified low- and high-temperature populations. Dotted red lines represent aromatic residues and the yellow-shaded region are residues of H_{aro} .

347 PolyQ solvation dynamics contribute to ELF3-PrD condensate forming propensity

348 A key question to understand temperature sensing by ELF3 is how expansion of the polyQ tract en-
 349 hances the phase-separation propensity of ELF3-PrD, thus increasing the temperature sensitivity
 350 of condensation (Jung et al., 2020). One potential driver is the increasing availability of glutamine
 351 on the protein surface with increasing tract length. Plotting the SASA values for our HCG ensembles
 352 (Fig. 4A) highlights all three polyQ tracts as regions of persistent high solvent accessibility. While the
 353 SASA value of polyQ tracts is only moderately higher per residue than the average, indicated by a
 354 dashed black line, it persists for the length of the tract, creating patches of high solvent accessibility
 355 not seen elsewhere in the PrD. SASA values for these HCG ensembles were temperature-invariant
 356 for each case. Similar to the effect of increasing temperature in our REST2 simulations, increas-
 357 ing polyQ length enhances the overall solvent accessibility of the protein. PolyQ tract expansion
 358 provides a direct and tunable means to enhance PrD solvent-accessibility.

359 Solvation dynamics serve as a determining factor in condensate formation behavior, in part by
 360 determining which regions remain accessible to the solvent and other proteins, and changes in
 361 temperature can alter these dynamics. To characterize how hydration differs across ELF3-PrD we
 362 obtained radial distribution functions (RDFs) of water oxygen atoms around ELF3-PrD as a whole
 363 and around each individual residue. Values for a per-residue solvation plot were obtained by using
 364 each residue as the reference point for an RDF calculation of water oxygen atoms and plotting the
 365 height of the first peak, representing the most tightly bound water molecules, referred to as the

first solvation shell (Fig. 4B). At 405K, many residues exhibit a decrease in local solvation, with the most prominent and continuous decreases seen in the three polyQ tracts. Further investigation of ELF3-PrD polyQ interactions at high temperature show a tradeoff wherein a decrease in polyQ-water hydrogen bonding is compensated by an increase in protein-protein hydrogen bonds, even though the number of polyQ-proximal water molecules increases (Fig. 4C). This is consistent with studies that have suggested water is a poor solvent for glutamine (*Khare et al., 2005*). This characteristic may influence condensation, either by promoting polyQ-protein interactions or altering the solubility of the protein.




Figure 4. Cluster formation and monomer conformation change are driven by similar aromatic residues. **A.** The solvent-accessible surface area is shown for each HCG polyQ condition at 290K. The variable polyQ tract is highlighted in sky blue and the second and third polyQ tracts in a lighter blue. The black dashed line demarcates the average SASA value for the system. **B.** Per-residue hydration plot of ELF3-PrD WT at 290K (blue) and 405K (red). Each value represents the maximum height of the first peak of an RDF between the given residue and solvent oxygen atoms. The aromatic helix, H_{aro} , is highlighted in yellow and the polyQ tracts in sky blue. **C.** A scatter plot relating change in hydrogen bonds between the variable polyQ tract and water with the change in hydrogen bonds between polyQ and protein from the wild-type REST2 simulation at 290K. Breaking of polyQ-water hydrogen bonds is correlated with an increase in polyQ-protein bonds.

PolyQ tract length modulates temperature-responsiveness of PrD clustering

We next shift our focus from the structural ensembles of ELF3-PrD monomers to the condensation dynamics of the system. Performing coarse-grained Martini simulations provides a more direct way to identify interactions which drive protein-protein interactions of the PrD and to observe the ways in which variables like polyQ tract length and sequence mutations modulate the formation and stability of clusters in response to temperature. We first examine the impact of temperature on contact dynamics of the 100 monomers that comprise our condensate simulations to see how they align with our all-atom simulations. The contact difference map seen in Fig. 5A provides an understanding of how intra-protein interactions change with increasing temperature. The blue

regions indicate interactions between an N-terminal region of the PrD with centrally located aromatics at 290K which are broken at 340K. Corresponding red lines in the central region indicate an enhancement of local interactions between aromatic residues within this region. This breaking of long-range contacts at increased temperature is similar to the temperature-induced conformational shift observed in the all-atom simulations. Here, F527 interacts in a temperature-dependent manner with F458 instead of the H_{aro} region (Fig. 5B), as was observed in the REST2 simulations. These Martini simulations do not capture transient helices, which may account for the lack of interaction with H_{aro} , but the trend is very similar nonetheless. By plotting the per-residue SASA of each of the 100 PrD monomers in Fig. 5C, a clear increase is seen across all residues, with the biggest peaks aligning with the aromatic residues. Based on a similar observation in the all-atom systems, we inferred that these residues would participate in inter-protein interactions, however, here we can observe these interactions directly. The inter-protein contact difference map in Fig. 5D indicate an increase in interaction between centrally-located aromatics at 340K as compared to 290K. An illustration of a WT PrD cluster forming at increased temperature is seen in Fig. 5E. The intra- and inter-protein difference maps are strikingly similar, suggesting the same interactions are responsible for monomer dynamics drive protein-protein interactions. We further explore the intra-protein interaction propensities of the aromatic residues in Fig. S14 which demonstrates the degree to which interactions between aromatic residues control the dynamics of this low complexity domain. The overall shape of the interaction profiles are nearly identical with only their scales varying, suggestive of the universal nature of aromatic contacts in these types of proteins. This plot provides some insight into the similarities observed in the intra- and inter-protein contact maps.

Our coarse-grained simulations allow us to measure the temperature dependence of ELF3-PrD clustering and how polyQ length and sequence mutations impact size and stability of clusters. Following the procedure used by previous protein condensate studies, we track the number of monomers comprising the largest cluster as a proxy for condensation propensity (*Benayad et al., 2021; Joseph et al., 2021a*). Cluster membership is determined using a cutoff of 0.5nm between backbone beads of individual monomers. The convergence of 100 ELF3-PrD monomers into a few large clusters is illustrated for each of these conditions at 290K in Fig. 6A with conformers taken from a wild type simulation at around the 0, 1 and 3 μ s. We take a closer look at the temperature dependence of cluster formation for the wild-type system in Fig. 6B. Here, there is a clear increase in maximum cluster size given a temperature increase from 290K to 300K. This represents a biologically relevant temperature range and is close to the temperatures used in experiments (*Jung et al., 2020*). Clustering occurs more slowly at 320K and only briefly reaches the size seen at 300K before the end of the 3 μ s simulation, but is still enhanced relative to 290K. At 340K clustering no longer occurs on a large scale with the largest clusters being around 10 monomers in size and transient in nature. We can gain insights into the driving force of temperature-dependent condensation by examining the net change in total contacts per residue of the 7Q system at 300K vs. 290K during the last microsecond of simulation, when the clusters are most fully formed (Fig. 6C). While an increase in contacts is observed for every residue at 300K, aromatic residues exhibit dramatic gains, clearly visible as large spikes in the net contact plot. As clusters start to form, the polyQ tract expels water molecules in exchange for increasing polyQ-protein interactions, as shown for the 7Q system at 300K in Fig. 6D.

Expansion of the polyQ tract to 19Q (Fig. S15) results in similar trends seen in the wild type at 290K at 300K, with enhancement of clustering observed at 300K resulting in clusters moderately larger than those exhibited by the wild type. At 320K, the 19Q system exhibits no enhancement of clustering compared with the 290K condition, revealing a difference between the 19Q and 7Q systems. Cluster formation is not observed at 340K, a trend seen in each ELF3-PrD variant simulated here. Interestingly, the variable polyQ tract of the 19Q system exhibits fewer protein-water interactions than the WT when normalized by tract length (Fig. S16). Removal of the variable polyQ tract, as in the 0Q system (Fig. S17), results in smaller clusters in the 290K condition compared with wild type. Enhancement is again observed at 300K, but now the 320K condition exhibits an increase




Figure 5. Temperature-dependent conformation change enhances SASA of CG monomers and promotes cluster formation driven by central aromatic residues. **A.** The intra-protein contact difference map between the 340K and 290K WT Martini condensate simulations. Values in red represent increases in contact value at 340K relative to 290K and blue represents a decrease. The entire 3 μ s trajectories were used to create these difference maps. **B.** Distance distributions between residue F527 and F458 at 290K (grey) and 340K (red) from WT CG condensate simulations. Distances are taken from intra-protein distances of each monomer. **C.** The solvent accessible surface area is plotted for all 100 monomers of both the 290K WT condensate simulation (black) and the 340K WT condensate simulation (red). **D.** The inter-protein contact difference map between the 340K and 290K WT Martini condensate simulations. As with Fig. 5A, contacts enhanced at high temperature are in red with those broken in blue. **E.** An illustration of the typical time evolution of an ELF3-PrD condensate simulation at 300K with simulation snapshots representing approximately 0 ns, 800 ns, and 3 μ s, respectively.

of cluster formation on par with that at 300K. These results indicate polyQ expansion enhances the condensation of ELF3-PrD in the biologically relevant temperature range of 290K to 300K and removal of the polyQ tract does not result in an inability to form clusters, but reduces the condensation at 290K while maintaining stability at a higher temperature of 320K when compared 7Q and 19Q.

In addition to polyQ length, we performed a set of simulations probing the dynamics of cluster formation of the F527A mutant in response to temperature (Fig. S18). As with the polyQ variants, an increase in maximum cluster size is observed at 300K as compared with 290K. However, a marked

decrease in cluster size is observed at 320K. This data suggests a more narrow range of effective clustering temperatures for the F527 mutant compared with the 7Q wild type system. These simulations suggest removal of one of the key drivers of protein-protein contacts, F527, narrows the range of temperatures at which clusters are stable and conducive to liquid-liquid phase separation. Due to its prominent role in temperature-responsive conformation change, we expected that F527A mutant would result in more expanded condensation-prone conformations, however, removal of F527 eliminates one of the key drivers of protein-protein interaction. This impedes to the ability of the F527A ELF3-PrD mutant to form stable condensates and reinforces the importance of sequence context in condensation of low-complexity domains.




Figure 6. Martini 3 simulations capture the temperature of condensate formation of ELF3-PrD. **A.** The number of ELF3-PrD clusters over time for the 7Q, 0Q, 19Q and F527A systems at 290K is illustrated with standard deviation. A cluster is defined using a 0.5nm cutoff between backbone beads. **B.** The number of ELF3-PrD monomers in the largest cluster from the wild-type simulations at four different temperatures. **C.** The number of water molecules interacting with the variable polyQ tract is demonstrated for 7Q at 300K in black. The concurrent number of protein interactions with the variable polyQ tract is shown in red. **D.** The net difference in protein contacts per residue at 300K vs 290K is plotted for the wild type ELF3PrD. Aromatic residues are denoted by dashed red lines.

Discussion

We have used varying levels of simulation accuracy to understand the molecular basis for the ELF3-PrD-based temperature response in plants. Together, these methods have enabled us to propose

mechanisms by which ELF3-PrD responds to temperature to promote protein-protein interaction and condensation. Initial sequence-based methods allowed us to put the ELF3-PrD in the context of other IDPs: a ‘tadpole’-like weak poly-ampholite (*Mao et al., 2010*) with some sequence characteristics in line with those seen for other LCST proteins (*Martin and Mittag, 2018b; Dignon et al., 2019*). Furthermore CALVADOS indicated some temperature dependent changes in radius of gyration, but CALVADOS and ALBATROSS gave different results for values like the Flory scaling exponent. However, we sought a more molecular-level understanding of the temperature-dependent properties of this process by building HCG ensembles at different temperatures and with different polyQ lengths to find that the variable polyQ was flanked by temperature-sensitive helices. These initial HCG results also indicated the importance of aromatic residues – specifically the F527. Extensive molecular simulations with REST2 allowed us to observe the temperature-dependent conformation landscape of the ELF3-PrD, including crucial long-range aromatic interactions not captured by HCG, like that of F527 and H_{aro} . Note that with an initial preprint the HCG and REST2 data of the atomistic ELF3-PrD simulations were made public at an early stage, and several groups analyzed these terabytes of data to test their methods, and found similar corroborating analyses to those presented here (*Viegas et al., 2024; González-Delgado et al., 2024*). Finally, coarse-grained Martini simulations showed that the expansion of the variable polyQ tract from 7Q to 19Q enhances condensate formation at the biologically relevant temperature of 300K, while the 0Q variant demonstrated smaller clusters at 290K and increased stability at 320K, potentially indicating an increase in the critical temperature required for phase separation. Additionally, the F527A mutant exhibited a decrease in cluster stability above 300K relative to the other variants studied, likely due to the removal of an important ‘sticker’ residue. These CG simulations further highlighted the importance of the aromatic residues for condensate formation, especially a central aromatic cluster with marked changes in contacts at different temperatures.

Overall, this molecular approach has helped understand the interplay of the varied sequence contexts of the ELF3-PrD and how they effect transient structures and temperature dependence. Here we found that aromatic residues, especially F527 and the central aromatic cluster, are key to condensate formation for this IDP, consistent with other analyses of condensate-forming IDPs (*Martini et al., 2020*). Furthermore, the transient temperature-dependent helices flanking the polyQ’s points to the importance of the sequence context around the polyQ’s. It is possible it is exactly this context that has reinforced these domains and the plasticity of polyQ motifs for adaptive temperature sensing, even when excessive polyQ length leads to disease (*Reiner et al., 2011*). These transient helices are reminiscent of SLiMs found in other IDP systems (*Davey et al., 2012*). While SLiMs have been seen to play a role in condensate formation, for example in TDP43, here we see a decrease in helicity as temperature increases, so this SLiM formation anti-correlates with condensate formation. The breakdown of these transient helices at high temperatures exposes aromatic residues to enhance condensate formation at higher temperatures (Fig. S22). Furthermore, a possible role for these putative SLiMs around the polyQ motifs is to enhance protein-protein and protein-DNA interaction with the rest of the EC complex at lower temperatures, when they are more stable.

Understanding the role of these transient helices in condensate formation from molecular simulation is complicated by the pros and cons of the various simulation methods used. The HCG and REST2 simulation both consistently found these polyQ-adjacent temperature-dependent helices, actually more prominent in the more accurate REST2 simulations (though we acknowledge reports of REST2 sampling overly compact states (*Zhang and Chen, 2022*)). These helices, however, are not well-captured in the coarse-grained simulations necessary to simulate condensate formation, as the only option is to explicitly enforce them with an elastic network model. However, we believe that the overall properties of these coarse-grained simulations are encouraging, as the contact maps and radius of gyration were overall very similar to those seen in the REST2 simulations. Furthermore, that Martini 3 was able to show increase in condensate-forming propensity at higher temperatures for the WT ELF3-PrD and an increase in condensate-forming propensity for a longer

polyQ tract is encouraging for the relevance of this force-field for studying LCST systems. For example, our attempts at using other single-residue-per-bead forcefields with implicit solvent like HPS (Dignon *et al.*, 2019) lead to much more compact conformations than we observed in atomistic simulation (Fig. S19), and especially noting the importance of protein-water interactions for LCST systems, we chose to not go this route.

The importance of understanding the ELF3-PrD temperature sensing mechanism in plants has increased since its original discovery (Jung *et al.*, 2020) as more similar systems have been described (Zhao *et al.*, 2017; Bohn *et al.*, 2024), and the threats of a warming planet on plant and crop health have become ever clearer. The results we present here help elucidate some of the molecular origins of this temperature sensitivity, however it is clear there is much work still to do. For example the simplified system we have studied here of the 173 residue ELF-PrD without the rest of the 695 residue protein and in the absence of the members of the EC (LUX, ELF4, and DNA) begs for further studies within a more realistic context, though ELF3-PrD condensate formation has been seen *in vitro*. Relatedly, both *in vitro* and *in vivo* experimental validation of the molecular mechanisms we postulate is an essential path forward. Overall, though, we believe that this study has taken the cutting edge of computational capabilities in the field of biomolecular condensate toward understanding a biologically important LCST.

Methods

Hierarchical Chain Growth Method and Simulation Details

The sequence for *Arabidopsis thaliana* ELF3 was obtained from UniProt. The PrD sequence (residues 432 to 604) was isolated and divided into 57 segments. Each segment was five amino acids in length including two residues that overlap with the previous segment and two residues that overlap with the subsequent segment. Each of the 57 segments was capped with an N-terminal acetyl group and a C-terminal N-methyl group to neutralize interactions of the charged ends of the protein. Segments were parametrized using the Amber03ws forcefield (Best *et al.*, 2014) and placed in a periodic rectangular box with 1nm of padding on each side of the protein and solvated with TIP4p2005 water molecules (Abascal and Vega, 2005).

Each segment was minimized until the maximum force reduction dropped below 1000 kJ/mol/nm per minimization step. For each of the 57 segments, 24 replicates were equilibrated to various temperatures ranging from 290K to 405K for 10ns. For this step, temperature effects were implemented using the v-rescale thermostat and the Berendsen barostat (Berendsen *et al.*, 1984). Finally, replica exchange MD was run for each segment for 100ns using GROMACS version 2021.1 (Abraham *et al.*, 2015). For this production run, the v-rescale thermostat was used to maintain temperature and the Parrinello-Rahman barostat (Parrinello and Rahman, 1980) was used to maintain a pressure of 1 bar.

Once the fragment simulations were finished, we used the hierarchical chain growth (HCG) application, developed by Pietrek *et al.*, to construct a 7Q ELF3-PrD ensemble from these 57 segments consisting of 32,000 unique conformations (Pietrek *et al.*, 2020; Stelzl *et al.*, 2022). Conformers were randomly chosen from each segment and joined in a pairwise fashion to create a set of fragment dimers. These were then joined into tetramers, octamers, and so on until the full 57-segment ELF3-PrD was reconstructed. This process was repeated until we had obtained ensembles with polyQ tracts of 0, 7, 13 and 19 residues in length each at temperatures of 290K, 300K, 320K and 415K for a total of 16 ensembles. Due to the two-residue overlap, we were required to run 19 segments made up of residues C-terminal to the polyQ region in order build the 0Q, 13Q and 19Q systems.

REST2 Simulation Details

The HCG method enabled us to quickly look at large ensembles of different polyQ conditions for potential structural differences and pointed us towards some regions and residues of interest. The

usefulness of the HCG method, for our purposes, is limited by the absence of long-range interactions. In order to accurately study the dynamics of ELF3-PrD we turn to REST2 (Replica Exchange with Solute Tempering), an all-atom enhanced sampling technique capable of exploring longer effective time scales than traditional all-atom MD ((*Jo and Jiang, 2015*)). Much like temperature replica exchange (T-REMD), REST2 runs multiple replicas in parallel, each at a different effective temperature. Every few hundred MD steps, replicas attempt to swap with one another with the probability of acceptance tied to the energy difference between replicates ((*Wang et al., 2011b*)). Unlike T-REMD where temperatures values are set for each replica, REST2 modulates the strength of protein-protein and protein-water interactions to achieve an effective temperature, a feature which reduces the number of replicas required compared with T-REMD. While REST2 can still be resource intensive, it is one of few methods able to provide Boltzmann-distributed ensembles of IDPs at biologically relevant timescales.

REST2 simulations ((*Wang et al., 2011b*)) were performed of the full-length wild-type ELF3-PrD, the ELF3-PrD F527A mutant and the 0Q ELF3-PrD with the variable polyQ tract removed. For the initial structure of these simulations, we chose an especially helical conformer from our HCG wild-type ensemble created at 290K. For the F527A mutant, F527 was manually edited to alanine by removing side-chain atoms and changing the residue name in the structure file. Each system was inserted into a rectangular periodic box with 1 nm of padding around each side of the protein and treated with the Amber03ws forcefield ((*Best et al., 2014*)) and the TIP4p2005 water model ((*Abascal and Vega, 2005*)). The wild-type system contained 2606 protein atoms, 122623 TIP4P water molecules, 439 Na atoms and 342 Cl atoms for a total of 490489 atoms. The F527A mutant matched this setup but with 2596 protein atoms for a total of 493769 atoms. The 0Q system contained 2487 protein atoms, 210607 water molecules, 237 Na atoms and 240 Cl atoms for a total of 846088 atoms. A 50ns equilibration run was performed for each system using the v-rescale thermostat and Berendsen barostat ((*Berendsen et al., 1984*)). After equilibration, each protein atom was marked within the base topology file as a “hot atom” and replicas were created by scaling interactions involving these “hot atoms” by a lambda value corresponding to temperatures in the range of 290K to 405K. The wild-type and F527A systems each had 20 replicas while 0Q had 12 replicas in order to achieve an exchange probability between 35% and 45%. REST2 production runs of each system were performed for 500ns using the v-rescale thermostat and Parrinello-Rahman barostat ((*Parrinello and Rahman, 1980*)). Convergence was verified using a split analysis of the radius of gyration values for each system at multiple temperatures (Fig. S20).

Martini Simulation Details

All condensate simulations contain 100 ELF3-PrD monomers in a 40 nm³ box. Monomers used for the initial configurations were obtained by performing an RMSD-based cluster analysis on previously run all-atom REST2 simulations of ELF3-PrD monomer. A representative structure was taken from 10 of the most populated clusters and each was converted to Martini format using Martini 3.0 and inserted into the box 10 times for a total of 100 monomers. Protein-water interactions were scaled up by 10%, a procedure shown to bring simulated proteins into better agreement with experimental measurements ((*Thomassen et al., 2022*)). Each initial configuration was equilibrated for 50ns to four different temperatures, 290K, 300K, 320K and 340K followed by production runs of 3 μ s. This process was performed for wildtype (7Q), 0Q, and 19Q ELF3-PrD, as well as the F527A mutant. All systems used explicit Martini water and contained 150mM NaCl. Three replicates were performed for each condition with the exception of wild type and F527A at 300K and 19Q at 340K, which have two replicates each.

Statistical Analysis of Ensemble Contact Maps

We used two types of contact analysis to identify mechanistically important interactions and characterize differences in our ELF3-PrD chain-growth ensembles. Both approaches, E-PCA and I-PCA ((*Johnson et al., 2018*)), are part of the CAMERRA family of contact analyses developed by Shen *et*

al. and seek to explain sources of conformational variance of protein ensembles. In E-PCA, contact maps from each conformation of the ensemble are used to produce a contact covariance matrix of each possible residue pair, i and j , with all other residue pairs, k and l . PCA is performed using this covariance matrix as input and the top PCs are returned. This method has been used to uncover coordinated interactions underlying mechanisms of allostery, cooperative ligand binding, and general collective motions of a variety of proteins ((*Lindsay et al., 2018; Clark et al., 2016; Johnson et al., 2015a,b*)). Due to the absence of long-range interactions in the HCG method used to generate our ensembles, E-PCA signals will be more prominent for local interactions. The dynamics of folded proteins tend to be dominated by the top few modes (PCs), however, intrinsically disordered proteins exhibit a slower eigen decay in which the first several PCs may contribute nearly equally (Fig. S21).

The second CAMERRA variant, I-PCA, provides information on the packing of residues and can identify regions of the sequence in which residues tend to interact more frequently. In this approach, individual residue pairs are considered against the rest of the protein, resulting in a covariance matrix of size $N \times N$ as opposed to $N^2 \times N^2$ for the E-PCA variant. This approach has been prominently used in studies of chromatin structure to differentiate transcriptionally active and inactive regions (euchromatin and heterochromatin, respectively) from Hi-C maps ((*Lajoie et al., 2015*)). When applied to natively structured proteins I-PCA tends to identify the individual domains of the protein and while IDPs do not contain structured domains, I-PCA can nevertheless identify dynamically associated domains ((*Lindsay et al., 2021*)).

Acknowledgments

We would like to acknowledge Lucy Reading-Ikkanda for providing the illustrations in Fig. 1A, Lukas Stelzl and Lisa Pietrek for developing the hierarchical chain growth method used here, and Javier González Delgado and Juan Cortes for characterizing the conformational landscape of the ELF3 REST2 simulations using their WARIO contact analysis method and subsequent discussions. VBPL is supported by the Brazilian agencies FAPESP (Grants 2023/02219-1 and 2022/07231-7) and National Council for Scientific and Technological Development – CNPq (Grant 310017/2020-3).

Data Availability

The datasets generated and analyzed during the current study are available for download:

HCG ensembles: https://users.flatironinstitute.org/~ccb/smbp/elf3_data/elf3prd_hcg.zip

REST2 trajectories: <https://zenodo.org/doi/10.5281/zenodo.13226984>

Martini trajectories: <https://zenodo.org/doi/10.5281/zenodo.13285376>

Analysis scripts and code: <https://github.com/flatironinstitute/ELF3>

Citations

References

- Abascal JLF**, Vega C. A general purpose model for the condensed phases of water: TIP4P/2005. The Journal of Chemical Physics. 2005; 123(23):234505. <https://doi.org/10.1063/1.2121687>, doi: 10.1063/1.2121687.
- Abraham MJ**, Murtola T, Schulz R, Páll S, Smith JC, Hess B, Lindahl E. GROMACS: High performance molecular simulations through multi-level parallelism from laptops to supercomputers. SoftwareX. 2015; 1-2:19–25. <https://www.sciencedirect.com/science/article/pii/S2352711015000059>, doi: 10.1016/j.softx.2015.06.001.
- Anwer MU**, Boikoglou E, Herrero E, Hallstein M, Davis AM, Velikkakam James G, Nagy F, Davis SJ. Natural variation reveals that intracellular distribution of ELF3 protein is associated with function in the circadian clock. eLife. 2014; 3. doi: 10.7554/eLife.02206, place: England.
- Bari KJ**, Prakashchand DD. Fundamental Challenges and Outlook in Simulating Liquid–Liquid Phase Separation of Intrinsically Disordered Proteins. The Journal of Physical Chemistry Letters. 2021; 12(6):1644–1656. <https://doi.org/10.1021/acs.jpclett.0c03404>, doi: 10.1021/acs.jpclett.0c03404, publisher: American Chemical Society.

- 648 **Benayad Z**, von Bülow S, Stelzl LS, Hummer G. Simulation of FUS Protein Condensates with an Adapted Coarse-
649 Grained Model. *Journal of Chemical Theory and Computation*. 2021; 17(1):525–537. [https://doi.org/10.1021/](https://doi.org/10.1021/acs.jctc.0c01064)
650 [acs.jctc.0c01064](https://doi.org/10.1021/acs.jctc.0c01064), doi: 10.1021/acs.jctc.0c01064, publisher: American Chemical Society.
- 651 **Berendsen HJC**, Postma JPM, van Gunsteren WF, DiNola A, Haak JR. Molecular dynamics with coupling to an
652 external bath. *The Journal of Chemical Physics*. 1984; 81(8):3684–3690. <https://doi.org/10.1063/1.448118>,
653 doi: 10.1063/1.448118, publisher: American Institute of Physics.
- 654 **Best RB**, Zheng W, Mittal J. Balanced Protein-Water Interactions Improve Properties of Disordered Proteins
655 and Non-Specific Protein Association. *Journal of chemical theory and computation*. 2014; 10(11):5113–5124.
656 <https://pubmed.ncbi.nlm.nih.gov/25400522>, doi: 10.1021/ct500569b, edition: 2014/10/16.
- 657 **Bohn L**, Huang J, Weidig S, Yang Z, Heidersberger C, Genty B, Falter-Braun P, Christmann A, Grill E. The tempera-
658 ture sensor TWA1 is required for thermotolerance in Arabidopsis. *Nature*. 2024 May; 629(8014):1126–1132.
- 659 **Box MS**, Huang BE, Domijan M, Jaeger KE, Khattak AK, Yoo SJ, Sedivy EL, Jones DM, Hearn TJ, Webb AAR,
660 Grant A, Locke JCW, Wigge PA. ELF3 Controls Thermoresponsive Growth in Arabidopsis. *Current Bi-*
661 *ology*. 2015; 25(2):194–199. <https://www.sciencedirect.com/science/article/pii/S0960982214014249>, doi:
662 10.1016/j.cub.2014.10.076.
- 663 **von Bulow S**, Tesei G, Lindorff-Larsen K. Prediction of phase separation propensities of disordered proteins
664 from sequence. *bioRxiv*. 2024; <https://www.biorxiv.org/content/early/2024/06/03/2024.06.03.597109>, doi:
665 10.1101/2024.06.03.597109.
- 666 **Choi JM**, Dar F, Pappu RV. LASSI: A lattice model for simulating phase transitions of multivalent proteins. *PLoS*
667 *computational biology*. 2019; 15(10):e1007028. doi: 10.1371/journal.pcbi.1007028, place: United States.
- 668 **Clark AK**, Wilder JH, Grayson AW, Johnson QR, Lindsay RJ, Nellas RB, Fernandez EJ, Shen T. The Promiscuity of
669 Allosteric Regulation of Nuclear Receptors by Retinoid X Receptor. *The journal of physical chemistry B*. 2016;
670 120(33):8338–8345. doi: 10.1021/acs.jpcc.6b02057.
- 671 **Das RK**, Pappu RV. Conformations of intrinsically disordered proteins are influenced by linear sequence
672 distributions of oppositely charged residues. *Proceedings of the National Academy of Sciences*. 2013;
673 110(33):13392. <http://www.pnas.org/content/110/33/13392.abstract>, doi: 10.1073/pnas.1304749110.
- 674 **Davey NE**, Van Roey K, Weatheritt RJ, Toedt G, Uyar B, Altenberg B, Budd A, Diella F, Dinkel H, Gibson TJ.
675 Attributes of short linear motifs. *Molecular BioSystems*. 2012; 8(1):268–281. [http://dx.doi.org/10.1039/](http://dx.doi.org/10.1039/C1MB05231D)
676 [C1MB05231D](http://dx.doi.org/10.1039/C1MB05231D), doi: 10.1039/C1MB05231D, publisher: The Royal Society of Chemistry.
- 677 **Dignon GL**, Zheng W, Best RB, Kim YC, Mittal J. Relation between single-molecule properties and phase behavior
678 of intrinsically disordered proteins. *Proceedings of the National Academy of Sciences*. 2018; 115(40):9929–
679 9934. <https://doi.org/10.1073/pnas.1804177115>, doi: 10.1073/pnas.1804177115, publisher: Proceedings of
680 the National Academy of Sciences.
- 681 **Dignon GL**, Zheng W, Kim YC, Mittal J. Temperature-Controlled Liquid–Liquid Phase Separation of Disor-
682 dered Proteins. *ACS Central Science*. 2019; 5(5):821–830. <https://doi.org/10.1021/acscentsci.9b00102>, doi:
683 10.1021/acscentsci.9b00102, PMID: 31139718.
- 684 **González-Delgado J**, Bernadó P, Neuvial P. WARIO: Weighted families of contact maps to characterize confor-
685 mational ensembles of (highly-)flexible proteins. *Research Square*. 2024 Apr; 64(8).
- 686 **Jo S**, Jiang W. A generic implementation of replica exchange with solute tempering (REST2) algorithm in NAMD
687 for complex biophysical simulations. *Computer Physics Communications*. 2015; 197:304–311. [https://www.](https://www.sciencedirect.com/science/article/pii/S001046551500329X)
688 [sciencedirect.com/science/article/pii/S001046551500329X](https://www.sciencedirect.com/science/article/pii/S001046551500329X), doi: 10.1016/j.cpc.2015.08.030.
- 689 **Johnson QR**, Lindsay RJ, Nellas RB, Fernandez EJ, Shen T. Mapping Allostery through Computational Glycine
690 Scanning and Correlation Analysis of Residue–Residue Contacts. *Biochemistry*. 2015; 54(7):1534–1541. <https://doi.org/10.1021/bi501152d>, doi: 10.1021/bi501152d, publisher: American Chemical Society.
- 692 **Johnson QR**, Lindsay RJ, Petridis L, Shen T. Investigation of Carbohydrate Recognition via Computer Simulation.
693 *Molecules* (Basel, Switzerland). 2015; 20(5):7700–7718. doi: 10.3390/molecules20057700.
- 694 **Johnson QR**, Lindsay RJ, Shen T. CAMERRA: An analysis tool for the computation of conformational dynamics by
695 evaluating residue-residue associations. *Journal of computational chemistry*. 2018; 39(20):1568–1578. doi:
696 10.1002/jcc.25192, place: United States.

- 697 **Joseph JA**, Espinosa JR, Sanchez-Burgos I, Garaizar A, Frenkel D, Collepardo-Guevara R. Thermodynam-
698 ics and kinetics of phase separation of protein-RNA mixtures by a minimal model. *Biophysical Jour-*
699 *nal*. 2021; 120(7):1219–1230. <https://www.sciencedirect.com/science/article/pii/S000634952100117X>, doi:
700 <https://doi.org/10.1016/j.bpj.2021.01.031>.
- 701 **Joseph JA**, Reinhardt A, Aguirre A, Chew PY, Russell KO, Espinosa JR, Garaizar A, Collepardo-Guevara R. Physics-
702 driven coarse-grained model for biomolecular phase separation with near-quantitative accuracy. *Nature*
703 *computational science*. 2021; 1(11):732–743. doi: 10.1038/s43588-021-00155-3, place: United States.
- 704 **Jung JH**, Barbosa AD, Hutin S, Kumita JR, Gao M, Derwort D, Silva CS, Lai X, Pierre E, Geng F, Kim SB, Baek S, Zubi-
705 eta C, Jaeger KE, Wigge PA. A prion-like domain in ELF3 functions as a thermosensor in Arabidopsis. *Nature*.
706 2020; 585(7824):256–260. <https://doi.org/10.1038/s41586-020-2644-7>, doi: 10.1038/s41586-020-2644-7.
- 707 **Kerbler SM**, Wigge PA. Temperature Sensing in Plants. *Annual Review of Plant Biology*. 2023; 74(1):341–366.
708 <https://www.annualreviews.org/doi/10.1146/annurev-arplant-102820-102235>, doi: 10.1146/annurev-arplant-
709 102820-102235.
- 710 **Khare SD**, Ding F, Gwanmesia KN, Dokholyan NV. Molecular Origin of Polyglutamine Aggregation in Neu-
711 rodegenerative Diseases. *PLOS Computational Biology*. 2005; 1(3):e30. <https://doi.org/10.1371/journal.pcbi.0010030>, doi: 10.1371/journal.pcbi.0010030.
- 712 **Lajoie BR**, Dekker J, Kaplan N. The Hitchhiker's guide to Hi-C analysis: practical guidelines. *Methods* (San Diego,
713 Calif). 2015; 72:65–75. doi: 10.1016/j.ymeth.2014.10.031.
- 714 **Lancaster AK**, Nutter-Upham A, Lindquist S, King OD. PLAAC: a web and command-line application to identify
715 proteins with prion-like amino acid composition. *Bioinformatics* (Oxford, England). 2014; 30(17):2501–2502.
716 <https://pubmed.ncbi.nlm.nih.gov/24825614>, doi: 10.1093/bioinformatics/btu310, edition: 2014/05/13.
- 717 **Lin J**, Xu Y, Zhu Z. Emerging Plant Thermosensors: From RNA to Protein. *Trends in Plant Sci-*
718 *ence*. 2020; 25(12):1187–1189. <https://www.sciencedirect.com/science/article/pii/S1360138520302545>, doi:
719 <https://doi.org/10.1016/j.tplants.2020.08.007>.
- 720 **Lindsay RJ**, Mansbach RA, Gnanakaran S, Shen T. Effects of pH on an IDP conformational ensemble explored by
721 molecular dynamics simulation. *Biophys Chem*. 2021; 271:106552. doi: 10.1016/j.bpc.2021.106552, edition:
722 20210126.
- 723 **Lindsay RJ**, Pham B, Shen T, McCord RP. Characterizing the 3D structure and dynamics of chromosomes
724 and proteins in a common contact matrix framework. *Nucleic acids research*. 2018; 46(16):8143–8152. doi:
725 10.1093/nar/gky604.
- 726 **Liu X**, Covington M, Fankhauser C, Chory J, Wagner D. ELF3 Encodes a Circadian Clock-Regulated Nuclear
727 Protein That Functions in an Arabidopsis PHYB Signal Transduction Pathway. *Plant Cell*. 2001; 13(6):1293–
728 1304. <http://www.plantcell.org/cgi/doi/10.1105/tpc.13.6.1293>, doi: 10.1105/tpc.13.6.1293.
- 729 **Lotthammer JM**, Ginell GM, Griffith D, Emenecker RJ, Holehouse AS. Direct prediction of intrinsically disor-
730 dered protein conformational properties from sequence. *Nature Methods*. 2024; <https://doi.org/10.1038/s41592-023-02159-5>, doi: 10.1038/s41592-023-02159-5.
- 731 **Mao AH**, Crick SL, Vitalis A, Chicoine CL, Pappu RV. Net charge per residue modulates conformational en-
732 sembles of intrinsically disordered proteins. *Proceedings of the National Academy of Sciences*. 2010;
733 107(18):8183–8188. <https://www.pnas.org/doi/abs/10.1073/pnas.0911107107>, doi: 10.1073/pnas.0911107107.
- 734 **Maristany MJ**, Gonzalez AA, Collepardo-Guevara R, Joseph JA. Universal predictive scaling laws of phase sepa-
735 ration of prion-like low complexity domains. *bioRxiv*. 2023; <https://www.biorxiv.org/content/early/2023/06/14/2023.06.14.543914>, doi: 10.1101/2023.06.14.543914.
- 736 **Martin EW**, Holehouse AS, Peran I, Farag M, Incicco JJ, Bremer A, Grace CR, Soranno A, Pappu RV, Mittag T.
737 Valence and patterning of aromatic residues determine the phase behavior of prion-like domains. *Science*
738 (New York, NY). 2020; 367(6478):694–699. doi: 10.1126/science.aaw8653, place: United States.
- 739 **Martin EW**, Mittag T. Relationship of Sequence and Phase Separation in Protein Low-Complexity
740 Regions. *Biochemistry*. 2018; 57(17):2478–2487. <https://doi.org/10.1021/acs.biochem.8b00008>, doi:
741 [10.1021/acs.biochem.8b00008](https://doi.org/10.1021/acs.biochem.8b00008), publisher: American Chemical Society.
- 742 **Martin EW**, Mittag T. Relationship of sequence and phase separation in protein low-complexity regions. *Bio-*
743 *chemistry*. 2018 May; 57(17):2478–2487.
- 744 **Martin EW**, Mittag T. Relationship of sequence and phase separation in protein low-complexity regions. *Bio-*
745 *chemistry*. 2018 May; 57(17):2478–2487.

- 747 **Moses D**, Ginell GM, Holehouse AS, Sukenik S. Intrinsically disordered regions are poised to act as sensors of
748 cellular chemistry. *Trends in Biochemical Sciences*. 2023; 48(12):1019–1034. [https://linkinghub.elsevier.com/
749 retrieve/pii/S0968000423002049](https://linkinghub.elsevier.com/retrieve/pii/S0968000423002049), doi: 10.1016/j.tibs.2023.08.001.
- 750 **Nag N**, Chetri PB, Uversky VN, Giri R, Tripathi T. Chapter 31 - Experimental methods to study intrinsically disor-
751 dered proteins. In: Tripathi T, Dubey VK, editors. *Advances in Protein Molecular and Structural Biology Methods*
752 Academic Press; 2022.p. 505–533. <https://www.sciencedirect.com/science/article/pii/B9780323902649000313>,
753 doi: <https://doi.org/10.1016/B978-0-323-90264-9.00031-3>.
- 754 **Nusinow DA**, Helfer A, Hamilton EE, King JJ, Imaizumi T, Schultz TF, Farré EM, Kay SA. The ELF4-ELF3-LUX
755 complex links the circadian clock to diurnal control of hypocotyl growth. *Nature*. 2011; 475(7356):398–402.
756 <https://pubmed.ncbi.nlm.nih.gov/21753751>, doi: 10.1038/nature10182.
- 757 **Paik I**, Huq E. Plant photoreceptors: Multi-functional sensory proteins and their signaling networks. *Seminars*
758 *in cell & developmental biology*. 2019; 92:114–121. doi: 10.1016/j.semcdb.2019.03.007, place: England.
- 759 **Parrinello M**, Rahman A. Crystal Structure and Pair Potentials: A Molecular-Dynamics Study. *Physical Review*
760 *Letters*. 1980; 45(14):1196–1199. <https://link.aps.org/doi/10.1103/PhysRevLett.45.1196>, doi: 10.1103/Phys-
761 *RevLett.45.1196*, publisher: American Physical Society.
- 762 **Pietrek LM**, Stelzl LS, Hummer G. Hierarchical Ensembles of Intrinsically Disordered Proteins at Atomic Resolu-
763 tion in Molecular Dynamics Simulations. *Journal of Chemical Theory and Computation*. 2020; 16(1):725–737.
764 <https://doi.org/10.1021/acs.jctc.9b00809>, doi: 10.1021/acs.jctc.9b00809
- 765 **Quint M**, Delker C, Franklin KA, Wigge PA, Halliday KJ, van Zanten M. Molecular and genetic control of plant
766 thermomorphogenesis. *Nat Plants*. 2016 Jan; 2:15190.
- 767 **Quiroz FG**, Chilkoti A. Sequence heuristics to encode phase behaviour in intrinsically disordered protein
768 polymers. *Nature Materials*. 2015; 14(11):1164–1171. <https://www.nature.com/articles/nmat4418>, doi:
769 10.1038/nmat4418.
- 770 **Ramazzotti M**, Monsellier E, Kamoun C, Degl'Innocenti D, Melki R. Polyglutamine repeats are associated to spe-
771 cific sequence biases that are conserved among eukaryotes. *PloS one*. 2012; 7(2):e30824. doi: 10.1371/jour-
772 *nal.pone.0030824*.
- 773 **Regy RM**, Thompson J, Kim YC, Mittal J. Improved coarse-grained model for studying sequence dependent
774 phase separation of disordered proteins. *Protein science : a publication of the Protein Society*. 2021;
775 30(7):1371–1379. doi: 10.1002/pro.4094, place: United States.
- 776 **Reiner A**, Dragatsis I, Dietrich P. Genetics and neuropathology of Huntington's disease. *International review*
777 *of neurobiology*. 2011; 98:325–372. doi: 10.1016/B978-0-12-381328-2.00014-6, place: United States.
- 778 **Robustelli P**, Piana S, Shaw DE. Developing a molecular dynamics force field for both folded and disordered
779 protein states. *Proceedings of the National Academy of Sciences of the United States of America*. 2018;
780 115(21):E4758–E4766. doi: 10.1073/pnas.1800690115, place: United States.
- 781 **Romero P**, Obradovic Z, Li X, Garner EC, Brown CJ, Dunker AK. Sequence complexity of disordered
782 protein. *Proteins: Structure, Function, and Bioinformatics*. 2001; 42(1):38–48. [https://onlinelibrary.
783 wiley.com/doi/abs/10.1002/1097-0134%2820010101%2942%3A1%3C38%3A%3AAID-PROT50%3E3.0.
784 CO%3B2-3](https://onlinelibrary.wiley.com/doi/abs/10.1002/1097-0134%2820010101%2942%3A1%3C38%3A%3AAID-PROT50%3E3.0.CO%3B2-3), doi: [https://doi.org/10.1002/1097-0134\(20010101\)42:1<38::AID-PROT50%3E3.0.CO%3B2-3](https://doi.org/10.1002/1097-0134(20010101)42:1<38::AID-PROT50%3E3.0.CO%3B2-3), _eprint:
785 [https://onlinelibrary.wiley.com/doi/pdf/10.1002/1097-0134%2820010101%2942%3A1%3C38%3A%3AAID-
786 PROT50%3E3.0.CO%3B2-3](https://onlinelibrary.wiley.com/doi/pdf/10.1002/1097-0134%2820010101%2942%3A1%3C38%3A%3AAID-PROT50%3E3.0.CO%3B2-3).
- 787 **Ruff KM**, Roberts S, Chilkoti A, Pappu RV. Advances in Understanding Stimulus-Responsive Phase Behavior
788 of Intrinsically Disordered Protein Polymers. *Journal of Molecular Biology*. 2018 Nov; 430(23):4619–4635.
789 <https://linkinghub.elsevier.com/retrieve/pii/S0022283618306594>, doi: 10.1016/j.jmb.2018.06.031.
- 790 **Schramm A**, Bignon C, Brocca S, Grandori R, Santambrogio C, Longhi S. An arsenal of methods for the
791 experimental characterization of intrinsically disordered proteins – How to choose and combine them?
792 *Archives of Biochemistry and Biophysics*. 2019; 676:108055. [https://www.sciencedirect.com/science/article/
793 pii/S0003986119304527](https://www.sciencedirect.com/science/article/pii/S0003986119304527), doi: <https://doi.org/10.1016/j.abb.2019.07.020>.
- 794 **Sengupta P**, Garrity P. Sensing temperature. *Current Biology*. 2013; 23(8):R304–R307. [https://www.sciencedirect.
795 com/science/article/pii/S0960982213002753](https://www.sciencedirect.com/science/article/pii/S0960982213002753), doi: 10.1016/j.cub.2013.03.009.

- 796 **Song D**, Luo R, Chen HF. The IDP-Specific Force Field ff14IDPSFF Improves the Conformer Sampling of In-
797 trinsically Disordered Proteins. *Journal of Chemical Information and Modeling*. 2017; 57(5):1166–1178.
798 <https://doi.org/10.1021/acs.jcim.7b00135>, doi: 10.1021/acs.jcim.7b00135, publisher: American Chemical So-
799 ciety.
- 800 **Stelzl LS**, Pietrek LM, Holla A, Oroz J, Sikora M, Köfinger J, Schuler B, Zweckstetter M, Hummer G. Global Struc-
801 ture of the Intrinsically Disordered Protein Tau Emerges from Its Local Structure. *JACS Au*. 2022; 2(3):673–686.
802 <https://doi.org/10.1021/jacsau.1c00536>, doi: 10.1021/jacsau.1c00536, publisher: American Chemical Society.
- 803 **Tesei G**, Lindorff-Larsen K. Improved predictions of phase behaviour of intrinsically disordered proteins by
804 tuning the interaction range. *Open research Europe*. 2022; 2:94. doi: 10.12688/openreseurope.14967.2,
805 place: Belgium.
- 806 **Tesei G**, Schulze TK, Crehuet R, Lindorff-Larsen K. Accurate model of liquid-liquid phase behavior of intrinsically
807 disordered proteins from optimization of single-chain properties. *Proceedings of the National Academy of*
808 *Sciences of the United States of America*. 2021; 118(44). doi: 10.1073/pnas.2111696118, place: United States.
- 809 **Thines B**, Harmon FG. Ambient temperature response establishes ELF3 as a required component of the core
810 Arabidopsis circadian clock. *Proceedings of the National Academy of Sciences*. 2010; 107(7):3257–3262. <https://doi.org/10.1073/pnas.0911006107>, doi: 10.1073/pnas.0911006107, publisher: Proceedings of the National
811 Academy of Sciences.
- 812 **Thomassen FE**, Pesce F, Roesgaard MA, Tesei G, Lindorff-Larsen K. Improving Martini 3 for Disordered and
813 Multidomain Proteins. *Journal of Chemical Theory and Computation*. 2022; 18(4):2033–2041. <https://doi.org/10.1021/acs.jctc.1c01042>, doi: 10.1021/acs.jctc.1c01042, publisher: American Chemical Society.
- 814 **Totzeck F**, Andrade-Navarro MA, Mier P. The Protein Structure Context of PolyQ Regions. *PloS one*. 2017;
815 12(1):e0170801. doi: 10.1371/journal.pone.0170801.
- 816 **Valley CC**, Cembran A, Perlmutter JD, Lewis AK, Labello NP, Gao J, Sachs JN. The methionine-aromatic motif
817 plays a unique role in stabilizing protein structure. *The Journal of biological chemistry*. 2012; 287(42):34979–
818 34991. doi: 10.1074/jbc.M112.374504, place: United States.
- 819 **Vernon RM**, Chong PA, Tsang B, Kim TH, Bah A, Farber P, Lin H, Forman-Kay JD. Pi-Pi contacts are an overlooked
820 protein feature relevant to phase separation. *eLife*. 2018 feb; 7:e31486. <https://doi.org/10.7554/eLife.31486>,
821 doi: 10.7554/eLife.31486.
- 822 **Viegas RG**, Martins IBS, Sanches MN, Oliveira Junior AB, Camargo JBd, Paulovich FV, Leite VBP. ELViM: Explor-
823 ing biomolecular energy landscapes through multidimensional visualization. *J Chem Inf Model*. 2024 Apr;
824 64(8):3443–3450.
- 825 **Wang L**, Friesner RA, Berne BJ. Replica exchange with solute scaling: a more efficient version of replica exchange
826 with solute tempering (REST2). *J Phys Chem B*. 2011 aug; 115(30):9431–9438.
- 827 **Wang L**, Friesner RA, Berne BJ. Replica exchange with solute scaling: a more efficient version of replica exchange
828 with solute tempering (REST2). *The journal of physical chemistry B*. 2011; 115(30):9431–9438. <https://pubmed.ncbi.nlm.nih.gov/21714551>, doi: 10.1021/jp204407d, edition: 2011/07/07.
- 829 **Wu H**, Wolynes PG, Papoian GA. AWSEM-IDP: A Coarse-Grained Force Field for Intrinsically Disordered Pro-
830 teins. *The Journal of Physical Chemistry B*. 2018 Dec; 122(49):11115–11125. <https://doi.org/10.1021/acs.jpcb.8b05791>, doi: 10.1021/acs.jpcb.8b05791.
- 831 **Zagotta M**, Shannon S, Jacobs C, Meeks-Wagner D. Early-Flowering Mutants of Arabidopsis thaliana. *Functional*
832 *Plant Biology*. 1992; 19(4):411–418. <https://doi.org/10.1071/PP9920411>.
- 833 **Zhang C**, Gao M, Seitz NC, Angel W, Hallworth A, Wiratan L, Darwish O, Alkharouf N, Dawit T, Lin D, Egoshi R,
834 Wang X, McClung CR, Lu H. LUX ARRHYTHMO mediates crosstalk between the circadian clock and defense
835 in Arabidopsis. *Nature Communications*. 2019; 10(1):2543. <https://doi.org/10.1038/s41467-019-10485-6>, doi:
836 10.1038/s41467-019-10485-6.
- 837 **Zhang Y**, Chen J. Recalibrating replica exchange solute tempering for sampling disordered protein con-
838 formations. *Biophysical Journal*. 2022; 121(3):196a. <https://doi.org/10.1016/j.bpj.2021.11.1730>, doi:
839 10.1016/j.bpj.2021.11.1730, publisher: Elsevier.
- 840 **Zhao C**, Liu B, Piao S, Wang X, Lobell DB, Huang Y, Huang M, Yao Y, Bassu S, Ciais P, Durand JL, Elliott J, Ewert
841 F, Janssens IA, Li T, Lin E, Liu Q, Martre P, Müller C, Peng S, et al. Temperature increase reduces global yields
842 of major crops in four independent estimates. *Proceedings of the National Academy of Sciences*. 2017;
843 114(35):9326–9331. <https://pnas.org/doi/full/10.1073/pnas.1701762114>, doi: 10.1073/pnas.1701762114.