

HELMINTHOLOGIA, 53, 3: 211 - 223, 2016

Review

Checklist of sea turtles endohelminth in Neotropical region

M. R. WERNECK¹, R. J. DA SILVA²

¹BW Veterinary Consulting, Rua Ponciano Eugênio Duarte 203, Centro, Zip Code: 11680-000, Ubatuba, São Paulo State, Brazil,
*E-mail: max@bwvet.com.br; ²São Paulo State University (UNESP), Institute of Biosciences, Department of Parasitology,
Laboratory of Wild Animals Parasitology, Botucatu, São Paulo State, Brazil

Article info

Received December 17, 2015
Accepted May 6, 2016

Summary

This paper presents a list of parasites described in sea turtles from the Neotropical region. Through the review of literature the occurrence of 79 taxa of helminthes parasites were observed, mostly consisting of the Phylum Platyhelminthes with 76 species distributed in 14 families and 2 families of the Phylum Nematoda within 3 species. Regarding the parasite records, the most studied host was the green turtle (*Chelonia mydas*) followed by the hawksbill turtle (*Eretmochelys imbricata*), olive ridley turtle (*Lepidochelys olivacea*), loggerhead turtle (*Caretta caretta*) and leatherback turtle (*Derموchelys coriacea*). Overall helminths were reported in 12 countries and in the Caribbean Sea region. This checklist is the largest compilation of data on helminths found in sea turtles in the Neotropical region.
Keywords: checklist; helminths; marine turtles; parasites

Introduction

The Neotropical region corresponds to Central and South America, where five species of sea turtles are found: Green turtle (*Chelonia mydas* Linnaeus, 1758), loggerhead turtle (*Caretta caretta* Linnaeus, 1758), hawksbill turtle (*Eretmochelys imbricata* Linnaeus, 1766), Olive ridley turtle (*Lepidochelys olivacea* Eschscholtz, 1829) and leatherback turtle (*Derموchelys coriacea* Vandelli, 1761) (Lutz & Musick, 1996). The International Union for Conservation of Nature (IUCN, 2015) considers *C. caretta*, *C. mydas* and *L. olivacea* as endangered and both *D. coriacea* and *E. imbricata* as critically threatened.

Parasitological analyses of sea turtles have been conducted for many years and the results have contributed to a better understanding of the helminths of this group of hosts. Thus, the aim of this paper was to offer a comprehensive bibliographic review of the species of endohelminths found in sea turtles in the Neotropics.

Material and Methods

This checklist was developed using only scientific papers on endohelminths identified in sea turtles published between 1911 and 2016. Dissertations, theses and abstracts from conferences were not considered.

The data are presented in two forms: The first part of the paper presents a list of helminths (and synonyms) separated by family, genus and species and includes the infection site, location, infected host and author of the reference. The second part presents the hosts and a list of helminths reported in these hosts. For the classification, the taxonomic proposal presented by the World Register of Marine Species (WoRMS, 2015) was used.

Results

This checklist records the occurrence of 79 taxa of parasites. Se

venty six represent the phylum Platyhelminthes (one from the order Aspidogastriida, 11 from the order Diplostomida and 64 from the order Plagiorchiida) represented by 14 families. Three represent the phylum Nematoda, in which two families have been found in the sea turtles *C. mydas*, *C. caretta*, *E. imbricata*, *L. olivacea* and *D. coriacea*. Overall helminths are reported in 12 countries and three are reported for the region in which the present study took place (i.e. the Caribbean Sea).

Parasite-Host list

Phylum Platyhelminthes Gegenbaur, 1859

Class Trematoda Rudolphi, 1808

Subclass Aspidogastrea Faust & Tang, 1936

Order Aspidogastriida Skrjabin & Guschanskaja, 1962

Superfamily Apidogastroidea Poche, 1907

Family Aspidogastridae Poche, 1907

Lophotaspis vallei (Stossich, 1899)

Site of infection: Esophagus and stomach.

Host and distribution: *C. caretta* from Brazil (Araújo, 1941)

Subclass Digenea Carus, 1863

Order Plagiorchiida La Rue, 1957

Suborder Echinostomata Szidat, 1939

Super Family Echinostomatoidea Looss, 1902

Family Calycodidae Dollfus, 1929

Calycodes anthos (Braun, 1899) Looss, 1901.

Site of infection: Small and large intestine.

Host and distribution: *C. caretta* from Brazil (Werneck *et al.*, 2008a), *C. mydas* from Panamá (Caballero *et al.*, 1955) and Brazil (Binoti *et al.*, 2016); *D. coriacea* from Uruguay (Werneck *et al.*, 2012) and *L. olivacea* from México (Pérez-Ponce de Leon *et al.*, 1996).

Calycodes caborjoensis Fischthal e Acholonu, 1976

Site of infection: small intestine of *E. imbricata* from Puerto Rico (Fischthal & Acholonu, 1976; Dyer *et al.*, 1995a).

Familia Rhytidodidae Odhner, 1926

Rhytidodes gelatinosus (Rudolphi, 1819) Looss, 1901

Site of infection: Stomach and intestine.

Host and distribution: *C. caretta* from Brazil (Viana, 1924; Travassos *et al.*, 1969); *C. mydas* from Panamá (Caballero, 1954) and Brazil (Werneck & Silva, 2015) and *E. imbricata* from Puerto Rico (Fischthal & Acholonu, 1976; Dyer *et al.*, 1995a) and Cuba (Vigueras, 1955).

Rhytidodoides intestinalis Price, 1939

Site of infection: Gall bladder.

Host and distribution: *C. mydas* from Panamá (Caballero, 1954) and Costa Rica (Santoro *et al.*, 2006).

Rhytidodoides similis Price, 1939

Site of infection: Gall bladder.

Host and distribution: *C. mydas* from Panamá (Caballero, 1954), Costa Rica (Santoro *et al.*, 2006) and Brazil (Werneck *et al.*, 2015a).

Suborder Hemiurata Markevitsch 1951

Superfamily Hemiuroidea Looss, 1899

Family Sclerodistomidae Odner, 1927

Prosorichis psenopsis Yamaguti, 1934

Site of infection: Stomach.

Host and distribution: *L. olivacea* from México (Pérez-Ponce de Leon *et al.*, 1996).

Suborder Pronocephalata Olson, Cribb, Tkach, Bray & Littlewood, 2003

Superfamily Paramphistomoidea Fischoeder, 1901

Family Cladorchiidae Fischoeder, 1901

Schizamphistomum erratum Blair, 1983

(Synonym: *Schizamphistomoides erratum* Blair, 1983)

Site of infection: Intestine.

Host and distribution: *C. mydas* from Costa Rica (Santoro *et al.*, 2006).

Schizamphistomoides spinulosum (Looss, 1901) Stunkard, 1925

Site of infection: Large intestine.

Host and distribution: *C. mydas* from Panamá (Caballero *et al.*, 1955).

Schizamphistomum scleroporium (Creplin, 1844) Looss, 1912

(Synonym: *Schizamphistomoides chelonei* Gupta 1961)

Site of infection: Stomach and intestine.

Host and distribution: *C. mydas* from Brazil (Werneck & Silva 2015), Costa Rica (Santoro *et al.*, 2006), Cuba (Groschaft *et al.*, 1977) and Trinidad (The West Indies) (Gupta, 1961) and *E. imbricata* from Puerto Rico (Fischthal & Acholonu, 1976).

Schizamphistomum sp. Looss, 1912

Site of infection: Intestine.

Host and distribution: *C. mydas* from Brazil (Binoti *et al.*, 2016) and Puerto Rico (Dyer *et al.*, 1995b).

Family Microscaphidiidae Looss, 1900

Angiodictyum anteroporium Chattopadhyaya, 1972 (Taxon inquirendum)

Site of infection: Large intestine.

Host and distribution: *E. imbricata* from Puerto Rico (Dyer *et al.*, 1995a).

Angiodictyum longum Blair, 1986

Site of infection: Small intestine.

Host and distribution: *C. mydas* from Brazil (Werneck & Silva 2015; Binoti *et al.*, 2016).

- Angiodictyum mooreae* Dyer, Williams & Bunkley-Williams, 1995
Site of infection: Large intestine.
Host and distribution: *E. imbricata* from Puerto Rico (Dyer *et al.*, 1995c).
- Angiodictyum parallelum* (Looss, 1901) Looss, 1902
(Synonym: *Microscaphidium parallelum* Looss, 1901)
Site of infection: Small and large intestine.
Host and distribution: *C. mydas* from Brazil (Werneck & Silva, 2015) and Puerto Rico (Dyer *et al.*, 1991) and *E. imbricata* from Puerto Rico (Dyer *et al.*, 1995c).
- Deuterobaris chelonei* Gupta, 1961.
Site of infection: Intestine.
Host and distribution: *C. mydas* from Trinidad (The West India) (Gupta, 1961).
- Deuterobaris intestinalis* Mehrotra, 1973
Site of infection: Intestine.
Host and distribution: *C. mydas* from Costa Rica (Santoro *et al.*, 2006).
- Deuterobaris proteus* (Brandes, 1891) Looss, 1900
Site of infection: Esophagus, stomach, small and large intestine.
Host and distribution: *C. mydas* from Brazil (Werneck & Silva, 2015) and Puerto Rico (Dyer *et al.*, 1991; Dyer *et al.*, 1995b).
- Microscaphidium reticulare* (Van Beneden, 1859) Looss, 1900
(Synonym: *Microscaphidium caballeroi* Groschafft, 1977, *Microscaphidium japonicum* Oguro, 1941 and *Monostomum reticulare* Van Beneden, 1859)
Site of infection: Small and large intestine.
Host and distribution: *C. mydas* from Brazil (Werneck & Silva, 2015), Costa Rica (Santoro *et al.*, 2006), Cuba (Groschafft *et al.*, 1977) and Puerto Rico (Dyer *et al.*, 1995b).
- Microscaphidium warui* Blair, 1986
Site of infection: Urinary bladder.
Host and distribution: *C. mydas* from Costa Rica (Santoro *et al.*, 2006).
- Microscaphidium aberrans* Looss, 1902
Site of infection: Intestine.
Host and distribution: *C. mydas* from Panamá (Caballero, 1954)
- Neotangium travassosi* Ruiz, 1943
[Synonym: *Octangium travassosi* (Ruiz, 1943) Yamaguti, 1958, *Neotangium trinidadii* Gupta 1962]
Site of infection: Stomach and small and large intestine.
Host and distribution: *E. imbricata* from Puerto Rico (Fischthal & Acholonu, 1976).
- C. mydas* from Brazil [Ruiz 1943 (see Muniz-Pereira *et al.*, 2009); Travassos *et al.*, 1969; Werneck & Silva, 2015; Binoti *et al.*, 2016)], Caribbean Sea (Gupta, 1962) and Trinidad (The West Indies) (Gupta, 1961).
- Octangium hyphalum* Blair, 1987
Site of infection: Intestine.
Host and distribution: *C. mydas* from Costa Rica (Santoro *et al.*, 2006).
- Octangium sagitta* (Looss, 1899) Looss, 1902
[Synonym: *Microscapha sagitta* Looss, 1899, *Octangium hasta* Looss, 1902, *Octangium takanoi* Kobayashi, 1921 (see Blair, 1987)].
Site of infection: Stomach and small and large intestine.
Host and distribution: *C. mydas* from Puerto Rico (Dyer *et al.*, 1991) and *E. imbricata* from Puerto Rico (Fischthal & Acholonu, 1976; Dyer *et al.*, 1995c).
- Polyangium linguatula* (Looss, 1899)
[Synonym: *Microscaphidium linguatula* Looss, 1899, *Nephrobium colymbi* Poche, 1926, *Polyangium colymbi* (Poche, 1926), *Polyangium longiseminale* Chattopadhyaya, 1972 and *Polyangium miyajimai* Kobayashi, 1915].
Site of infection: Small and large intestine.
Host and distribution: *C. mydas* from Brazil (Teixeira de Freitas & Lent, 1938; Travassos *et al.*, 1969; Werneck & Silva, 2015; Binoti *et al.*, 2016), Costa Rica (Santoro *et al.*, 2006), Cuba (Groschafft *et al.*, 1977) and Puerto Rico (Dyer *et al.*, 1991; Dyer *et al.*, 1995c).
- Superfamily Pronocephaloidea Looss, 1899**
Family Pronocephalidae Looss, 1899
Adenogaster serialis Looss, 1901
Site of infection: Small and large intestine.
Host and distribution: *C. mydas* from Panamá (Caballero *et al.*, 1955) and Peru (Tantalean *et al.*, 1992); *E. imbricata* from Costa Rica (Santoro *et al.*, 2015) and Cuba (Vigueras, 1955) and *L. olivacea* from México (Pérez-Ponce de Leon *et al.*, 1996; Vivaldo *et al.*, 2006; Vivaldo *et al.*, 2009).
- Charaxicephaloides polyorchis* Groschafft & Tenora, 1978.
Site of infection: Stomach.
Host and distribution: *C. mydas* from Costa Rica (Santoro *et al.*, 2009a) and Cuba (Groschafft & Tenora, 1978; Groschafft *et al.*, 1977).
- Charaxicephaloides* sp. Groschafft & Tenora, 1978
Site of infection: Stomach.
Host and distribution: *C. mydas* from Costa Rica (Santoro *et al.*, 2006).

- Charaxicephalus robustus* Looss, 1901
Site of infection: Esophagus and stomach.
Host and distribution: *C. mydas* from Brazil (Binoti *et al.*, 2016) and Costa Rica (Santoro *et al.*, 2006).
- Cricocephalus americanus* Viguera, 1955
Site of infection: Stomach.
Host and distribution: *E. imbricata* from Cuba (Vieguera, 1955).
- Cricocephalus albus* (Kuhl & Van Hasselt, 1822)
(Synonym: *Cricocephalus delitescens* Looss, 1899 and *Cricocephalus koidzumii* Kobayashi, 1921).
Site of infection: Esophagus, stomach and small intestine.
Host and distribution: *C. mydas* from Brazil (Ruiz, 1946; Travassos *et al.*, 1969; Werneck & Silva, 2015; Binoti *et al.*, 2016), Costa Rica (Santoro *et al.*, 2006), Panama (Caballero *et al.*, 1955) and Trinidad (The West Indies) (Gupta, 1961); *E. imbricata* from Costa Rica (Santoro *et al.*, 2015) and Puerto Rico (Fischthal & Acholonu, 1976).
- Cricocephalus megastomum* Looss, 1902
Site of infection: Esophagus, stomach and small intestine.
Host and distribution: *C. mydas* from Brazil (Werneck & Silva, 2015; Binoti *et al.*, 2016), Costa Rica (Santoro *et al.*, 2006) and Cuba (Groschafft *et al.*, 1977); *E. imbricata* from Puerto Rico (Fischthal & Acholonu, 1976; Dyer *et al.*, 1995a).
- Cricocephalus resectus* Looss, 1902
Site of infection: Esophagus and stomach.
Host and distribution: *C. mydas* from Costa Rica (Santoro *et al.*, 2006).
- Cricocephalus vitallani* (Gupta, 1962)
(Synonym: *Neocricocephalus vitallani* Gupta, 1962)
Site of infection: Intestine.
Host and distribution: *C. mydas* from Caribbean Sea (Gupta, 1962).
- Desmogonius desmogonius* Stephens, 1911
Site of infection: Esophagus, stomach and intestine.
Host and distribution: *C. mydas* from "Caribbean waters" (Coil & Reid, 1965), Costa Rica (Santoro *et al.*, 2006) and Jamaica (Stephens, 1911).
- Diaschistorchis pandus* (Braun, 1901) Johnstone, 1913
Site of infection: Stomach and small intestine.
Host and distribution: *C. mydas* from Brazil (Werneck & Silva, 2015); *E. imbricata* from Brazil (Werneck *et al.*, 2015b), Cuba (Viguera, 1955) and Puerto Rico (Fischthal & Acholonu, 1976; Dyer *et al.*, 1995a).
- Metacetabulum invaginatum* Teixeira de Freitas & Lent 1938
Site of infection: Stomach and small intestine.
Host and distribution: *C. mydas* from Brazil (Teixeira de Freitas & Lent, 1938; Ruiz, 1946; Travassos *et al.*, 1969; Werneck & Silva, 2015; Binoti *et al.*, 2016); *E. imbricata* from Porto Rico (Fischthal & Acholonu, 1976).
- Pleurogonius americanus* Caballero, Zerecero & Grocott, 1955
[Synonym: *Pyelosomum americanum* (Caballero, Zerecero & Grocott, 1955)]
Site of infection: Intestine.
Host and distribution: *C. mydas* from Panamá (Caballero *et al.*, 1955).
- Pleurogonius laterouteus* Fischthal & Acholonu, 1976
Site of infection: Large intestine.
Host and distribution: *E. imbricata* from Porto Rico (Fischthal & Acholonu, 1976).
- Pleurogonius linearis* Looss, 1901
[Synonym: *Pyelosomum lineare* (Looss, 1901)]
Site of infection: Small and large intestine.
Host and distribution: *C. mydas* from Brazil (Ruiz, 1946; Travassos *et al.*, 1969; Werneck & Silva, 2015; Binoti *et al.*, 2016) and Costa Rica (Santoro *et al.*, 2006); *E. imbricata* from Costa Rica (Santoro *et al.*, 2015), México (Caballero & Zerecero, 1950) and Porto Rico (Fischthal & Acholonu, 1976).
- Pleurogonius longiusculus* Looss, 1901
[Synonym: *Pyelosomum longiusculum* (Looss, 1901)]
Site of infection: Esophagus, stomach, small and large intestine.
Host and distribution: *C. mydas* from Brazil (Ruiz, 1946; Travassos *et al.*, 1969; Werneck & Silva, 2015; Binoti *et al.*, 2016), Costa Rica (Santoro *et al.*, 2006) and Panamá (Caballero, 1954; Caballero *et al.*, 1955).
- Pleurogonius grocotti* Caballero, 1954
Site of infection: Intestine.
Host and distribution: *C. mydas* from Panamá (Caballero, 1954).
- Pleurogonius chelonii* Mehra, 1939
(Synonym: *Pleurogonius mehrai* Ruiz, 1946).
Site of infection: Intestine.
Host and distribution: *C. mydas* from Trinidad (The West Indies) (Gupta, 1961).
- Pleurogonius longibursatus* Viguera, 1955
[Synonym: *Pyelosomum longibursatum* (Viguera, 1955)].
Site of infection: Intestine.
Host and distribution: *E. imbricata* from Cuba (Viguera, 1955).

Pleurogonius puertoricensis Fischthal & Acholonu, 1976

Site of infection: Large intestine.

Host and distribution: *E. imbricata* from Puerto Rico (Fischthal & Acholonu, 1976).

Pleurogonius sindhii Mehra, 1939

[Synonym: *Pyelosomum sindhii* (Mehra, 1939)].

Site of infection: Intestine.

Host and distribution: *C. mydas* from Costa Rica (Santoro *et al.*, 2006) and Panamá (Caballero *et al.*, 1955).

Pleurogonius solidus Looss, 1901

[Synonym: *Pyelosomum solidum* (Looss, 1901); *Glyphicephalus solidus* Looss, 1901].

Site of infection: Intestine.

Host and distribution: *C. mydas* from Costa Rica (Santoro *et al.*, 2006) and *E. imbricata* from Cuba (Vigueras, 1955).

Pleurogonius tortugueroi Santoro, Greiner, Morales & Rodríguez-Ortiz, 2007.

Site of infection: Large intestine.

Host and distribution: *C. mydas* from Costa Rica (Santoro *et al.*, 2007).

Pleurogonius trigonocephalus (Rudolphi, 1809) Looss, 1901.

[Synonym: *Pyelosomum trigonocephalum* (Rudolphi, 1809), and *Monostoma trigonocephalum* Rudolphi, 1809].

Site of infection: Esophagus, stomach, small intestine, large intestine and liver.

Host and distribution: *C. mydas* from Brazil (Travassos *et al.*, 1969; Werneck & Silva, 2015); *E. imbricata* from Costa Rica (Santoro *et al.*, 2015), Cuba (Vigueras, 1955) and Puerto Rico (Fischthal & Acholonu, 1976).

Pleurogonius sp.

Site of infection: Intestine.

Host and distribution: *C. mydas* from British West Indies (Greiner *et al.*, 1980) and Costa Rica (Santoro *et al.*, 2006).

Pleurogonius lobatus (Looss, 1901)

(Synonym: *Glyphicephalus lobatus* Looss, 1901).

Site of infection: Stomach and small and large intestine.

Host and distribution: *C. mydas* from Brazil (Ruiz, 1946; Travassos *et al.*, 1969; Werneck & Silva, 2015; Binoti *et al.*, 2016), Costa Rica (Santoro *et al.*, 2006), Panamá (Caballero *et al.*, 1955), and Puerto Rico (Dyer *et al.*, 1991); *E. imbricata* from Costa Rica (Santoro *et al.*, 2015) and Puerto Rico (Fischthal & Acholonu, 1976); *L. olivacea* from México (Pérez-Ponce de León *et al.*, 1996).

Pleurogonius latus Fischthal & Acholonu 1976

(Synonym: *Glyphicephalus latus* Fischthal & Acholonu 1976).

Site of infection: Small intestine (occasionally stomach and large intestine).

Host and distribution: *E. imbricata* from Puerto Rico (Fischthal & Acholonu, 1976).

Pleurogonius stenobursatum (Fischthal & Acholonu, 1976) Blair, 2005

[Synonym: *Pyelosomum stenobursatum* (Fischthal & Acholonu, 1976) Pérez ponce de León & Brooks, 1995, and *Epibathra stenobursata* Fischthal & Acholonu 1976].

Site of infection: Large intestine.

Host and distribution: *E. imbricata* from Puerto Rico (Fischthal & Acholonu, 1976).

Pronocephalus obliquus Looss, 1899

(Synonym: *Pronocephalus mehrai* Chattopadhyaya, 1972).

Site of infection: Esophagus, stomach, small intestine, large intestine and liver.

Host and distribution: *C. mydas* from Brazil (Travassos *et al.*, 1969; Werneck & Silva, 2015; Binoti *et al.*, 2016) and Costa Rica (Santoro *et al.*, 2006); *E. imbricata* from Brazil (Werneck *et al.*, 2015b).

Pronocephalus trigonocephalus Looss, 1899

Site of infection: Esophagus, stomach and intestine.

Host and distribution: *C. caretta* from Brazil (Viana, 1924); *C. mydas* from Brazil (Ruiz, 1946; Travassos *et al.*, 1969; Binoti *et al.*, 2016) and Panamá (Caballero *et al.*, 1955).

Pyelosomum cochlear Looss, 1899

Site of infection: Urinary bladder.

Host and distribution: *C. mydas* from Brazil (Werneck & Silva, 2015; Binoti *et al.*, 2016), Costa Rica (Santoro *et al.*, 2006), Panamá (Caballero, 1954) and Puerto Rico (Dyer *et al.*, 1995b; Dyer *et al.*, 1991); *L. olivacea* from Brazil (Werneck *et al.*, 2015c).

Pyelosomum crassum (Looss, 1901) Ruiz, 1946

(Synonym: *Glyphicephalus crassa* Looss, 1901).

Site of infection: Small and large intestine.

Host and distribution: *C. mydas* from Brazil (Ruiz, 1946; Travassos *et al.*, 1969; Werneck & Silva, 2015), and *E. imbricata* from Costa Rica (Santoro *et al.*, 2015).

Pyelosomum posterorchis Oguro, 1936

Site of infection: Small intestine.

Host and distribution: *C. mydas* from Panamá (Caballero *et al.*, 1955); *E. imbricata* from Costa Rica (Santoro *et al.*, 2015) and Puerto Rico (Fischthal & Acholonu, 1976).

Pyelosomum renicapite (Leidy, 1856)

[Synonym: *Astororchis renicapite* (Leidy, 1856), *Monostomum nephrocephalum* Diesing, 1858, *Monostomum sphargidis* MacCallum, 1921 and *Pyelosomum longicaecum* Luhman, 1935]

Site of infection: Small and large intestine.

Host and distribution: *C. caretta* from Brazil (Werneck *et al.*, 2008a); *D. coriacea* from Brazil and Uruguay (Werneck *et al.*, 2012) and Puerto Rico (Dyer *et al.*, 1995b); *L. olivacea* from México (Pérez-Ponce de Leon *et al.*, 1996).

Rameshwarotrema uterocrescens Rao, 1975

Site of infection: Esophagus, stomach, small and large intestine.

Host and distribution: *C. mydas* from Brazil (Werneck & Silva, 2015) and Costa Rica (Santoro *et al.*, 2006); *E. imbricata* from Puerto Rico (Dyer *et al.*, 1995a).

Ruicephalus minutus (Ruiz, 1946) Skrjabin, 1955

(Synonym: *Pronocephalus minutus* Ruiz, 1946)

Site of infection: Stomach and small intestine.

Host and distribution: *C. mydas* from Brazil (Ruiz, 1946; Travassos *et al.*, 1969; Werneck & Silva, 2015; Binoti *et al.*, 2016).

Suborder Xiphidiata Olson Cribb, Tkach, Bray & Littlewood, 2003

Superfamily Gorgoderoidea Looss, 1901

Family Gorgoderidae Looss, 1899

Subfamily Anaporrhutinae Looss, 1901

Plesiochorus cymbiformis (Rudolphi, 1819) Looss, 1901

[Synonym: *Phyllodistomum cymbiforme* (Rudolphi, 1819; *Plesiochorus cymbiformis elongatus* Pigulevsky, 1953)]

Site of infection: Urinary bladder and small intestine.

Host and distribution: *C. mydas* from Brazil (Binoti *et al.*, 2016) and Panamá (Caballero, 1954); *E. imbricata* from Puerto Rico (Fischthal & Acholonu, 1976); *L. olivacea* from Costa Rica (Santoro & Morales, 2007).

Superfamily Microphalloidea Ward, 1901

Family Pachypsolidae Yamaguti, 1958

Pachypsolus irroratus (Rudolphi, 1819) Looss, 1902

[Synonym: *Pachypsolus branchus* Barker, 1922, *Pachypsolus lunatus* Looss, 1901, *Pachypsolus puertoricensis* Fischthal & Acholonu 1976 and *Pachypsolus ovalis* Linton, 1910].

Site of infection: Stomach and intestine.

Host and distribution: *C. mydas* from Panamá (Caballero *et al.*, 1955); *E. imbricata* from Porto Rico (Fischthal & Acholonu, 1976); *L. olivacea* from México (Pérez-Ponce de Leon *et al.*, 1996) and Costa Rica (Santoro & Morales, 2007).

Superfamily Plagiorchioidea Lühe, 1901

Family Brachycoeliidae Looss, 1899

Cymatocarpus solearis (Braun, 1899) Braun, 1901

(probably Synonym: *Cymatocarpus undulatus* Looss, 1899)

Site of infection: Esophagus, stomach and small intestine.

Host and distribution: *C. mydas* from Brazil (Werneck & Silva, 2015) and Mexico (Caballero, 1959); *E. imbricata* from Brazil (Werneck *et al.*, 2015b)

Family Plagiorchiidae Lühe, 1901

Enodiotrema megachondrus (Looss, 1899) Looss, 1901

Site of infection: Small intestine.

Host and distribution: *C. mydas* from Brazil (Werneck *et al.*, 2016a); *E. imbricata* from Cuba (Groschaft *et al.*, 1977); *L. olivacea* from Costa Rica (Santoro & Morales, 2007) and México (Pérez-Ponce de Leon *et al.*, 1996; Vivaldo *et al.*, 2006).

Enodiotrema reductum Looss, 1901

Site of infection: Small intestine.

Host and distribution: *C. mydas* from Panamá (Caballero, 1954); *E. imbricata* from Costa Rica (Santoro *et al.*, 2015) and Puerto Rico (Fischthal & Acholonu, 1976; Dyer *et al.*, 1995a).

Family Styphlotrematidae Baer, 1924

Styphlotrema solitaria (Looss, 1899) Odhner, 1911

Site of infection: Esophagus, stomach and small intestine.

Host and distribution: *E. imbricata* from Brazil (Werneck & Silva, 2012), Cuba (Groschaft *et al.*, 1977) and Puerto Rico (Fischthal & Acholonu, 1976).

Family Telorchiidae Looss, 1899

Orchidasma amphiorchis (Braun, 1899)

Site of infection: Stomach and small intestine.

Host and distribution: *C. caretta* from Argentina (Boero & Led, 1974) and Brazil (Werneck *et al.*, 2008a); *C. mydas* from Brazil (Teixeira de Freitas & Lent, 1938; Travassos *et al.*, 1969; Binoti *et al.*, 2016), México (Caballero & Zerecero, 1950; Caballero, 1962); Panama (Caballero *et al.*, 1955), and Peru (Tantalean *et al.*, 1992); *E. imbricata* from Puerto Rico (Fischthal & Acholonu, 1976); *L. olivacea* from México (Pérez-Ponce de Leon *et al.*, 1996).

Order Diplostomida Olson, Cribb, Tkach, Bray & Littlewood, 2003

Suborder Diplostomata Olson, Cribb, Tkach, Bray & Littlewood, 2003

Superfamily Schistosomatoidea Stiles & Hassall, 1898

Family Spirorchiidae Stunkard, 1921

Amphiorchis amphiorchis Price, 1934

Site of infection: Blood vessels of large intestine.

Host and distribution: *E. imbricata* from Puerto Rico (Fischthal & Acholonu, 1976).

Amphiorchis caborojoensis Fischthal & Acholonu, 1976

Site of infection: Blood vessels of lung, liver, small intestine, body wash and heart.

Host and distribution: *E. imbricata* from Puerto Rico (Dyer *et al.*, 1995a; Fischthal & Acholonu, 1976) and Brazil (Werneck *et al.*, 2008b; Dutra *et al.*, 2012; Werneck *et al.*, 2015b).

Amphiorchis indicus Mehrotra, 1973

Site of infection: Esophagus, stomach, small intestine, large intestine and liver.

Host and distribution: *C. mydas* from Brazil (Werneck & Silva, 2013; Werneck & Silva, 2015).

Amphiorchis solus (Simha & Chattopadhyaya, 1970) Platt, 2002 (Synonym: *Squaroacetabulum solus* Simha & Chattopadhyaya, 1970)

Site of infection: Intestine and heart.

Host and distribution: *C. mydas* from Brazil (Werneck *et al.*, 2011; Werneck & Medeiros, 2016) and Costa Rica (Santoro *et al.*, 2006).

Carettacola stunkardi (Martin & Bamberger, 1952)

[Synonym: *Haemoxenicon stunkardi* Martin & Bamberger, 1951, *Carettacola chelonenecon* (Martin & Bamberger, 1952), and *Haemoxenicon chelonenecon* Martin & Bamberger, 1952)].

Site of infection: Blood vessels of urinary bladder, heart, body wash and liver.

Host and distribution: *C. mydas* from Brazil (Werneck *et al.*, 2013) and Panamá (Caballero *et al.*, 1955); *E. imbricata* from Brazil (Werneck *et al.*, 2008b; Werneck *et al.*, 2015b) and Costa Rica (Santoro *et al.*, 2015).

Hapalotrema postorchis Rao, 1976

Site of infection: Great vessels and heart.

Host and distribution: *C. mydas* from Brazil (Werneck *et al.*, 2015d) and Costa Rica (Santoro *et al.*, 2006); *E. imbricata* from Brazil (Werneck *et al.*, 2014).

Hapalotrema synorchis Luhman, 1935

Site of infection: Heart.

Host and distribution: *E. imbricata* from Puerto Rico (Fischthal & Acholonu, 1976).

Learedius learedi Price, 1934

Site of infection: Esophagus, stomach, small and large intestine, liver, gall bladder, heart, lung, body wash, kidney.

Host and distribution: *C. mydas* from Bermuda (Rand & Wiles, 1985), Brazil (Werneck *et al.*, 2006; Werneck & Silva, 2015; Binoti *et al.*, 2016), British West Indies (Greiner *et al.*, 1980), Costa Rica (Santoro *et al.*, 2006), Panama (Caballero *et al.*, 1955), Puerto Rico (Dyer *et al.*, 1991), México (Cordeiro-Tapia *et al.*, 2004; Inohuye-Rivera *et al.*, 2004); *E. imbricata* from Puerto Rico (Dyer *et al.*, 1995a).

Learedius orientalis Mehra, 1939

Site of infection: Heart.

Host and distribution: *C. mydas* from Puerto Rico (Dyer *et al.*, 1995a); *E. imbricata* from Puerto Rico (Fischthal & Acholonu, 1976).

Monticellius indicum Mehra, 1939

Site of infection: Heart.

Host and distribution: *C. mydas* from Brazil (Werneck *et al.*, 2008c; Werneck & Silva, 2015) and Costa Rica (Santoro *et al.*, 2006; Santoro *et al.*, 2009b); *E. imbricata* from Brazil (Werneck *et al.*, 2015e).

Neospirochis schistosomatoides Price 1934

Site of infection: Heart chambers and Aorta.

Host and distribution: *C. mydas* from Bermuda (Rand & Wiles, 1985) and Brazil (Werneck *et al.*, 2016b).

Neospirochis sp.

Site of infection: Small intestine, liver, heart, lung, body wash.

Host and distribution: *C. mydas* from Brazil (Werneck & Silva, 2015).

Phylum Nematoda Rudolphi, 1808

Class Chromadorea Inglis, 1983

Subclass Chromadoria Adamson, 1987

Order Rhabditida Chitwood, 1933

Suborder Spirurina Railliet & Henry, 1915

Infraorder Ascaridomorpha De Ley & Blaxter 2002

Superfamily Ascaridoidea Baird, 1853

Family Anisakidae Skrjabin & Karokhin, 1945

Sulcascaaris sulcata (Rudolphi, 1819)

Site of infection: Esophagus, stomach and small intestine.

Host and distribution: *C. caretta* from Brazil (Werneck *et al.*, 2008a) and Uruguay (Lent & Teixeira de Freitas, 1948); *C. mydas* from Brazil (Teixeira de Freitas & Lent, 1946).

Anisakis larvae

Site of infection: Body wash.

Host and distribution: *E. imbricata* from (Werneck *et al.*, 2015b).

Superfamily Cosmocercoidea Travassos, 1935

Family Kathlaniidae Travassos, 1918

Kathlania leptura (Rudolphi, 1819)

Site of infection: Small and large intestine.

Host and distribution: *C. caretta* from Brazil (Werneck *et al.*, 2008a); *C. mydas* from Brazil (Travassos, 1918).

Tonaudia freitasi Vicente & Santos, 1968.

Site of infection: Stomach.

Host and distribution: *C. mydas* from Brazil (Vicente & Santos, 1968).

Nematode larvae

Site of infection: Esophagus, stomach, small and large intestine, liver.

Host and distribution: *C. mydas* from Brazil (Werneck & Silva, 2015).

Host-Parasite list

Chelonia mydas

Digeneans

Family Calycodidae

- *Calycodes anthos*
-

Family Rhytidodidae

- *Rhytidodes gelatinosus*
- *Rhytidodoides intestinalis*
- *Rhytidodoides similis*

Family Cladorchiidae

Schizamphistomum erratum

Schizamphistomum scleroporium

Schizamphistomoides spinulosum

Family Microscaphidiidae

- *Angiodictyum longum*
- *Angiodictyum parallelum*
- *Deuterobaris chelonei*
- *Deuterobaris intestinalis*
- *Deuterobaris proteus*
- *Microscaphidium aberrans*
- *Microscaphidium reticulare*
- *Microscaphidium warui*
- *Neoctangium travassosi*
- *Octangium hyphalum*
- *Octangium sagitta*
- *Polyangium linguatula*

Family Pronocephalidae

- *Adenogaster serialis*
- *Charaxicephaloides polyorchis*
- *Charaxicephalus robustus*
- *Cricocephalus albus*
- *Cricocephalus megastomum*
- *Cricocephalus resectus*
- *Cricocephalus vitallani*
- *Desmogonius desmogonius*
- *Diaschistorchis pandus*
- *Metacetabulum invaginatum*
- *Pleurogonius americanus*
- *Pleurogonius chelonii*
- *Pleurogonius grocottii*
- *Pleurogonius linearis*
- *Pleurogonius longiusculus*
- *Pleurogonius sindhii*

- *Pleurogonius solidus*
- *Pleurogonius tortugueroi*
- *Pleurogonius trigonocephalus*
- *Pleurogonius lobatus*
- *Pronocephalus obliquus*
- *Pronocephalus trigonocephalus*
- *Pyelosomum cochlear*
- *Pyelosomum crassum*
- *Pyelosomum posterorchis*
- *Rameshwarotrema uterocrescens*
- *Ruicephalus minutus*

Family Gorgoderidae

- *Plesiochorus cymbiformis*

Family Pachypsolidae

- *Pachypsolus irroratus*

Family Brachycoeliidae

- *Cymatocarpus solearis*

Family Plagiorchiidae

- *Enodiotrema reductum*

Family Telorchidae

- *Orchidasma amphiorchis*

Family Spirorchidae

- *Amphiorchis indicus*
- *Amphiorchis solus*
- *Carettacola stunkardi*
- *Hapalotrema postorchis*
- *Learedius learedi*
- *Learedius orientalis*
- *Monticellius indicum*
- *Neospororchis schistosomatoides*

Nematodes

Family Anisakidae

- *Sulcascaaris sulcata*

Family Kathlaniidae

- *Kathlania leptura*.
- *Tonaudia freitasi*

Eretmochelys imbricata

Digeneans

Family Calycodidae

- *Calycodes caborojoensis*

Familia Rhytidodidae

- *Rhytidodes gelatinosus*

Family Cladorchiidae

- *Schizamphistomum scleroporum*

Family Microscaphidiidae

- *Angiodictyum anteroporum*
- *Angiodictyum mooreae*
- *Angiodictyum parallelum*
- *Neoctangium travassosi*
- *Octangium sagitta*

Family Pronocephalidae

- *Adenogaster serialis*
- *Cricocephalus albus*
- *Cricocephalus americanus*
- *Cricocephalus megastomum*
- *Diaschistorchis pandus*
- *Metacetabulum invaginatum*
- *Pleurogonius laterouteus*
- *Pleurogonius latus*
- *Pleurogonius linearis*
- *Pleurogonius lobatus*
- *Pleurogonius longibursatus*
- *Pleurogonius puertoricensis*
- *Pleurogonius solidus*
- *Pleurogonius trigonocephalus*
- *Pleurogonius stenobursatum*
- *Pronocephalus obliquus*
- *Pyelosomum crassum*
- *Pyelosomum posterorchis*
- *Rameshwarotrema uterorescens*

Family Gorgoderidae

- *Plesiochorus cymbiformis*

Family Pachypsolidae

- *Pachypsolus irroratus*

Family Brachycoeliidae

- *Cymatocarpus solearis*

Family Plagiorchiidae

- *Enodiotrema megachondrus*
- *Enodiotrema reductum*

Family Styphlotrematidae

- *Styphlotrema solitaria*

Family Telorchiidae

- *Orchidasma amphiorchis*

Family Spirorchiidae

- *Amphiorchis amphiorchis*
- *Amphiorchis caborojoensis*
- *Carettacola stunkardi*
- *Hapalotrema postorchis*
- *Hapalotrema synorchis*
- *Learedius learedi*
- *Learedius orientalis*
- *Monticellius indicum*

Nematodes

Family Anisakidae

- *Anisakis larvae*

Lepidochelys olivacea

Digeneans

Family Calycodidae

- *Calycodes anthos*

Family Sclerodistomidae

- *Prosorchis psenopsis*

Family Pronocephalidae

- *Adenogaster serialis*
- *Pleurogonius lobatus*
- *Pyelosomum cochlear*
- *Pyelosomum renicapite*

Family Gongoderidae

- *Plesiochorus cymbiformis*

Family Pachypsolidae

- *Pachypsolus irroratus*

Family Plagiorchiidae

- *Enodiotrema megachondrus*

Family Telorchiidae

- *Orchidasma amphiorchis*

Caretta caretta

Aspidogastrids

Family Aspidogastridae

- *Lophotaspis vallei*

Digeneans

Family Calycodidae

- *Calycodes anthos*

Familia Rhytidodidae

- *Rhytidodes gelatinosus*

Family Pronocephalidae

- *Pronocephalus trigonocephalus*
- *Pyelosomum renicapite*

Family Telorchidae

- *Orchidasma amphiorchis*

Nematodes

Family Anisakidae

- *Sulcascaris sulcata*

Family Kathlaniidae

- *Kathlania leptura*

Dermochelys coriacea

Digeneans

Family Calycodidae

- *Calycodes anthos*

Family Pronocephalidae

- *Pyelosomum renicapite*

Discussion

This checklist demonstrates the occurrence of 79 taxa of helminth parasites. Seventy six represent the phylum Platyhelminthes (one from the order Aspidogastrida, 11 from the order Diplostomida and 64 from the order Plagiorchiida) represented by 14 families. Three represent the phylum Nematoda, Five species of sea turtles (*C. mydas*, *C. caretta*, *E. imbricata*, *L. olivacea* and *D. coriacea*) had records of helminths in the Neotropical region. The host with the largest number of records of parasites was *C. mydas* (records of only the genus were excluded from the calculation), representing 62 parasite species, followed by *E. imbricata* (42 species), *L. olivacea* (10 species), *C. caretta* (8 species) and *D. coriacea* (2 species). Overall helminths are reported in 12 countries and three are reported for the region in which

the present study took place (i.e. the Caribbean Sea).

Among the countries in which helminths are reported, Brazil, Costa Rica and Panamá have the largest number of records for *C. mydas*. *Lepidochelys olivacea* is the best studied sea turtle from Mexico, with eight records of parasite species. Porto Rico stands out for the large number of parasite species reported in *E. imbricata*, such as *C. caborjoensis*, *A. anteroporum*, *A. mooreae*, *P. laterouterus*, *P. puertoricensis*, *P. latus*, *P. sternobursatum*, *A. amphiorchis* and *H. synorchis*, which are only found in this country in this region.

Dermochelys coriacea had the fewest reports of parasites from the region with only *C. anthos* and *P. renicapite* being detected. This species of turtle has been studied the least of all marine turtles regarding its parasites around the world (see Werneck et al., 2012). This paper presents the largest compilation of data on helminth parasites in sea turtles in the Neotropical region published to date.

Acknowledgements

The authors wish to express their special thanks to Dr. Paula Baldassin, Daniel Clifton and the reviewers for critical reading and suggestions to improve the manuscript.

References

- ARAÚJO, T.L. (1941): Nota sobre um trematoide Aspidogastridae de tartaruga marinha. *Bol. Ind. Anim.*, 4: 3 – 4
- BINOTI, E., GOMES, M.C., DE CALAIS JUNIOR, A., WERNECK, M.R., MARTINS, I.V.F. BOELONI, J.N. (2016): Helminth fauna of *Chelonia mydas* (Linnaeus, 1758) in the south of the State of Espírito Santo and description of tissue injury. *Helminthologia*, 53(2): 195 – 199. DOI: 10.1515/helmin-2016-0012
- BLAIR, D. (1987): A revision of the subfamily Octangiinae (Platyhelminthes: Digenea: Microscaphidiidae) parasitic in marine turtles (Reptilia: Chelonia). *Aust. J. Zool.* 35: 75 – 92
- BOERO, J.J., LED, J.E. (1974): Parasites of the authochthonous fauna. *Rev. de Agronomía y de Veterinária*, 3: 16 – 17
- CABALLERO, C.E. (1954): Helmintos de la República de Panamá. X. Algunos tremátodos de *Chelonia mydas* (L.) tortuga marina comestible del Océano Pacífico del norte. *An. Inst. Biol. Univ. Nac. Auton. Mex.*, 8: 31 – 58
- CABALLERO, E.C. (1959): Tremátodos de las tortugas de México VII. Descripción de un trematodo digeneo que parasita a tortugas marinas comestibles del puerto de Acapulco, Guerrero. *An. Inst. Biol./ Univ. Nac. Auton. Mex.*, 33: 159 – 166
- CABALLERO, C.E. (1962): Tremátodos de las tortugas de México. X. Presencia de *Orchidasma amphirochis* (Braun, 1899) Looss, 1900 en una tortuga marina, *Chelone mydas* de las Costas del estado de Tamaulipas, México. *An. Inst. Biol./ Univ. Nac. Auton. Mex.*, 33: 47 – 55
- CABALLERO, C.E., ZERECERO M.C. (1950): Tremátodos de las tortugas de México VI. *Ver. Med. Vet. Y Paras.*, 9: 123 – 132

- CABALLERO, C.E., ZERECERO M.C., GROCOTT, R.G. (1955): Helmintos de la República de Panamá. XI. Tremátodos de *Chelonia mydas* (L) tortuga marina comestible del Océano Pacífico del norte. 2ª parte. *An. Inst. Biol./ Univ. Nac. Auton. Mex.*, 26: 149 – 191
- COIL, W.H., REID, W.A. (1965): *Desmogonius desmogonius* Stephens, 1911 (pronocephalidae: Digenea), a redescription with observations on egg filament formation. *Z. Parasitenkd.*, 25: 506 – 509
- CORDEIRO-TAPIA, A.A., GARDNER, S.C., ARELLANO-BLANCO, J., INOHUYE-RIVERA, R.B. (2004): *Learedius learedi* infection in black turtles (*Chelonia mydas agassizii*), Baja California sur, Mexico. *J. Parasitol.*, 90(3): 645 – 647. DOI: 10.1645/GE-165R
- DUTRA, G.H.P., SILVA, A.N., NASCIMENTO, C.L., WERNCK, M.R. (2012): Lesões macroscópicas e histopatológicas da infecção por helmintos da Family Spirorchiidae em *Eretmochelys imbricata* Linnaeus 1758 (Testudines, Cheloniidae): relato de um caso no litoral brasileiro. *Natural Resources, Aquidabã*, 2(1): 83 – 89. DOI: 10.6008/ESS2237-9290.2012.001.0006
- DYER, W.G., WILLIAMS, E.H.J., BUNKLEY-WILLIAMS, L. (1995c): *Angiodictyum mooreae* n.sp. (Digenea: Microscaphidiidae) and Other digeneans from na Atlântic Hawksbill turtle *Eretmochelys imbricata* from Puerto Rico. *J. Aquat. Anim. Health*, 7(1): 38 – 41. DOI: 10.1577/1548-8667(1995)007<0038:AMNSDM>2.3.CO;2
- DYER, W.G., WILLIAMS, E.H.J., BUNKLEY-WILLIAMS, L. (1995b): Digenea of the green turtle (*Chelonia mydas*) and the leatherback turtle (*Dermochelys coriacea*) from Puerto Rico. *Caribb. J. Sci.*, 31(3-4): 269 – 273
- DYER, W.G., WILLIAMS, E.H.J., BUNKLEY-WILLIAMS, L. (1991): Some Digeneans (Trematoda) of the green turtle, *Chelonia mydas* (Testudines: Cheloniidae) from Puerto Rico. *Proc. Helminthol. Soc. Wash.*, 58(2): 176 – 180
- DYER, W.G., WILLIAMS, E.H.J., BUNKLEY-WILLIAMS, L., MOORE, D. (1995a): Some digeneans (Trematoda) of the hawksbill turtle, *Eretmochelys imbricata imbricata* (Testudines: Cheloniidae) from Puerto Rico. *Proc. Helminthol. Soc. Wash.*, 62: 13 – 17
- FISCHTHAL, J.H., ACHOLONU, A.D. (1976) Some digenetic trematodes from the Atlantic hawksbill turtle, *Eretmochelys imbricate imbricata* (L) from Puerto Rico. *Proc. Helminthol. Soc. Wash.*, 43(2): 174 – 185
- GREINER, E.C., FORRESTER, D.J., JACOBSON, R. (1980): Helminths of mariculture-reared green turtles (*Chelonia mydas mydas*) from Grand Cayman, British West Indies. *Proc. Helminthol. Soc. Wash.*, 47(1): 142 – 144
- GROSCHAFT, J., TENORA, F. (1978): *Charaxicephaloides polyorchis* Gen Nov, Sp Nov (Trematoda: Charaxicephalinae) from *Chelonia mydas mydas* (Testudinata) in Cuba. *Vestnik Cs. Spol. Zool.*, 42(2): 108 – 111
- GROSCHAFT, J., OTERO, A.C., TENORA, F. (1977): Trematodes (Trematoda) from Cuban turtles *Chelonia mydas mydas* (L) and *Eretmochelys imbricata imbricata* (L) Testudinata-Cheloniidae). *Acta Univ. Agriculturae*, 25(4): 155 – 167
- GUPTA, S.P. (1961): On some trematodes from the intestine of the marine turtle, *Chelone mydas* from the Caribbean Sea. *Can. j. zool.*, 39: 293 – 298
- GUPTA, S.P. (1962): On two new trematodes from the intestine of the marine turtles, *Chelone mydas*, from the Caribbean Sea. *Indian j. Helminthol.*, 14(2): 71 – 76
- INOHUYE-RIVEIRA, R., CORDEIRO-TAPIA, A., ARELLANO-BLANCO, J., GARDNER, S.C. (2004): *Learedius learedi* Price, 1934 (Trematoda: Spirorchiidae), in black turtle (*Chelonia mydas agassizii*) hearts from Magdalena Bay, Baja California Sur, Mexico. *Comp. Parasitol.*, 71(1): 37 – 41. DOI: 10.1654/4113
- IUCN- International Union for Conservation of Nature (2015): Red List of Threatened Species. Version 2015-4. Retrieved March 21, from: www.iucnredlist.org
- LENT, H., TEIXEIRA DE FREITAS, J.F. (1948): Uma coleção de nematódeos, parasitos de vertebrados do Museu de História natural de Montevidéu. *Mem. Inst. Oswaldo Cruz*, 46(1), 1 – 71
- LUTZ, P.L., MUSICK, J.A. (1996): *The biology of sea turtles*. Washington, DC, CRC Press, 446 pp.
- MUNIZ-PEREIRA, L.C., VIEIRA, F.M., LUQUE, J.L. (2009): Checklist of helminth parasites of threatened vertebrate species from Brazil. *Zootaxa*, 2123, 1 – 45
- PÉREZ-PONCE DE LEÓN, G. P., GARCÍA-PRIETO, L., LEÓN-RÉGAGNON, V. (1996): Gastrointestinal Digenetic trematodes of olive ridley's turtle (*Lepidochelys olivacea*) from Oaxaca México. Taxonomy and infracommunity structure. *J. Helminthol. Soc. Wash.*, 63(1): 76 – 82
- RAND, T.G., WILES, M. (1985): Histopathology of infection by *Learedius learedi* Price, 1934 and *Neosporichis schistomatoides* Price, 1934 (Digenea: Spirorchiidae) in Wild Green turtles, *Chelonia mydas* L., from Bermuda. *J. Wildl. Dis.*, 21(4): 461 – 463
- RUIZ, J.M. (1943): *Neoctangium travassosi* gen. N., sp. (Trematoda: Paramphistomoidea) parasito de quelônio marinho. Chave dos gêneros da Family Microscaphidiidae Travassos, 1922. *Memo. Inst. Butantan*, 17: 35 – 45
- RUIZ, J.M. (1946): Pronocephalidae (Trematoda) Estudo das espécies brasileiras e revisão da Família. *Memo. Inst. Butantan*, 19: 249 – 372
- SANTORO, M., BRANDMAYR, O., GREINER, E.C., MORALES, J.A., RODRIGUES-ORTIZ, B. (2009a): Redescription of *Charaxicephaloides polyorchis* Groschaft ant Tenora 1978 (Digenea: Pronocephalidae) from the Green turtle *Chelonia mydas* in Costa Rica. *Helminthologia*, 46(2): 97 – 99. DOI: 10.2478/s11687-009-0019-6
- SANTORO, M., GREINER, E.C., MORALES, J.A., RODRIGUES-ORTIZ, B. (2007): A new pronocephalid, *Pleurogonius tortugueroi* n. sp. (Digenea), from the intestine of Green sea turtles (*Chelonia mydas*) in Costa Rica. *Parassitologia*, 49(1-2): 97 – 100
- SANTORO, M., GREINER, E.C., MORALES, J.A., RODRIGUES-ORTIZ, B. (2006): Digenetic trematode community in nesting green sea turtles (*Chelonia mydas*) from Tortuguero National Park, Costa Rica. *J. Parasitol.*, 92(6): 1202 – 1206. DOI: 10.1645/GE-866R.1
- SANTORO, M., GREINER, E.C., MORALES, J.A., RODRIGUES-ORTIZ, B. (2009b): Redescription of *Monticellius indicum* Mehra, 1939 (Digenea: Spirorchiidae) from the heart of Green Sea Turtle (*Chelonia mydas*) in Costa Rica. *Open Parasitol J.*, 3: 4 – 8
- SANTORO, M., MORALES, J.A. (2007): Some digenetic trematodes of

- the olive ridley sea turtle, *Lepidochelys olivacea* (Testudines, Cheloniidae) in Costa Rica. *Helminthologia*, 44(1): 25 – 28
- SANTORO, M., MORALES, J.A., BOLAÑOS, F., CHAVES, G., DE STEFANO, M. (2015): Helminths of hawksbill turtle (*Eretmochelys imbricata*) from the Pacific coast of Costa Rica. *Helminthologia*, 52(1): 67 – 70. DOI: 10.1515/helmin-2015-0012
- STEPHENS, J.W.W. (1911): *Desmogonius desmogonius*, a new species and genus of monostome flukes. *Ann. Trop. Med. Parasit.*, 5: 497 – 500
- TANTALEÁN, M., SARMIENTO, L., HUIZA, A. (1992): Digeneos (Trematoda) del Perú. *Boletín de Lima (Perú)*, 14: 47 – 84
- TEIXEIRA DE FREITAS, J.T., LENT, H. (1946): “*Porrocaecum sulcatum*” (Rudolphi, 1819). *Rev. Bras. Biol.*, 6(2): 235 – 238
- TEIXEIRA DE FREITAS, J.T., LENT, H. (1938): Sobre alguns trematodos parasitos de *Chelone mydas* (L.), principalmente Paramphistomoides. *Mem. Inst. Oswaldo Cruz*, 33(1): 79 – 97
- TRAVASSOS, L. (1918): Informações sobre a familia Kahlhanidae n. nom. *Rev. Bras. Biol.*, 2: 83 – 88
- TRAVASSOS, L., FREITAS, T., KOHN, A. (1969): Trematódeos do Brasil. *Mem. Inst. Oswaldo Cruz*, 67: 1 – 886
- VIANA, L. (1924): Tentativa de catalogação das espécies brasileiras de trematódeos. *Mem. Inst. Oswaldo Cruz*, 17: 95 – 227
- VICENTE, J.J., SANTOS, E. (1968): Terceira espécie do género “*Tonaudia*” Travassos, 1918, (Nemaloda, Kathlaniidae). *Atas Soc. Biol. Rio J.*, 12(2): 55 – 56
- VIGUERAS, P. (1955): Contribución al conocimiento de la fauna helmintológica cubana. *Mem. Soc. Cub. Hist. Nat.*, 22: 21 – 71
- VIVALDO, S.G., MÁRQUEZ, L.J.G., SARABIA, D.O., GARCÍA, J.L.V., CASA, F.C. (2009): Pathology in the Olive Ridley turtles (*Lepidochelys olivacea*) that arrived to the shores of Cuyutlan, Colima, Mexico. *Vet. Mex.*, 40: 69 – 78
- VIVALDO, S.G., SARABIA, D.O., SALAZAR, C.P., HERNÁNDEZ, A.G., LEZAMA, J.R. (2006): Identificación de parásitos y epibiontes de la tortuga Golfi na (*Lepidochelys olivacea*) que arribó a playas de Michoacán y Oaxaca, México. *Vet. Mex.*, 37(4): 431 – 440
- WERNECK, M.R., AMORIM, D.B., BLAIR, D. (2015c): Olive Ridley Sea Turtle *Lepidochelys olivacea* (Eschscholtz, 1829) from Brazil as a New Host of *Pyelosomum cochlear* Looss 1899 (Digenea: Pronocephalidae). *Comp. Parasitol.*, 82(1): 144 – 147. DOI: 10.1654/4739.1
- WERNECK, M.R., BALDASSIN, P., D’AZEREDO, F., TRAZZI, A., BERGER, B.C. (2014): The Hawksbill Sea Turtle *Eretmochelys imbricata* Linnaeus 1758 (Testudines, Cheloniidae) as new host of *Hapalotrema postorchis* Rao, 1976 (Digenea: Spirorchiidae). *Comp. Parasitol.*, 81(1): 75 – 78. DOI: <http://dx.doi.org/10.1654/4662.1>
- WERNECK, M.R., BALDASSIN, P., TORRES, F., TRAZZI, A., BERGUER, B. (2013): Report of *Carettacola stunkardi* (Martin & Bamberger, 1952) Dailey, Fast & Balazs, 1991 (Digenea: Spirorchiidae) infecting Green turtle *Chelonia mydas* Linnaeus, 1758 (Testudines, Cheloniidae) in Brazil. *Braz. J. Biol.*, 73(3): 675 – 676
- WERNECK, M.R., BECKER, J.H., GALLO, B.M.G., SILVA, R.J. (2006): *Learedius learedi* Price 1934 (Digenea, Spirorchiidae) in *Chelonia mydas* Linnaeus 1758 (Testudines, Cheloniidae) in Brazil: case report. *Arq. Bras. Med. Vet. Zootec.*, 58(4): 550 – 555
- WERNECK, M.R., BINOTI, E., MARTINS, I.V.F., CALAIS JÚNIOR, A., GOMES, M.C., BOELONI, J.N., TRAZZI, A., BERGUER, B.C. (2015a): Occurrence of *Rhytidodoides similis* Price, 1939 (Digenea: Rhytidodidae) and lesions due to spirorchiid eggs in a Green Turtle *Chelonia mydas* Linnaeus, 1758 (Testudines, Cheloniidae) from Brazil. *Comp. Parasitol.*, 82(2): 291 – 295. DOI: 10.1654/4747.1
- WERNECK, M.R., CONTI, L.M., BERGER, B. (2016a): Report of *Enodiotrema megachondrus* (Looss, 1899) Looss, 1901 (Digenea: Plagiorchiidae) in a green turtle *Chelonia mydas* Linnaeus, 1758 (Testudines, Cheloniidae) from Brazil. *Helminthologia*. Advance online publication. DOI 10.1515/helmin-2016-0019
- WERNECK, M.R., GALLO, B.M.G., LIMA, E.H.S.M., SILVA, R.J. (2011): Occurrence of *Amphiorchis solus* Simha & Chattopadhyaya, 1970 (Digenea, Spirorchiidae) infecting Green turtle *Chelonia mydas* Linnaeus, 1758 (Testudines, Cheloniidae) in Brazil. *Comp. Parasitol.*, 78(1): 200 – 203. DOI: 10.1654/4435.1
- WERNECK, M.R., GALLO, B.M. G., SILVA, R.J. (2008c): First report of *Monticellius indicum* Mehra, 1939 (Digenea: Spirorchiidae) infecting *Chelonia mydas* Linnaeus, 1758 (Testudines: Cheloniidae) from Brazil. *Braz. J. Biol.*, 68(2): 455 – 456
- WERNECK, M.R., GALLO, B.M.G., SILVA, R.J. (2008b): Spirorchiids (Digenea: Spirorchiidae) infecting a Hawksbill sea turtle *Eretmochelys imbricata* (Linnaeus 1758) from Brazil. *Arq. Bras. Med. Vet. Zootec.*, 60(3): 663 – 666
- WERNECK, M.R., LIMA, E.H.S.M., PIRES, T., SILVA, R.J. (2015b): Helminth parasites of the Juvenile Hawksbill Turtle *Eretmochelys imbricata* (Testudines, Cheloniidae) in Brazil. *J. Parasitol.*, 101(4): 130 – 134. DOI: <http://dx.doi.org/10.1645/13-479.1>
- WERNECK, M.R., MEDEIROS, L.S. (2016): Report of the fourth specimen of *Amphiorchis solus* (Simha & Chattopadhyaya, 1970) 46 years after the original description. *Helminthologia*. Advance online publication. DOI 10.1515/helmin-2016-0020
- WERNECK, M.R., SILVA, R.J. (2013): Occurrence of *Amphiorchis indicus* Gupta & Mehrotra, 1981 (Digenea, Spirorchiidae) infecting Green Turtle *Chelonia mydas* Linnaeus, 1758 (Testudines, Cheloniidae) in Brazil. *Braz. J. Biol.*, 73(1): 225 – 227
- WERNECK, M.R., SILVA, R.J. (2015): Some helminth parasites of juvenile green turtles *Chelonia mydas* (Testudines, Cheloniidae) in Brazil. *J. Parasitol.*, 101(6): 713 – 716. DOI: 10.1645/15-780
- WERNECK, M.R., SILVA, R.J. (2012): *Styphlotrema solitaria* Looss, 1899 (Digenea, Styphlotrematidae) infecting *Eretmochelys imbricata* (Linnaeus 1758) (Testudines, Cheloniidae) in Brazil. *Neotrop. Helminthol.*, 6(1): 121 – 125
- WERNECK, M.R., SOUZA, G., BERGUER, B.C. (2016b): *Neosporichis schistosomatoides* Price 1934 (Digenea: Spirorchiidae) infecting a Green Turtle, *Chelonia mydas* Linnaeus, 1758 (Testudines, Cheloniidae), from Brazil. *Helminthologia*, 53(1): 94 – 98. DOI: 10.1515/helmin-2015-0074
- WERNECK, M.R., SOUZA, G., BERGER, B.C., TRAZZI, A., RIBEIRO, R., SILVA, M.A., LEANDRO, H.J., CARVALHO, E.C.Q. (2015d): Pathologi-

- cal changes by *Hapalotrema postorchis* Rao 1976 (Digenea: Spirorchiidae) in a green turtle *Chelonia mydas* Linnaeus 1758 (Testudines, Cheloniidae) from Brazil. *Helminthologia*, 52(2): 148 – 154. DOI: 10.1515/helmin-2015-0027
- WERNECK, M.R., SOUZA, V.R., TRAZZI, A., BERGER, B.C. (2015e): *Moniticellius indicum* Mehra, 1939 (Digenea: Spirorchiidae) in a Hawksbill Turtle, *Eretmochelys imbricata* Linnaeus 1766 (Testudines, Cheloniidae) from Brazil. *Comp. Parasitol.*, 82(1): 155 – 157. DOI: 10.1654/4716.1
- WERNECK, M.R., THOMAZINI C.M., MORI, E.S., GONÇALVES, V.T. GALLO, B.M., SILVA, R.J. (2008a): Gastrointestinal helminth parasites of Loggerhead turtle *Caretta caretta* Linnaeus 1758 (Testudines, Cheloniidae) in Brazil. *PanamJAS*, 3(3): 351 – 354
- WERNECK, M.R., VERISSIMO, L., BALDASSIN, P., GAGLIARDI, F., TADASHI, E., VANDERLINDE, J. BAPTISTOTTE, C., MELO M.T.D., LIMA, E.H.S.M., GALLO, B.M. G., SILVA, R.J. (2012): Digenetic Trematodes of *Deremochelys coriacea* From the Southwestern Atlantic Ocean. *Marine Turtles Newsletter*, 132: 13 – 14
- WoRMS- World Register of Marine Species (2015): Retrieved November 25, from: <http://www.marinespecies.org/aphia.php?p=search>