

**UNIVERSIDADE ESTADUAL PAULISTA  
FACULDADE DE CIÊNCIAS AGRÁRIAS E VETERINÁRIAS  
CÂMPUS DE JABOTICABAL**

**INFLUENCE OF THE STARTING TIME OF PRESLAUGHTER  
FEED WITHDRAWAL ON WELFARE OF PIGS AND PORK  
QUALITY**

**Filipe Antônio Dalla Costa**  
Médico Veterinário

2016

**UNIVERSIDADE ESTADUAL PAULISTA  
FACULDADE DE CIÊNCIAS AGRÁRIAS E VETERINÁRIAS  
CÂMPUS DE JABOTICABAL**

**INFLUENCE OF THE STARTING TIME OF PRESLAUGHTER  
FEED WITHDRAWAL ON WELFARE OF PIGS AND PORK  
QUALITY**

**Filipe Antônio Dalla Costa**

**Orientador: Dr. Luigi Faucitano**

**Co-orientador: Prof. Dr. Mateus José Rodrigues Paranhos da Costa**

Dissertação apresentada à Faculdade de Ciências Agrárias e Veterinárias – Unesp, Campus de Jaboticabal, como parte das exigências para a obtenção do título de Mestre em Zootecnia.

**2016**

D144i Dalla Costa, Filipe Antônio  
Influence of the starting time of feed withdrawal interval  
preslaughter on welfare of pigs and pork quality / Filipe Antônio Dalla  
Costa. -- Jaboticabal, 2016  
xiii, 97 p.; 28 cm

Dissertação (mestrado) - Universidade Estadual Paulista,  
Faculdade de Ciências Agrárias e Veterinárias, 2016

Orientador: Luigi Faucitano

Banca examinadora: Cristiane Gonçalves Titto, Vivian Feddern  
Bibliografia

1. Behaviour. 2. Fasting. 3. Meat quality. 4. Pigs. 5. Stress. I. Título.  
II. Jaboticabal - Faculdade de Ciências Agrárias e Veterinárias.

CDU: 637.5:636.4

Ficha catalográfica elaborada pela Seção Técnica de Aquisição e Tratamento da Informação –  
Serviço Técnico de Biblioteca e Documentação - UNESP, Câmpus de Jaboticabal.

**Folha de Aprovação**  
**FORNECIDO PELA PÓS-GRADUAÇÃO**

## **DADOS CURRICULARES DO AUTOR**

**Filipe Antônio Dalla Costa**, nascido em 21 de novembro de 1990 na cidade de Concórdia-SC. Formado no ano de 2013 no curso de Medicina Veterinária pela Universidade do Estado de Santa Catarina, iniciou seus estudos em bem-estar animal e qualidade de carne durante estágios extracurriculares na graduação. Em 2014 ingressou no mestrado pelo Programa de Pós-Graduação em Zootecnia na Faculdade de Ciências Agrárias e Veterinárias da Universidade Estadual Paulista “Júlio de Mesquita Filho”, em Jaboticabal/SP, como bolsista do Conselho Nacional Científico e Tecnológico - CNPq e integrante do Grupo de Estudos e Pesquisas em Etologia e Ecologia Animal (Grupo ETCO).

“Os loucos que acham que podem mudar o mundo, são os que efetivamente o fazem.”

*Steve Jobs*

Aos meus pais, que se sacrificaram dia e noite em prol da minha educação para que eu pudesse estar aqui alcançando meus objetivos e sempre entenderam minhas decisões. A minha família, que mesmo eu estando distante, sempre me apoiou.

A todos os colaboradores rurais e da indústria que batalham todos os dias para melhorar as condições de manejo pré-abate.

**Dedico.**

## AGRADECIMENTOS

Primeiramente agradeço aos meus pais, Osmar e Tânia, que sempre me ensinaram todos os princípios morais de um homem, incentivando-me e me apoiando a estudar e atingir todos os meus objetivos mesmo que esses nos fizessem ficar distante. Pai, você todos os dias me dá um exemplo de homem honesto, trabalhador e dedicado à família. Mãe, muito obrigado por me aconselhar da melhor maneira sempre e dar força para seguir em frente sempre.

A todos os meus familiares que, embora saudosos, souberam compreender todos os momentos distantes e a importância disso.

Ao meu orientador, Dr. Luigi Faucitano, que aceitou me orientar e, mesmo à distância, confiou no nosso trabalho. Pelos conselhos e tempo dedicado à preparação deste trabalho. Pela oportunidade de fazer parte do seu grupo de pesquisa no Agri-food and Agriculture Canada (desde a graduação) e incentivar todos os estudantes a evoluírem na pesquisa científica.

Ao meu co-orientador, Professor Mateus Paranhos, por abrir as portas do Grupo ETCO onde vivi, amadureci e tive experiências incríveis trabalhando. Pela sua dedicação incansável ao trabalho do grupo e incentivos à educação e melhoria de todos os jovens que passam por aqui. Pela oportunidade de conhecer, ver, rever e aprender com grandes pesquisadores, que fazem a diferença no bem-estar animal: Temple Grandin, Donald Broom, Neville Gregory, Mohan Raj, Antonio Velarde, Antoni Dalmau, Karen Schwartzkopf-Genswein, Luigi Faucitano, Osmar Dalla Costa, Xavier Mantega, Márcia del Campo, Linda Keeling, Jens Jung.

Ao Dr. Osmar Dalla Costa, meu pai e conselheiro, que me orientou durante a minha caminhada profissional nesses dois anos de trabalho e me incentivou a produzir o melhor possível. Pelo seu trabalho e esforços para melhorar a suinocultura brasileira e condições de trabalho de todos os colaboradores da indústria.

Aos integrantes do exame de qualificação (Dra. Charli Ludtke e Professora Hirasilva Borba) e banca de defesa pela disponibilidade, atenção e valiosíssimas contribuições para melhoria do meu trabalho.

À Dra. Charli Ludtke, por permitir que fizesse parte da equipe STEPS, como

estagiário, onde realizei meu primeiro estágio na área de bem-estar animal e confirmei minhas aspirações profissionais e paixão pela área.

Aos meus companheiros do Grupo ETCO que me receberam de braços abertos desde o primeiro dia, sou grato pelas conquistas que tivemos, conhecimentos adquiridos, amizades, e esforços sem limites para melhorar as condições de produção animal. Agradeço especialmente àqueles que além de colegas de trabalho se tornaram amigos muito queridos e dividimos horas de dedicação à pesquisa: Steffan Edward e Monique Carvalhal.

Aos meus amigos que acompanharam toda essa maravilhosa etapa da minha vida de perto e de longe, tanto no Brasil quanto no exterior.

Obrigado meus excelentes professores por sempre me incentivarem a estudar e tornar minhas noites mais curtas com suas provas, questões difíceis e seus altos níveis de exigência, e por pegarem no meu pé por algum deslize mesmo que minúsculo esse fosse. Vocês são os responsáveis não só pela educação acadêmica, mas também pela formação de pessoas e, a cima disso, profissionais competentes que auxiliarão toda a cadeia produtiva agropecuária e a saúde humana, garantindo altos níveis de bem-estar animal, segurança na produção de alimentos, nutrição, sanidade animal e prevenções de enfermidades e doenças.

## SUMMARY

	<b>Page</b>
<b>Abstract</b> .....	iv
<b>Table List</b> .....	v
<b>Figure List</b> .....	vi
<b>Acronyms and Abbreviations List</b> .....	vii
<b>CHAPTER 1 – GENERAL CONSIDERATIONS</b> .....	1
1. INTRODUCTION.....	1
2. LITERATURE REVIEW.....	2
2.1. Animal welfare.....	2
2.2. General principles of stress.....	3
2.3. Behaviour.....	4
2.3.1. Normal and abnormal behaviour.....	4
2.3.2. Postural behaviour.....	5
2.3.3. Fighting behaviour.....	6
2.4. Physiology of stress response.....	8
2.4.1. Physiological parameters of stress.....	8
2.4.1.1. Cortisol.....	8
2.4.1.2. Lactate and Creatine kinase (CK).....	9
2.4.1.3. Heart and respiration rate.....	9
2.4.1.3. Body temperature.....	10
2.5. Animal losses.....	11
2.6. Carcass quality.....	12
2.6.1 Carcass yield.....	12
2.6.2. Skin lesions.....	15
2.7. Conversion of muscle into meat.....	15
2.8. Meat quality.....	16
2.8.1. Meat pH.....	17
2.8.2. Meat colour.....	18
2.8.3. Drip loss.....	20
2.9. Meat quality categories.....	21
2.10. The effects of preslaughter handling on animal welfare and meat quality..	23

2.10.1. Preslaughter feed withdrawal.....	23
2.10.2. Physiological parameters.....	24
2.11. Loading.....	25
2.11.1. Exiting the farm gate.....	26
2.11.2. Loading onto the truck.....	28
2.12. Transporting pigs.....	29
2.12.1. Vehicle design .....	29
2.12.2. Transport density.....	31
2.13. Unloading.....	32
2.14. Lairage.....	34
2.15. Handling pigs to slaughter.....	36
2.16. Stunning Methods.....	37
2.16.1. Electrical stunning.....	37
2.16.2. Carbon-dioxide (CO <sub>2</sub> ) stunning.....	38
2.17. Slaughter methods.....	38
REFERENCES.....	40
CHAPTER 2. EFFECTS OF APPLYING PRESLAUGHTER FEED WITHDRAWAL AT THE ABATTOIR ON BEHAVIOUR, BLOOD PARAMETERS AND MEAT QUALITY IN PIGS.....	70
CHAPTER 3. FINAL CONSIDERATIONS AND IMPLICATIONS.....	95


**Agriculture et Agroalimentaire Canada**

Direction générale de la science et de la technologie

 Centre de recherche et de développement de Sherbrooke  
 (CRDS)  
 2000, rue College  
 Sherbrooke QC J1M 0C8

 Courriel: [pauline.bilodeau@agr.gc.ca](mailto:pauline.bilodeau@agr.gc.ca)  
 Téléphone: 819 780-7118  
 Télécopieur: 819 564-5507

**Agriculture and Agri-Food Canada**

Science and Technology Branch

 Sherbrooke Research and Development Centre  
 (SRDC)  
 2000 College street  
 Sherbrooke QC J1M 0C8

 E-mail: [pauline.bilodeau@agr.gc.ca](mailto:pauline.bilodeau@agr.gc.ca)  
 Phone : 819 780-7118  
 Fax : 819 564-5507

Sherbrooke, 10 décembre 2015

Objet: Certification CIPA

Le comité institutionnel de protection des animaux (CIPA) du Centre de recherche et de développement de Sherbrooke (anciennement *Centre de recherche et de développement sur le bovin laitier et le porc*) certifie que le projet intitulé:

« *Influence de la mise à jeun avant l'abattage appliquée à la ferme ou à l'abattoir sur la perte de poids, la réponse comportementale, les caractéristiques du tractus gastro-intestinale et la variation de la qualité de la carcasse et de la viande de porc* »

a reçu l'approbation du comité institutionnel de protection des animaux lors de la réunion régulière du 12 février 2013 (projet #419).

Bien à vous,

 Mme Pauline Bilodeau,  
 Présidente du CIPA

 Sherbrooke, December 10<sup>th</sup>, 2015

Object: ACC Certification

This letter is to certify that the project entitled:

*"Influence of preslaughter feed withdrawal as either applied at the farm or in lairage on body weight losses, behavioral response, gastro-intestinal tract characteristics and carcass and meat quality variation in pigs"*

has received the approval of the Sherbrooke Research and Development Center (formerly *Dairy and Swine Research and Development Centre*) Animal Care Committee (ACC) at the meeting held February 12<sup>th</sup>, 2013 (project #419).

Sincerely,

 Mrs. Pauline Bilodeau,  
 President of the local Animal Care Committee

## **INFLUENCE OF THE STARTING TIME OF PRESLAUGHTER FEED WITHDRAWAL ON WELFARE OF PIGS AND PORK QUALITY**

**ABSTRACT** – The objective of this study was to evaluate the effects of starting time of feed withdrawal interval preslaughter at the farm (FARM) or at the plant (PLANT) on animal losses, behaviour during loading and lairage, blood parameters (lactate and creatine kinase) and meat quality traits (pH, colour and drip loss) in pigs. A total of 700 pigs was distributed into two treatments groups. The FARM group was fasted for 18 h before the transport plus 6 h before slaughter, including 2 h transport and 4 h lairage, and the PLANT group was fasted for 24 h (considering 2 h transport plus 22 h at lairage), totalizing 24 h of fasting time before slaughter for both groups. The treatments neither influenced transport losses nor blood parameters. However, PLANT pigs had longer fights ( $P < 0.05$ ) than FARM pigs at lairage and produced darker ( $P = 0.02$ ) and drier ( $P = 0.03$ ) loins. Based on these results, it is recommended apply feed withdrawal time before loading, which showed improvements to pigs' welfare, due to the lower number of fights during lairage at the slaughter plant, and produced better pork quality.

**Keywords:** animal welfare, animal losses, fasting, stress, preslaughter

## TABLE LIST

### CHAPTER 1

Table 1. Pork quality based on parameters of ultimate pH (pHu), drip loss (DL), subjective colour (JCS), and objective colour (L\*) (CORREA et al., 2007)

Table 2. Feed withdrawal interval applied in different countries (FAUCITANO, 2007; DALLA COSTA; DALLA COSTA, 2014a)

### CHAPTER 2

Table 1. Description of pig behaviors and interventions of the handler during loading (adapted from Weschenfelder et al., 2012)

Table 2. Effects of feed withdrawal as mostly applied on-farm (FARM) or at slaughter plant (PLANT) on the frequency of pig behaviours and interventions of handlers at loading

Table 3. Effects of feed withdrawal as mostly applied on-farm (FARM) or at slaughter plant (PLANT) on exsanguination blood parametersa

Table 4. Effects of feed withdrawal as applied on-farm or at the slaughter plant on meat quality traits as assessed in the Longissimus thoracis (LT), Semimembranosus (SM) and Adductor (AD) muscles

## FIGURE LIST

### CHAPTER 1

Figure 1. The coverage of animal welfare definitions (adapted from PARANHOS DA COSTA; OLIVEIRA and RUEDA, 2013).

Figure 2. Animal physiological reaction to a stressor (adapted from CARRAGHER; MATTHEWS, 1996).

Figure 3. Changes in glycogen, phosphocreatine (CP), ATP, and lactic acid concentrations during *post-mortem* metabolism (Weschenfelder, 2013).

Figure 4. Variation in salivary cortisol levels ( $\mu\text{g/dL}$ ) of pigs during preslaughter handling (adapted from DALLA COSTA et al., 2008).

### CHAPTER 2

Figure 1. Behaviour (back transformed least square means and confidence interval) at the beginning of lairage (after 20 min shower) of pigs being fasted for 24 h starting at the farm or the slaughter plant. a) percentage of pigs lying down, b) number of fights, c) total duration of fights. † $P < 0.10$ , \* $P < 0.05$ , \*\* $P < 0.01$  difference between treatments within lairage periods. For PLANT pigs only, black bars lacking a common superscript letter differ ( $P < 0.05$ )

Figure 2. Behaviour (back transformed least square means and confidence interval) at the end of lairage (from 120 min to the beginning of the last shower) of pigs being fasted for 24 h starting at the farm or the slaughter plant. a) percentage of pigs lying down, b) number of fights, c) total duration of fights. \* $P < 0.05$ , \*\*\* $P < 0.001$  difference between treatments within lairage periods

## ACRONYMS AND ABBREVIATIONS LIST

ACTH: Adrenocorticotropic Hormone

ATP: Adenosine Triphosphate

CK: Creatine Kinase

CO<sub>2</sub>: Carbon Dioxide

CRH: Corticotropin-releasing Hormone

DFD: Dark, Firm and Dry

DOA: Dead-on-arrival

JCS: Japanese Colour Scale

NAI: Non-ambulatory, Injured

NANI: Non-ambulatory, Non-injured

PC: Phosphocreatine

PFN: Pale, Firm, Non-exudative

pH<sub>u</sub>: ultimate pH

PSE: Pale, Soft, Exudative

RFN: Red, Firm, Dry

RSE: Reddish-pink, Firm, exudative

## CHAPTER 1 – General Considerations

### 1. Introduction

Feed withdrawal is a recommended practice for on-farm preparation before pigs' transport. This practice consists in denying pigs the access to feed, while keeping free access to water (DALLA COSTA; DALLA COSTA, 2014a). The appropriate application of feed withdrawal benefits both producers and slaughterhouse managers, as it helps to reduce transport losses (GUÀRDIA; GISPERT; DIESTRE, 1996; STEWART et al., 2008) and carcass contamination due to lower risk of gut contents spillage during carcass evisceration (SAUCIER et al., 2007), besides improving pork quality (FAUCITANO; CHEVILLON; ELLIS, 2010a). However, Aalhus (1992) and Viau and Champagne (1998) reported that this practice is not always consistently applied by the producers who prefer keeping pigs on feed until departure and starting fasting interval from farm departure to the slaughterhouse, as confirmed by a survey conducted at swine farms in Quebec, reporting that only 15% of pigs were fasted before transport (VIAU; CHAMPAGNE, 1998). This practice implies that to respect the fasting interval needed to obtain empty stomachs at slaughter a longer lairage time must be applied at the slaughterhouse.

The causes of the misapplication of feed withdrawal on-farm are associated to the split marketing production system which is very common at finishing farms in North America and in some European countries, such as France and Spain. It implies the removal of the heaviest 25 to 50% of pigs from a pen to market them 1-2 weeks earlier than the other penmates. The advantage of this practice is the reduction of production costs and the opportunity to ship batches with a more uniform market weight to the abattoir (SCROGGS et al., 2002). Whereas, the disadvantage is the risk of slower growth rate of pigs remaining in the pen that were also withdrawn of feed. A solution can be the use of shipping rooms where the heaviest pigs can be transferred ahead of time before loading, with local feed withdrawal drawing the feed there (VIAU; CHAMPAGNE, 1998).

The objectives of this study were to assess the effects of starting time of

preslaughter feed withdrawal at the farm or at the plant on animal losses during transport, as well as behaviour during loading and lairage, besides evaluating blood parameters, carcass traits and pork quality.

## 2. Literature review

### 2.1. Animal welfare

According to Hughes (1976), animal welfare is a state of complete mental and physical health, where there is harmony between animal and environment. Similarly, Broom (1986) defined welfare of an individual as its state as regards its attempts to cope with its environment. While the Broom's definition refers to how well or how badly coping attempts succeed, the Hughes's definition refers only to situations when animals succeed in its attempts to cope with the environment. Therefore, when animal welfare can be measured from a state of very poor to very good (BROOM, 1991), the Broom's definition which is broader (Figure 1) is considered as more useful by the scientific community.




Figure 1. The coverage of animal welfare definitions (adapted from PARANHOS DA COSTA; OLIVEIRA and RUEDA, 2013).

The industry is giving more attention to animal welfare not only because of opportunities offered by new markets, but also due to financial losses related to the production of meat with quality defects, such as PSE (pale, soft, exudative) and DFD (dark, firm, dry) pork (FRASER; BROOM, 1997) .

## 2.2. General principles of stress

Stress is defined as an internal organism's state of perturbation on physiological homeostasis or psychological well-being (COMMITTEE ON ALLEVIATION OF DISTRESS IN LABORATORY ANIMALS, 2008). It occurs when the organism fails in coping with environmental changes, resulting in over taxation of the control systems and, consequently, compromised fitness (BROOM, 1988). In response to stress, the organism makes some significant biological changes as showed by the behavioural and physiological adaptive responses. When stress results in a "biological cost", it is considered as distress (MOBERG, 2000). Distress occurs when the adaptation processes fail and the organism cannot return to its physiological and/or psychological homeostasis (MOBERG, 1987; CARSTENS; MOBERG, 2000). Strong challenges may overload the pigs' capacity to cope with the stressor and result in death (HEMSWORTH et al., 2015).

Behaviour changes are the first and most efficient response to a stressor for the animal (VON HOLST, 1998; MOBERG, 2000). However, under given circumstances, behavioural changes alone may not be enough or appropriate to cope with a stressor. In this case, the stress response done by the autonomic nervous and neuroendocrine systems is intensified. The stress response is multifactorial and may be influenced by a combination of several factors, such as the type and duration of a stress event, animal age, genetics, coping patterns and social status of the animals (VON HOLST, 1998; SALAK-JOHNSON; McGLONE, 2007). The physiological reaction to a stressor is shown in Figure 2.




Figure 2. Animal physiological reaction to a stressor (adapted from CARRAGHER; MATTHEWS, 1996).

### 2.3. Behaviour

In response to a potential stressful stimulus, animals develop some adaptation mechanism to keep the homeostatic balance (GRANDIN, 1998; MOBERG, 2000). First, the animals show behaviour reactions to the environment (MOBERG, 2000). The behavioral reaction is quick, especially in acute stressful situations involving fear (DE PASSILLÉ et al., 1995). For example, increased vocalisation and greater number of slips and falls are behavioural signs of the pig response to stressful situations, e.g. rough handling during preslaughter procedures (WARRISS et al., 1994; GRANDIN, 1998a; CORREA et al., 2010; DOKMANOVIC et al., 2014).

#### 2.3.1. Normal and abnormal behaviour

Since it measures animal own decision-making process using non-invasive methods, animal behaviour is an important tool for animal welfare assessments (DAWKINS, 2004).

Naturally, pigs can interact with their environment through their sensory

abilities, such as hearing, vision, touch, olfaction and taste (HELD et al., 2009). This interaction results in behaviour patterns, such as sexual receptiveness and spatial intrusion (HELD et al., 2009). Normal behavior is defined as the collection of behaviours animals have developed during the evolutionary process and would express in a natural environment (LIDFORS; BERG; ALGERS, 2005). Basically, pigs perform three types of natural behaviours: 1) behaviour they want to express (e.g. nest building at the end of gestation); 2) behaviour performed only under specific conditions (e.g. panting or shivering under thermal stress); and, 3) behaviour they do not want to express (e.g. separation from a group; OLSSON; WÜRBEL; MENCH, 2011).

On the other hand, when normal behaviour is hindered, the abnormal behaviour and stress may be induced (LIDFORS et al., 2005). Abnormal behaviour is defined as an uncommon response to a specific combination between motivational elements and stimulus (JENSEN, 1986) and indicates a failure in the animal attempt to cope with the environment (JENSEN, 1986; LIDFORS et al., 2005). According to Jensen (1986) and Mormède (2008), abnormal behaviours in pigs may be classified as: 1) apathy or boredom, indicating a non-adaptation to a novel environment; 2) aggression, induced by competition for food and social hierarchy; 3) stereotypies, which is an unvaried sequence of movements repeated with no obvious purpose due to poor environmental stimulation and frustration; and 4) cannibalism (i.e. tail biting), induced by poor environmental stimulation.

### **2.3.2. Postural behaviour**

Observation of postural behaviour is a method to determine stress levels imposed to animals during preslaughter handling (EWBANK; BRYANT, 1972; MOSS, 1978). Pigs can seat or lie down during lairage in order to relax or rest due to physical fatigue (BRANDT; AASLYNG, 2015). The shorter time to lie down at lairage may be an indicative of physical fatigue caused by the additive stress effect during preslaughter handling (BRANDT et al., 2013). Usually, pigs seat down within 1 hour of lairage (MOSS, 1978; BRANDT et al., 2013). There is an effect of feed withdrawal on behavior during lairage. Brown et al. (1999) observed a greater activity of fighting

and drinking on arrival at the lairage pen in pigs fasted for longer time (18 vs. < 1 h). Fernandez et al. (1995) reported that fed pigs spent significantly more time lying down (10 min) than 24 h fasted ones (5 min). The increased proportion of pigs lying down at lairage is associated with the reduction in fighting behaviour, as shown by previous studies (MOSS, 1978; FERNANDEZ et al., 1995; BROWN et al., 1999). Independently of the reason, in order to relax or rest, pigs lying seem to be a welfare advantage, because it reduces the number of fights resulting in lower risk of skin damage (BRANDT et al., 2013).

Rabaste et al. (2007) found an effect of group size on lying down behaviour, where pigs kept in smaller groups (10 vs. 30 pigs) tended to spend more time lying down. The environmental temperature during the lairage period influences lying behaviour (FREAQUEZA et al., 1998; EKESBEO, 2011). Pigs lie close together during cold stress, whereas when it is warm, they lie without direct body contact (EKESBEO, 2011). Fraqueza et al. (1998) observed a greater percentage of pigs lying down at 35°C for 0.5 or 3 hours than at 20°C, which is correspondent to the thermoregulatory behaviour during attempts to increase body heat conduction to the floor. Weeding et al. (1993) also verified a larger number of pigs lying down for longer periods when kept in lairage above 25°C and not cooled by showering.

### **2.3.3. Fighting behaviour**

Fighting is a normal social behaviour in group of pigs, and is easily observed when unknown pigs are mixed into a group (RHIM et al., 2015). Usually, fighting behaviour occur in order to establish a clear dominance hierarchy (EWBANK, 1976). Situations such as mixing groups and competition by resources (i.e. food shortage, high density and restricted feed space) may increase fights (EWBANK; BRYANT, 1972; LØVENDAHL et al., 2005). Aggression can also occur due to fearful stress, and as a spontaneous response to sudden pain (GREGORY, 2004). During preslaughter handling, no mixing is a practical recommendation. However, under commercial condition, mixing is a common practice, which causes intensive aggression in commercial housing systems for pigs (RHIM et al., 2015). Generally, pigs can be mixed at the farm during fasting in the expedition room or loading into the

truck, or even at the slaughter plant at the lairage. The fighting behaviour sometimes results in damages to health and induces physiological stress responses (MEESE; EWBANK, 1972; EWBANK; BRYANT, 1972; AREY; EDWARD, 1998; PITTS et al., 2000). Physical activity in lairage is the most potent factor stimulating muscle glycogenolysis in pigs. Brandt et al (2013) found a significant correlation between creatine kinase at exsanguination and skin damage score. Creatine kinase measurement can be an indicator of skin or muscle damage and it can be obtained from just one of the measurements (BRANDT et al., 2013). Problems associated to fighting include high bruises scores on carcass what affects meat quality (STOOKEY; GONYOU, 1994; WARRISS, 1996; GISPERT et al. 2000), and represent a significant economic cost to pork chain and serious animal welfare issue (AYO et al., 1998; O'CONNELL; BEATTIE, 1999).

Most of the fights occur during the first 60 min of lairage (MOSS, 1978; GEVERINK et al., 1996; FRAQUEZA et al., 1998). In mixed groups, Meese and Ewbank (1973) suggested that fights are practically eliminated after 24 hours and the social order is defined within 48 hours. The fasting condition affects fighting behaviour in pigs. Brown et al. (1999) found a greater number of fights on arrival at the lairage pen in pigs fasted for longer time (18 vs. < 1 h). Fernandez et al. (1995) reported a greater activity in fasted pigs (24 vs. 0 h) than fed ones. However, the authors did not found difference between treatments for the latency before the first aggressive interaction. Pigs fight more when kept in lairage for longer periods (overnight to > 24 h vs. 3 h) as a result of mixing and frustration due feed withdrawal (MURRAY; JONES, 1994; WARRISS et al., 1998b; GUÀRDIA et al., 2009).

The skin bruises are direct associated to stress suffered by pigs, as demonstrated by higher concentrations of cortisol and creatine phosphokinase in blood after slaughter (WARRISS; BROWN, 1985; WARRISS, 1996). Fernandez et al. (1995) suggested that the activation of the hypothalamic-pituitary-adrenal axis in response to both novel environment and aggressive encounters is not influenced by food deprivation. However, literature is scarce on the effect of feed withdrawal on sympatho-adrenal reactivity to stress.

Fighting behaviour can be direct or indirect measured by behaviour observations or counting the lesions on the skin or carcass, respectively. However,

because observing behaviour is time consuming and impractical, counting skin or carcass lesion is more efficient, quicker and easier to determine fighting in pigs.

## **2.4. Physiology of the stress response**

The autonomic system (i.e. the sympatho-adrenal medullary) and the hypothalamo-pituitary adrenal axis are activated in order to protect the biological system and to cope with the stressor through an increased synthesis of catecholamines and glucocorticoids (ULRICH-LAI; HERMAN, 2009; TURNER et al. 2012). The first activation occurs through a rapid response to stress, so called "alarm", "emergency syndrome" or also "fight-or-flight" (CANNON, 1929; MOBERG, 2000).

The stressful stimulus, external or internal, is transmitted to the hypothalamus which secretes the corticotropin-releasing hormone (CRH). This hormone is carried by the hypothalamo-hypophyseal portal system to the anterior lobe of the pituitary gland which is responsible for the synthesis and secretion of adrenocorticotrophic hormone (ACTH). Its principal effects are stimulation of the cortex of the adrenal glands, resulting in synthesis and release of glucocorticoids (cortisol) and mineralocorticoids (adrenaline and noradrenaline) into blood (MATTERI; CARROLL; DYER, 2000; ULRICH-LAI; HERMAN, 2009).

### **2.4.1. Physiological parameters of stress**

#### **2.4.1.1. Cortisol**

Cortisol is the major glucocorticoid used for the assessment of pigs' welfare. Cortisol levels can be a blood indicator of long-term psychological stress response (fear, alert) rising in less than 30 min in response to a stressor and returning to the basal level in 3 h (NYBERG et al., 1988; BRADSHAW et al., 1996; EKKEL et al., 1997; RUIS et al., 2001; MORMÈDE, 2008). Cortisol is secreted by the cortex of adrenal gland following the activation of the hypothalamic–pituitary–adrenal (HPA) axis in response to a stressor (WILLIAMS, 1983; NORMAN; LITWACK, 1997;

BOLANDER, 2004). However, a particular attention must be paid in the interpretation of blood cortisol variation, as its blood levels follow a circadian rhythm, reaching a peak in the early morning and declining steadily during daylight hours and reaching the lowest level in the evening (RUIS et al., 1997; HAY et al., 2000; MORMÈDE, 2008).

#### **2.4.1.2. Lactate and creatine kinase**

Lactate concentration in blood is a very short-term indicator of physical stress which reaches the peak within 4 min and returns to basal level in 2 h (ANDERSON, 2010). Lactate is produced from pyruvate during normal aerobic metabolism of body muscle tissues. However, in anaerobic conditions glucose is broken down and oxidized to pyruvate, which in turn is converted into lactate in the striated muscle and liver (VOET; VOET, 1995; HARRIS, 2004).

The creatine kinase (CK) enzyme is also an indicator of physical stress and muscle damage, but in the long-term as it takes up to 6 h to achieve the peak and returns to basal levels 48 h later (ANDERSON, 2010). The tissue of skeletal muscle depends on the availability of adenosine triphosphate (ATP) for a proper functioning (DAROIT; BRANDELLI, 2008). Generally, the muscle stores energy in a phosphocreatine (PC) form which can be rapidly converted into ATP through a reaction catalysed by CK (VOET; VOET, 1995). The ATP is consumed during high metabolic activity and is regenerated thanks to the CK catalysing the transphosphorylation reaction between PC and adenosine diphosphate (WALLIMANN et al., 1992; WYSS et al., 1992).

#### **2.4.1.3. Heart and respiration rate**

Heart rate variability has been largely used as a welfare indicator in animal production that increases in stressful situations (TERLOUW et al., 2004; BROOM; FRASER, 2007; MILLS et al., 2010) and during physical activity (PÖSÖ; PUOLANNE, 2005). Independently the stress type, either psychological (i.e. presence of a predator or human contact) or physical (i.e. during loading and

unloading), the heart rate measurement is a very rapid stress indicator (WARRISS, 2010) that provides the number of heart beats per unit time (MILLS et al., 2010). This measurement is defined as tachycardia to an acceleration of heart beats induced by the release of epinephrine from the sympathetic nervous system in response to a disturbing situation or physical activity; and bradycardia expressed by deceleration of heart beats in response to an emotional situation, such as fear or relaxing. During preslaughter handling, heart rate is increased during the use of electrical goads (KUCHENMEISTER et al., 2005; CORREA et al., 2010), at loading (van PUTTEN; ELSHOF, 1978), transport (GEVERINK et al., 1998; WARRISS, 1998) and according to social stress (de JONG et al., 2000).

The respiration rate is a stress indicator influenced by factor such as emotional disturbance without body activity (MELLOR; MURRAY, 1989; LORSCHY, 1997), physical exercise and fatigue (ZHANG et al., 1992; RITTER et al., 2007) and heat stress (FORREST et al., 1968). The heat stress increases breathing rate in order to control body temperature (MOUNT, 1972; ZHANG et al., 1992; BROWN-BRANDL et al., 2001) through the evaporative heat loss. Sweating is not an efficient tool to cool pigs; therefore, in situations they do not have an external source of water for evaporation from body surfaces, they use the panting for evaporative cooling (INGRAM, 1965; HALEY et al., 2008b). So, the breathing rate and panting behaviour can be used as a stress indicator.

#### **2.4.1.4. Body temperature**

Body temperature is an indirect indicator of the metabolic rate (WEBB, 1995) that provides information about welfare status due to its impacts on pigs' physiology. Thermoregulation is controlled by the hypothalamus that acts as a thermostat (CHARKOUDIAN, 2003; CAMPBELL, 2011) and is fundamental to maintenance of homeostasis (TERRIEN et al., 2011). The stress response increases body temperature due to the stimulation of the autonomic nervous system by the hypothalamus, that induces changes in the peripheral vascular tone and blood flow on the animals' body (BLESSING, 2003; MITCHELL, 2013). During a stressful situation, there is an activation of HPA axis that results in alterations in heat

production and dissipation by increasing the blood catecholamines and cortisol levels, and blood flow (SCHAEFER et al., 2002).

In heat stress, the blood flow to skin increases to 6 to 8 L/min due to a peripheral vasodilatation caused by the sympathetic vasodilator system (JOHNSON; PROPPE, 1996; as cited by CHARKOUDIAN, 2003). While in cold stress, the sympathetic vasoconstrictor system reduces blood flow to skin by shifting blood flow from the periphery to deep veins (ROWELL, 1983), which decreases the heat dissipation (CHARKOUDIAN, 2003).

The thermal equilibrium of pigs result comes from the balance between heat production and heat loss. Basically, in animals, this control is done by sweating (heat stress) and shivering (cold stress; CHARKOUDIAN, 2003). However, as pigs do not have functional sweat glands, the thermal regulation of pigs under heat stress mostly depends on wallowing, thermal panting (evaporative cooling) and heat dissipation (peripheral blood flow; INGRAM, 1965). Thermal stress is associated with other variables, such as: heart rate, rectal temperature, blood pressure and urinary excretion (PATIENCE et al., 2003).

## **2.5. Animal losses**

Losses of pigs during or after transport are not only indicative of a welfare problem, but also represent an economical loss for the pork production chain (SUTHERLAND et al., 2008; RITTER et al., 2009; DALLA COSTA; DALLA COSTA, 2014b). Transport losses are defined on arrival at the plant as dead-on-arrival (DOA), as non-ambulatory injured pigs (NAI), and non-ambulatory non-injured pigs (NANI; SUTHERLAND et al., 2008; RITTER et al., 2009). While NAI pigs are easily identified by a body injury (e.g. lameness due to broken leg or claw problem), NANI pigs present signs of acute stress, such as difficulty in moving, open-mouth breathing, skin discoloration, muscle tremors and abnormal vocalizations (ANDERSON et al., 2002; ELLIS et al., 2003).

The incidence of transport losses can be affected by several factors, such as poor preparation of pigs on farm (no or too short fasting interval prior to transport), handling at loading and unloading at the slaughterhouse (i.e. use of electric prods;

PEETERS et al., 2004; CORREA et al., 2010), environmental conditions (CLARK, 1979; TARRANT, 1989), health problems (TARRANT, 1989), and transport conditions (TARRANT, 1989; WERNER; REINERS; WICKE, 2005; RITTER et al., 2008). The proportion of fatigued pigs is greater during the winter, while dead pigs represent the greatest portion of total losses during summer (FITZERLAND et al. 2009). The author also found an increase of 7.55-fold in total losses when transport density was changed from the minimum (212.38 kg/m<sup>2</sup>) to the maximum (338.64 kg/m<sup>2</sup>) density. Similarly, Ritter et al. (2006) found that total of injured pigs, fatigued pigs, and total losses were reduced by 0.35, 0.37, and 0.52%, respectively, when transport density was reduced from 0.48 to 0.39 m<sup>2</sup>/pig.

Some potential causes of pigs dead-on-arrival are animal genetics, slaughter weight and increased production size (TARRANT, 1989; MURRAY; JOHNSON, 1998; WARRISS, 1998; ELLIS et al., 2003; WERNER; REINERS; WICKE, 2005; VECEREK et al., 2006). Fatigued pigs display signals such as open-mouth breathing, skin discoloration, muscle tremors, and abnormal vocalizations (ANDERSON et al., 2002), which are characteristic of acute-stress. A positive association ( $r = 0.81$ ) between dead and non-ambulatory pigs (HAMILTON et al., 2003) suggests a common cause for dead and non-ambulatory pigs (RITTER et al., 2009).

Fasting the pigs before loading reduce losses during transport (SMITH, 1937; GISPERT et al., 2000; RITTER, 2007; AVERÓS et al., 2008). Pigs transported with full stomach can suffer motion sickness and vomiting (BRADSHAW et al., 1996; GRANDIN, 2014). In this case, their cortisol levels are raised resulting from travel sickness (GISPERT et al., 2000). Averós et al. (2008) reported a double risk of mortality (0.3 vs. 0.54%) for not fasted pigs before loading (0 vs. <12; 12-18; >18 h). A feed withdrawal of 12-18 h before loading is considered suitable to prevent transport losses and meat quality defect (WARRISS, 1985, 1992; CHEVILLON, 1994; GUÀRDIA et al. 2004, 2005).

## **2.6. Carcass quality**

### **2.6.1. Carcass yield**

Some parameters such as carcass weight, fat depth, muscle depth, length,

shape, distribution of joints, sex, and presence of skin lesions have to be considered to estimate carcass quality. The carcass yield is calculated according to the proportion between the weight of the carcass live weight (WARRISS, 2010a), such as the equation:

$$\text{Carcass Yield} = (\text{Carcass weight/live weight}) \times 100.$$

Carcass yield is a useful measurement for pork carcass grade, which is estimated to be around 75% (FORTIN et al., 2003; POMAR et al., 2008; WARRISS, 2010a).

The carcass yield can be affected by fasting time (ELLIS, 1998; BEATTIE, et al., 2002; KEPHART; MILLS, 2005; FAUCITANO et al., 2006; Rodriguez et al. 2008; Faucitano et al. 2010a; PANELLA-RIERA et al., 2012), feed texture and meal frequency (FAUCITANO et al., 2006). The fasting time increase cold killing-out mainly due to a reduction in gut fill and offal weight (ELLIS, 1998; BEATTIE et al., 2002).

In the carcass, the proportion of tissues of interest is determined by the lean yield, which is calculated by dividing the weight of the tissues of interest by the overall or partial carcass weight (POMAR et al., 2008). In commercial conditions, the lean yield trait is estimated according to subjective (visual) or objective (caliper and optical probes) carcass measurements (KAUFFMAN; WARNER, 1993). In Canada, it is estimated by measuring fat thickness and muscle depth at the  $\frac{3}{4}$  last ribs using an optical probe (FORTIN et al., 2003). Lean yield is mostly affected by genetics (SELLIER, 1998; GISPERT et al., 2007), nutrition (ANDERSEN, 2000) and sex (LATORRE et al., 2003; 2004).

The live weight loss of pigs withdrawn of feed preslaughter may reach up to 5% in the first 24 h corresponding to around 5 kg body weight loss at an approximately rate of 0.2% per hour, or 0.25 kg/h (PELOSO, 2001). Carcass weight does not appear to be affected during the first 24 h as live weight losses are more associated to the excretion of urine and feces than to utilisation of body tissues (BEATTIE et al., 2002). Chevillon, Dubois and Vautier (2006) only found a significant effect of feed withdrawal on carcass weight loss from 24 h fasting on with losses

being estimated at approximately 100 g/h starting (CHEVILLON, 2001) corresponding to 1% decrease in carcass yield (FAUCITANO et al., 2006).

The small reduction in carcass weight (360 g/pig) and yield (0.33%) observed after 24 h fasting (CHEVILLON et al., 2006) corresponds to the removal of one meal (1.3-2 kg of feed) per day (KEPHART; MILLS, 2005; FAUCITANO et al., 2010). This feed cost can be thus saved by the producer as feed ingested by pigs in the last 10 h won't be converted into muscle tissue and carcass weight. Indeed, after its intake feed takes 4 h to 8 h to be absorbed in the small intestine and most of its nutrients take 9 h before being released into the blood circulation (WARRISS, 1985).

However, other studies failed to find effects on carcass weight losses even after 60 h feed withdrawal (ELLIS, 1998; TURGEON, 2003). The reason for this discrepancy in the results may be the differences in preslaughter stress levels, genotype or slaughter weight between studies (FAUCITANO et al., 2010a). According to Warriss (1985), transport to slaughter may contribute more to the live weight weight loss rate than fasting alone. Furthermore, researches showed that the recommended 24 h maximum fasting interval does not truly apply to pigs with high body energy stores (muscle glycolytic potential), such as pigs carrying the Hal<sup>n</sup> or RN<sup>-</sup> gene (MURRAY et al., 1989; BIDNER et al., 1999, 2004), and heavy pigs (> 170 kg; NANNI COSTA et al., 2002), for which a feed withdrawal time between 30 and 48 h appears more appropriate to reduce the risk of PSE pork production at no detriment for carcass weight or yield (MURRAY et al., 1989; LO FIEGO et al., 1997; BIDNER et al., 1999, 2004).

The nutritional condition of pigs can also influence weight losses during fasting time. Studying the effects of feed withdrawal time (4, 14 or 24 h), meal frequency (2 vs. 5 meals per day) and feed type (mash vs. pellet), Faucitano et al. (2006) reported a higher carcass yield in pigs fed a pelleted diet compared with a mash one (81.2 vs 79.3%), and in pigs fed 5 meals/day (80.7 vs 79.8%), regardless of the pre-slaughter feed withdrawal time applied. These findings were attributed to the differences in stomachs weight removed on the dressing line.

### **2.6.2. Skin lesions**

Skin lesions can be classified as superficial (scratches) or deep (bruises). Bruises are formed by the blood leaked from capillaries and trapped under the skin (ROBIN et al., 2015).

The presence of skin lesions is an important carcass quality trait for the market resulting in up to 6% carcass downgrading (MLC, 1985). According to North American pork industry estimates (VANSICKLE, 2002; SCHULTZ-KASTER; HILL, 2006; RIENDEAU, 2011), lesions alone contribute from \$0.08 to \$0.44 value loss per carcass. Their assessment also provides valuable information about animal welfare during preslaughter period (DALMAU et al., 2009; EUROPEAN FOOD SAFETY AUTHORITY, 2012) in which the most common sources of skin lesions are fighting (MOSS, 1978; GUISE; PENNY, 1989), rough handling (GEVERINK et al., 1996; RABASTE et al., 2007; FAUCITANO et al., 2008) and high density (GUISE; PENNY, 1989; BARTON-GADE; CHRISTENSEN, 1998).

The degree of carcass skin lesions is measured subjectively using a pictorial chart, either providing a general score based on the carcass appearance (1 = none to 5 = severe; MLC, 1985; Figure 4) or counting the bruises by anatomical location (ITP, 1996; BARTON-GADE et al., 1996a; Welfare Quality, 2009), and defining their shape to identify the infliction source (e.g., fighting, handling and overcrowding; ITP, 1996). Information about the bruise infliction can be taken by bruises' colour, with red bruises being the most recent ones (FAUCITANO, 2001).

### **2.7. Conversion of muscle into meat**

The muscles still metabolise energy, contract and produce heat after death. However, oxygen is not delivered to the cells and the metabolic end products are not removed, because there is no blood circulation (GREGORY, 1998). Under anaerobic conditions, lactic acid is produced from glucose either naturally present or derived from muscle glycogen to produce adenosine triphosphate (ATP; GREGORY, 1998; HUFF-LONERGAN; LONERGAN, 2005). The glycolysis process stops with the complete depletion of glycogen or, the accumulation of lactic acid when a very low

pH is reached (BOWKER et al., 2000; POSO; PUOLANNE, 2005). During this stage, the lactic acid accumulated in muscle decreases pH from a near neutral pH of 6.8-7.2 to about 5.6-5.8 (Figure 4).




Figure 3. Changes in glycogen, phosphocreatine (PC), ATP, and lactic acid concentrations during *post-mortem* metabolism (TARRANT; McLOUGHLIN; HARRINGTON, 1972).

The increased acidity causes a reduction in water binding ability and calcium release triggering muscle contraction. The cells do not have energy left and the metabolism stops at this stage. The intracellular pH arrives at its final level (GREGORY, 1998; BOWKER et. al., 2000; RYU; CHOI; KIM, 2005). Rigor mortis is a muscle state achieved when ATP is depleted being impossible to expulse  $\text{Ca}^{2+}$  from the sarcoplasm and depicts the beginning of permanent acto-myosin crossbridges (GREGORY, 1998; BOWKER et al., 2000).

## 2.8. Meat quality

Meat is defined as the final result of a sequence of biochemical and physical events during the conversion process from muscle into meat in the *post-mortem* period, which takes from 1 day to 2 weeks (HONIKEL, 2004). Furthermore, meat

quality is a generic term used to describe properties and perceptions of meat, such as combination of subjective and objective measurements of meat characteristics.

The following four factors can be associated with the abnormal *post-mortem* metabolism and meat quality variation: 1) the  $\text{Ca}^{2+}$  regulation; 2) the muscle ATPase activity; 3) glycogenolytic enzymes activity; and 4) muscle glycogen levels regulation. These factors in combination with muscle temperature changes influence the *post-mortem* glycolysis affecting pH fall rate and other meat quality traits, such as colour and water-holding capacity.

### 2.8.1. Meat pH

In normal conditions, during the conversion of muscle into meat muscle pH declines gradually from approximately a value of 7.0 to around 6.1 within 1 h after slaughter due to lactic acid accumulation (LAWRIE, 1966; BENDALL, 1973). It finally achieves the ultimate value ( $\text{pH}_u$ ) of 5.6-5.8 at 24 h *post-mortem* (SELLIER; MONIN, 1994).

However, *post-mortem* pH can vary depending on the glycogen content in the muscle at the time of slaughter. High levels of glycogen just prior to slaughter lead to higher lactic acid concentration in the muscle producing an approximately three-fold more rapid than normal pH decline (lower than 6 at 45 min *post-mortem*; WARRISS; BROWN, 1987; KÖHLER, 2001). This rapid pH fall combined with high muscle temperature (35-38°C), denatures myofibrillar proteins leading to a loss of their water binding capacity and PSE pork (GREGORY, 1998; NICOLAIEWSKY, 1998, TERRA, 1998).

When pigs are submitted to prolonged stress just before slaughter, because their glycogen reserves will be low at the time of slaughter, the muscle will not have enough substrate to produce lactic acid during the conversion of muscle into meat, resulting in higher ultimate pH ( $\text{pH}_u$ ) than 6, which is indicative of the DFD meat quality defect (TARRANT, 1989; CARRAGHER; MATTHEWS, 1996; GREGORY, 1998).

Feed withdrawal before slaughter is instrumental for raising muscle ultimate pH through the consumption of glycogen stores in the muscle (WARRISS, 1982;

JONES; ROMPALA; HAWORTH, 1985; EIKELENBOOM et al., 1991; MURRAY; JONES, 1994; TURGEON, 2003; GUÀRDIA et al., 2004, 2005; FAUCITANO et al., 2006; DALLA COSTA et al., 2010) resulting in improved water-holding capacity and colour (EIKELENBOOM et al., 1991; MURRAY; JONES, 1994; GISPERT et al., 2000; GUÀRDIA et al., 2005).

Faucitano et al. (2006) showed that 24 h feed withdrawal may increase the ultimate pH of 0.07 units in the *longissimus* muscle compared with 14 h. However, long fasting periods (> 22 h) may also increase the prevalence of DFD pork due to muscle glycogen exhaustion (EIKELENBOOM et al., 1991; GISPERT et al., 2000; GUÀRDIA et al., 2005). Whereas, short fasting periods may increase the risk of PSE pork production at a fasting time below 18 h (GUÀRDIA et al., 2004). However, other studies report little or no effects of preslaughter fasting on the ultimate pH value (DE SMET et al., 1996; MURRAY et al., 2001; BEATTIE et al., 2002; MORROW et al., 2002). The discrepancy in these findings indicates that the effects of feed withdrawal on pork ultimate pH are dependent on the pre-slaughter handling practices (i.e. mixing, long transport or lairage time) which are applied differently between studies (FERNANDEZ; TORNBERG, 1991; LEHESKA et al., 2003; BERTOL et al., 2005; FAUCITANO et al., 2006, 2010a) and the genotype (DE SMET et al., 1996). These effects are also muscle type-dependent, with higher ultimate pH value being more easily observable in locomotor muscles (i.e. shoulder and ham muscles) having a lower glycolytic potential (lower glycogen stores) and getting fatigued more rapidly (HAMBRECHT et al., 2005a; ROCHA et al., 2015).

### **2.8.2. Meat colour**

Meat colour is one of the most valuable meat quality properties for the pork industry (WARNER; KAUFFMAN; RUSSEL, 1993) as it may influence decision making of consumers at the retail level (BREWER; McKEITH, 1999; ANDERSEN; OKSBJERG; THERKILDSEN, 2005; GRUNERT, 2005).

Colour in meat is a perception of its optical properties. Meat is a translucent material (allowing light to pass through it). Therefore, light cast interact with the meat and part of it is absorbed and part scattered (WARRISS, 2010a). Absorption is

mainly influenced by the concentration and state of pigments, mainly myoglobin (WARNER, 1994; MANCINI; HUNT, 2005). A higher concentration of myoglobin produces a brighter red pork colour (YOUNG; WEST, 2001). The myoglobin forms are dependent from the availability of oxygen and the oxidation process. Myoglobin exists in three-forms: purple-red deoxymyoglobin in the absence of oxygen, bright red oxymyoglobin formed in the presence of air and brown metmyoglobin, resulting from myoglobin oxidation (TERRA, 1998). When freshly cut meat is exposed to air, the oxygenation or “blooming” takes 10-30 min resulting from the oxygen binding to myoglobin (CORNFORTH; JAYASINGH, 2004).

The structural properties in the superficial layers of the muscle influence the light scattering characteristics of meat. In meat, light scattering originates at the boundary between myofibrils and the sarcoplasm, and within the myofilament lattice of the myofibril, which happens due to instability of the sarcoplasmic and myofibrillar proteins (myosin). The aforementioned *post-mortem* muscle protein denaturation, in fact, leads to a “white precipitate” of denatured sarcoplasmic proteins accumulated on the myofibrils which disguise the red colour and make a semi-opaque background (JOO et al., 1999) reducing light diffusion into the material (MANCINI; HUNT, 2005). Due to that, there is a lower light absorption and a higher light reflection resulting in a lighter appearance producing PSE meat. Consequently, the light penetrates to a shallower depth and is adsorbed to a smaller extent by myoglobin. In PSE muscles, the increased scattered light occurs because of the reduced gaps between myofibrils. On the other hand, the dark red-pink colour of DFD pork is because the tissues are more translucent and scatter or reflect less light (BARTON-GADE; OLSEN, 1987; BREWER, McKEITH, 1999). The higher pH of DFD pork enhances the darkness because of less free water available in the muscle due to more water still bound to proteins and, as a consequence, light scattering is reduced (BREWER et al. 2001).

Sources of variation of pork colour include genetic selection, nutrition, pre-slaughter handling, *post-mortem* carcass handling (i.e. cooling rate and technique), processing and holding times and temperature before packaging or further use, packaging, distribution and marketing, including lighting and other display conditions (MANCINI; HUNT, 2005).

The  $L^*$ ,  $a^*$ ,  $b^*$  are coordinates of the tristimulus on the CIELAB colour space and can specify various colour space by determining the components of lightness and chromaticity. When colour is expressed in CIELAB,  $L^*$  defines lightness,  $a^*$  denotes red-greenness and  $b^*$  yellow-blueness; (CIELAB, 1976). Generally, these colour coordinates are measured by the Minolta Chromameter which evaluates the reflectance of the myoglobin and haemoglobin after exposure to oxygenation (BREWER et al., 2001). For instance, the greater the myoglobin concentration, the lower the lightness or  $L^*$  value. Nevertheless, the myoglobin state affects the lightness ratios being higher for oxidised and lower for oxygenated state (FERNANDEZ-LOPEZ; PEREZ-ALVAREZ; SAYAS-BARBERA, 2000).

The consumer's colour perception at the retail level can be simulated by the subjective colour assessment using the Japanese Colour Scales (JCS; NAKAI et al., 1975). The JCS are six plastic disks (from 1 = very pale to 6 = very dark) with a meat-like appearance developed by using objective colorimetry.

### **2.8.3. Drip loss**

Of all main muscle characteristics that affect appearance, shrinkage, processing properties, and palatability, drip loss is perhaps the most important for the meat industry economics and consumer perception (KAUFFMAN et al., 1986) due to loss of saleable weight and of a significant amount of proteins, water soluble vitamins and amino acids (112 mg of protein/mL of fluid; SAVAGE; WARRISS; JOLLEY, 1990).

Drip is defined as an aqueous solution purged from meat containing sarcoplasmic proteins, such as myoglobin which gives the characteristic red colour to it (SAVAGE et al., 1990).

Most animal muscles contain approximately 75% water held by 20% proteins. Most water (85%) is inside muscle cells, while the remaining 15% is located around them (HUFF-LONERGAN; LONERGAN, 2005). While half of the water contained in muscle cells is linked to muscle fiber proteins, the other half is linked to sarcoplasmic proteins. Water in the muscle presents a variable level of binding to the protein, with only 5% fully bound (HUFF-LONERGAN; LONERGAN, 2005). The mobility of water

is restricted by capillary forces and osmotic pressure in the live animal at normal pH. However, as mentioned before, when the pH falls close to the isoelectric point (pH 5.0 - 5.1) of meat proteins there is a considerable reduction in their hydration and in the ability to hold water tightly (TOLDRA, 2003).

The lateral shrinkage of muscle fibres is the single most important factor for drip loss formation (OFFER; COUSINS, 1992). The shrinkage of myofibrils and denaturation of proteins causes a dislocation of water within or out of the fiber. During rigor development there will be less space available among the muscle fibers for the retention of immobilised water which becomes free and flows out the fibrils space around fibres and fibres bundles (OFFER et al., 1992).

As it depends on the rate of pH drop in early *post-mortem* muscle, drip loss rate is influenced by several factors such as genetics (FUJII et al., 1991; HUFF-LONERGAN; LONERGAN, 2005), animal and carcass handling (JENNEN et al., 2007), temperature management *post-mortem* (JENNEN et al., 2007), nutrition (LYNCH et al., 1998; D'SOUZA et al., 2005; JENNEN et al., 2007) and processing (JENNEN et al., 2007).

Drip loss can be assessed by methodologies based on fluid absorption, such as the filter paper (KAUFFMAN et al., 1986), gravity, such as the bag and EZ-drip loss methods (HONIKEL, 1998; CHRISTENSEN, 2003; CORREA et al., 2007), and pressure, such as the filter paper press (GRAU; HAMM, 1953).

## **2.9. Meat quality categories**

Pork meat was initially divided into three quality categories based on pH, colour and drip loss variation, such as: normal (or RFN: red, firm, non-exudative), PSE and DFD pork. Pork classified as PSE has distinctly reduced processing and cooking yields (BOLES et al., 1991; PERSON et al., 2005), as well as decreased tenderness and juiciness (BENNETT et al., 1973). Pork classified as DFD appears very dark and sticky due to the self-cohesiveness of the myofibrils and their high water-binding capacity (FISHER, 2005). Furthermore, the high pH value, which is conducive to bacterial growth, results in shorter shelf-life for this meat type (FISCHER, 2005; HOLMER et al., 2009; FAUCITANO et al., 2010b).

PSE meat has been described as having an initial pH lower than 5.8 (SELLIER et al., 1994), pHu value between 5.5 and 5.7 (SELLIER et al., 1994), drip loss higher than 5% (WARNER; KAUFFMAN; GREASER, 1997; JOO et al., 1999) and Minolta colour L\* value higher than 50. Whereas, DFD pork is described as having a high pHu (> 6.0), very low drip loss (< 2%) and colour L\* values lower than 42 (WARNER, 1994; CORREA et al., 2007).

PSE and DFD conditions are commonly considered as a stress-related muscle disorders. The occurrence of PSE pork has been associated to animal exposure to acute stress just before slaughter, such as the use of electric prods, no or too short fasting duration or animal genetic background (Hal gene). Whereas, exposing animals to a long period of stress, such as long fasting periods combined with long transportation and lairage time and mixing cause the depletion of muscle glycogen reserves and may increase the incidence of DFD pork (VAN DE PERRE et al., 2010; WARRISS, 2010b).

However, two new quality categories have been added for a more reliable quality assessment considering the variation in either colour or drip loss, namely RSE (reddish-pink, firm, exudative) and PFN (pale, firm, non-exudative) pork (WARNER et al., 1993, 1997; VAN LAACK et al., 1994; Table 1).

Table 1. Pork quality based on parameters of ultimate pH (pHu), drip loss (DL), subjective colour (JCS), and objective colour (L\*) (CORREA et al., 2007)

<b>Quality classes<sup>1</sup></b>	<b>pHu</b>	<b>DL</b>	<b>JCS<sup>2</sup></b>	<b>L*</b>
PSE	<5,5	> 5%	1 – 1,5	>50
Moderate PSE	5,5 – 5,6	> 5%	2 – 3	≥ 50
PFN	5,5 - 5,8	< 5%	<3	> 50
RSE	5,6 - 5,8	>5%	3	42 – 50
RFN	5,6 - 5,8	2 – 5%	3	42 – 50
Moderate DFD	5,8 – 6,1	< 5%	3 – 4	42 – 45
DFD	> 6,1	<2%	≥ 4	≤ 42

<sup>1</sup> PSE (pale, soft, exudative), PFN (pale, firm, non-exudative), RSE (red, soft, exudative), RFN (red, firm, non-exudative), DFD (dark, firm, dry)

<sup>2</sup> Japanese Colour Standards from 1= pale to 6 = dark (NAKAI et al., 1975)

PFN pork has normal structure (firmness) and  $pH_u$  (5.5 – 5.8), but shows a paler colour, with an  $L^*$  value as low as the PSE meat ( $> 50$ ). The RSE meat is characterised by a normal  $pH_u$  value (between 5.6 and 5.8) and a reddish-pink colour ( $L^*$  between 42 and 50), but a drip loss as high as in the PSE meat class.

The PFN and RSE meat quality classes have been recently reported as major quality defects in Canada, representing more than 13% of all defects comparing to PSE (13%) and DFD pork (10%; MURRAY et al., 2001). Additionally, Faucitano et al. (2010b) reported that 47% of meat quality defects were classified as RSE meat, whereas PSE and DFD represented only 21 and 2%, respectively. However, despite the economic importance, the causes of RSE meat occurrence remain unknown (VAN LAACK et al., 1994).

## **2.10. The effects of preslaughter handling practices on animal welfare and meat quality**

### **2.10.1. Preslaughter feed withdrawal**

Feed withdrawal is a common practice for on-farm preparation before transport that is regulated by codes of practice (AAFC, 1993). It consists in removing the access to feed, while keeping free access to water, during a defined interval prior to slaughter. Communication between producer and slaughter plant is necessary to inform the loading time one day before loading and to prepare the animals to loading. Total feed withdrawal interval before slaughter has to be planned considering the on-farm feed withdrawal interval, time spent during loading, transport and unloading, and lairage period (DALLA COSTA; DALLA COSTA, 2014a).

Starting feed withdrawal time at the farm may result in economic benefits for the pork chain, such as feed saving (KEPHART; MILLS, 2005), better pigs' welfare during transport (BRADSHAW et al., 1996; GUÀRDIA et al., 2005; CORREA, 2011) resulting in decreased animal losses (GISPERT et al., 2000; AVERÓS et al., 2008), reduced carcass contamination due to a lower risk of gut contents spillage during carcass evisceration (FAUCITANO et al., 2010a) and improved pork quality due to greater easiness of handling (EIKELENBOOM; HOVING-BOLINK; SYBESMA, 1991; GUÀRDIA et al., 2004, 2005). It also helps reduce wastes to be disposed of at the

slaughter plant (MURRAY et al., 2001).

Overall, recommendations about the ideal duration of fasting preslaughter from 5 h in Canada (AAFC, 1993) to 48 h in USA (MILLER et al., 1997). Table 2 shows some feed withdrawal intervals recommended in different countries. Generally, a feed withdrawal of 12 hours is applied on-farm in Brazil. The Brazilian legislation recommends a feed withdrawal between 8-24 hours at the slaughter plant (INSTRUÇÃO NORMATIVA nº 3/2000; PORTARIA Nº 711). However, based on a review of literature reports, Faucitano et al. (2010a) suggest a period of 24 h between the last meal and slaughter as an acceptable compromise to obtain optimal carcass yield and pork quality.

Table 2. Feed withdrawal interval applied in different countries (FAUCITANO, 2007; DALLA COSTA; DALLA COSTA, 2014a)

Country	On-farm (hours)	Total (hours)
United Kingdom	4-8	18
Brazil	-	24
European Community	10	-
Canada	5	24
France	12-18	24-28

### 2.10.2. Physiological parameters

During preslaughter handling, Dalla Costa et al. (2008) reported an increase of 2.67 times in salivary cortisol level, rising from 0.267 on-farm to 0.714 after transport and unloading (Figure 3).




Figure 4. Variation in salivary cortisol levels (µg/dL) of pigs during preslaughter handling (adapted from DALLA COSTA et al., 2008).

Feed withdrawal period may influence cortisol levels in blood (PARROTT; MISSON, 1989; GISPERT et al., 2000; FAUCITANO et al., 2006). However, the literature results are contradictory. Gispert et al. (2000) reported lower level of cortisol in blood of pigs fasted on-farm from 12 to 18 h than those fasted for <12 h or >18 h. Whereas, other studies found increased blood cortisol levels starting from 15 h of fasting (HOUPTE et al., 1983; PARROT; MISSON, 1989). Faucitano et al. (2006) also found higher cortisol level in the urine of pigs fasted for 14 h compared with pigs fasted for 4 and 24 h. It is hard to explain how feed withdrawal affects the elevation of cortisol levels, as the effects are multifactorial (VON MICKWITZ, 1982), with cortisol levels being also affected by travel sickness during transport (BRADSHAW et al., 1996) and fighting during lairage (GISPERT et al., 2000), both responses due to increased demand for energy supply (Von MICKWITZ, 1982).

The concentrations of blood lactate and CK do not seem to be affected by fasting time (WARRISS et al., 1992; GISPERT et al., 2000). Leheska, Wulf and Maddock (2003) found no effect of fasting (48-h prior to harvest) or no fasting of pigs on lactate levels. Gispert et al. (2000) and Dalla Costa et al. (2008) reported no effects of fasting during on-farm feed withdrawal (< 12, 12-18 and > 18 h; 9, 12, 15 and 18 h, respectively) on lactate and CK levels. Edwards et al. (2010) reported an increase in lactate level prior to loading and transport from  $4.3 \pm 0.3$  mM (basal level) to  $7.4 \pm 0.4$  mM after pre-slaughter handling, movement, and stunning.

## **2.11. Loading**

Loading pigs onto the truck at the farm represents the greatest challenge pigs have to cope with pre-slaughter and also plays an important role in the pork industry economics because of its impact on animal losses, carcass condemnation and meat quality variation (WARRISS, 1998; GARCIA; McGLONE, 2014).

The stressfulness of the loading procedure results from the combination of different factors, such as group splitting in the finishing pen, distance moved from the pen to the load point, group size, handling system and eventually the design of the loading device (either ramp or quay).

The effects of loading procedures are included in those of the farm of origin,

which has been showed to be a major contributor for animal losses during transportation, general stress response and carcass and meat quality variation (BROWN et al., 1999; DALLA COSTA et al., 2007; DEWEY et al., 2009; FITZGERALD et al., 2009).

### **2.11.1. Exiting the farm gate**

Based on heart rate and blood cortisol variation, sorting pigs in the fattening pen and moving them to the loading area are the first intense stress and physical efforts which animals are exposed to during the loading process (BRADSHAW et al., 1996; CORREA et al., 2010; 2013). The major factors triggering this physiological response are group splitting in the finishing pen, walking a distance through the farm alley and interaction with the handler through narrow and enclosed spaces (handling).

Sorting pigs out of a pen can be very stressful to the group of pigs in the pen as it involves a close human-animal interaction and animal separation from the group (WAIBLINGER et al., 2006; JOHNSON et al., 2013). Group splitting at the time of loading is necessary to handle small groups of pigs through the alleys or to ship groups of uniform weight (split-marketing). Split marketing is very common at finishing farms in North America and in some European countries, such as France and Spain. It implies the removal of the heaviest 25 to 50% of pigs from a pen to market them 1-2 weeks earlier than the other pen-mates. The advantage of this practice is the reduction of production costs and the opportunity to ship batches with a more uniform market weight to the abattoir (SCROGGS et al., 2002).

In large swine units, animals are often imposed to walk long distances to get to the loading area. According to Ritter et al. (2008), the average distance over which pigs are moved to reach the loading area may be greater than 100 m. This situation may result in muscle fatigue during loading and transportation (RITTER et al., 2008). An increased frequency of open-mouth breathing and skin discoloration (signs of acute stress) at loading at the farm and at unloading at the slaughter plant have been reported in pigs moved over a long distance (up to 91 m) compared with those moved over a short one (< 24 m) to reach the loading area (RITTER et al., 2007,

2008). Therefore, the distance to the loading area should be minimized as much as possible (FAUCITANO, 1998) by using, for example, shipping rooms. Besides being a recommended practice for biosecurity and feed withdrawal control (CHEVILLON, 2005; CARR, 2006; FAUCITANO et al., 2010a), moving pigs to shipping pens improves animal welfare pre- and during transport. A 2 to 18 h wait in the shipping pen prior to loading was reported to reduce loading time, and pigs' heart rate and fatigue at loading, all resulting in 25% lower mortality rate during transport (CHEVILLON, 2005; GEESING et al., 2010).

At loading, pigs are usually moved using plastic sorting paddles and board, electric prods or flags. However, these tools do not have the same efficiency and the same effects on the behaviour and physiology of pigs during handling. Some authors describe the use of electric prods as being as stressful as the loading procedure (BECKER et al., 1985) and more aversive than 90% CO<sub>2</sub> gas inhalation (JONGMAN et al., 2000). Regulations and codes of practice recommend to limit, if not to avoid, the use of electric prods for pig handling at any stage of the preslaughter period (EC, 1993, 2002; CARC, 2001) because they increase moving time and animal losses (BENJAMIN et al., 2001; CORREA et al., 2010) besides making handling more difficult (BRUNDIGE et al., 1998; MCGLONE et al., 2004). They also produce a negative physiological (higher heart rate and blood lactate level) and behavioural (falls, slips, vocalization, overlaps, etc.) response (BRUNDIGE et al., 1998; BENJAMIN et al., 2001; BERTOL et al., 2002, 2005; HAMILTON et al., 2004; CORREA et al., 2010, EDWARDS et al., 2010), eventually resulting in poor pork quality (JONGMAN et al., 2000; CORREA et al., 2010). As Grandin (2001) points out, electric prods should be used only as last choice, and only when a fit animal refuses to move forward. Its abusive use may reflect the handler's lack of experience or training (FAUCITANO, 2001; CORREA et al., 2010; ROCHA et al., 2015) or the poor facilities design (DRIESSEN; GEERS, 2000; TORREY et al., 2013b).

Flags, paddles and plastic boards can be considered good alternatives to the use of electric prods. McGlone et al. (2004) compared the efficiency and effects of these tools, and concluded that the sorting board and the flag were the most efficient devices for moving pigs because they appear as solid and blocking walls. Whereas, when comparing the use of plastic boards combined with paddle and compressed air

prod along the alley and the loading ramp, Correa et al. (2010) reported a similar efficiency between these two handling tools, although the use of compressed air prod increased loading time due to increased behaviour of 180° turns, likely caused by the noise of the air compressor.

The correct choice of group size of pigs to move forward guarantees the smooth flow of pigs through the alleys and ramps, without stressing the animal and reducing the workload for handlers. Lewis and McGlone (2007) and Berry et al. (2009) reported a shorter loading time, a lower increase in heart rate and reduced animal losses on arrival at the plant when pigs were handled in groups of 5 to 6 compared to smaller or larger groups (< 5 or > 6 pigs).

### **2.11.2. Loading onto the truck**

Although widely used, loading ramps may be considered detrimental to the welfare of pigs if they do not meet some specific criteria (BROWN et al., 2005; KEPHART et al., 2010). Pigs showed to have difficulties, as showed by increased heart rate and slower pace, when negotiating steep ramps > 20° (van PUTTEN; ELSHOF, 1978; WARRISS et al., 1991).

The use of a staircase, including 8 cm high and 30 cm long steps, may be an alternative to reduce the slope of the ramp as it reduces slipping when the ramp is wet (i.e. presence of feces and urine; Grandin, 2000, 2008). Little spaced (15 cm) cleats also reduce slipperiness, especially on steep ramps, allowing an easier and safer movement of animals (WARRISS et al., 1991).

Given the difficult handling through ramps, alternative systems, such as a truck tailgate lift, hydraulic deck, and gantry have been developed and tested for their efficiency and effects on pig welfare at loading (CHRISTENSEN et al., 1994; BROWN et al., 2005; BERRY et al., 2012; CORREA et al., 2013; WESCHENFELDER et al., 2012, 2013). When compared to the ramp, the tail gate lift system showed to reduce the need of electric prods during handling (GRANDIN, 1991). However, other studies failed to find differences in heart rate, body temperature, blood cortisol concentration and meat quality in pigs loaded with a tailgate lift compared to pigs loaded with ramps (NANNI COSTA et al., 1999, 2002;

BROWN et al., 2005).

The loading gantry has also been tested as a promising alternative. It consists of an aluminum covered chute (7° slope) with two dual pivoting extension systems that allow proper positioning to both the barn and trailer. Berry et al. (2012) reported that when compared to a regular (19° slope) loading ramp, the use of the loading gantry reduced the electric prod use and produced fewer overlaps, slips, falls and balking incidences, pig vocalizations and transport losses (0.6 vs. 1%;  $P = 0.03$ ).

## **2.12. Transporting pigs**

Transportation is considered one of the most stressful event for pigs prior to slaughter (GEVERINK et al., 1998b) which may compromise pigs' welfare and pork quality (WARRISS; 1998, VON BORELL; SCHAFFER 2005; SCHWARTZKOPF-GENSEWIN et al., 2012). Major sources of stress during transport are ambient conditions (season of the year), vehicle design and loading density.

Season can affect the welfare of pigs during transport. Transport mortality rate and the percentage of panting pigs on arrival at the plant are reported to increase at temperatures above 20 and 17°C, respectively (SUTHERLAND et al., 2009; KEPHART et al., 2010). However, seasonal effects on total transport losses do not always correspond with increases in environmental temperature. Clark (1979) reported greater mortality losses in winter and Sutherland et al. (2009) found increased percentage of downers on arrival at the plant at ambient temperatures of 5°C and below.

### **2.12.1. Vehicle design**

Vehicle design plays a major role in the incidence of animal losses during transport as reported in a recent Canadian survey where vehicle design was included in the 16% transporter contribution to the incidence of transport mortality rate (DEWEY et al., 2009).

The design of the truck used to transport pigs differs around the world, ranging from small and single-decked trucks to large and triple-decked punch-hole trailers,

which can be equipped with fan-assisted ventilation depending on the geographic location and climate conditions. The results from comparative studies including trucks featuring ramps or hydraulic decks showed longer loading and unloading time, increased use of electric prods and higher proportion of animal losses in trailers equipped with internal ramps, such as the Canadian pot-belly (PB) trailer than in other vehicle types, equipped with hydraulic decks, such as a double-decked truck or a flat-deck trailer (ELLIS; RITTER, 2006; BARTON-GADE et al., 2007; RITTER et al., 2008; SUTHERLAND et al., 2009; KEPHART et al., 2010; CORREA et al., 2013).

The effects of vehicle design on animal welfare is also related to the variation in internal climate (temperature and humidity) resulting from differences in radiant heat load, the amount of water and heat produced by pigs, and the air flow pattern (LAMBOOIJ; SYBESMA, 1988). Heat stress is more important than cold stress, in terms of transport losses (SCHRAMA et al., 1996). Some trials conducted in Canada showed that the PB trailer was warmer than the flat-deck trailer either while stationary or moving due to the different pattern of the side openings and internal gates design (RITTER et al., 2008). Within the PB trailer, higher temperatures have been recorded in the front compartments of the middle and bottom deck (BROWN et al., 2011). The higher temperatures have been explained by the reduced ventilation (BROWN et al., 2011). Animal losses particularly increase when the vehicle is stationary under warm ambient conditions (RITTER et al., 2006) as in the absence of ventilation internal temperatures can be up to 9°C warmer than the external temperature (WESCHENFELDER et al., 2012; FOX et al., 2014). To improve the climate conditions in a stationary vehicle pigs should be cooled off by water sprinkling and/or fan-assisted ventilation (BROWN et al., 2011). Active ventilation contributes to reduce deaths during transport (NIELSEN, 1982). Whereas, when ambient temperature is above 20°C water sprinkling pigs for 5 min in a stationary truck before leaving the farm and during wait time at the plant proved to reduce drinking behaviour in lairage, physical fatigue at slaughter (lower blood lactate levels) and drip loss in pork meat (FOX et al., 2014; NANNONI et al., 2014). Christensen and Barton-Gade (1999) reported that when active ventilation and water misting/sprinkling are combined transport mortality rate can be significantly decreased as ventilation, besides removing the excessive humidity from the interior of the sprinkled truck,

increases evaporative cooling in pigs.

The effects of vehicle design on pork quality are inconclusive. A higher incidence of PSE (pale, soft, exudative) pork was reported in pigs transported in European featuring fixed decks and ramps (GUÀRDIA et al., 2004; LAMMENS et al., 2007). Whereas, Correa et al. (2013) reported no effect on pork quality when comparing the PB trailer with a double-decked truck equipped with an upper moving deck during 2 h travel time. Weschenfelder et al. (2013) studied the pork quality variation in pigs either transported with the PB trailer or the flat-deck trailer featuring a middle and upper moving decks and found higher  $pH_u$  values in the hams of pigs being transported with the PB trailer but only after short distances (45 min).

### **2.12.2. Transport density**

Transport density is the amount of space that pigs are provided on the truck during transportation. Even when loading densities are regulated by legislation, like in the case of the EU directive 95/29/EC, they are hardly met in practice as the chosen densities are frequently adjusted to transport conditions (weather and distance) and slaughter weight. In most EU countries and North America loading densities vary from 0.35 to 0.50  $m^2/100$  kg (WARRISS, 1998). Based on the measurements of the space needed for sternal recumbency, the minimum space required for slaughter pigs of 90-100 kg live-weight is equivalent to about 250  $kg/m^2$  (or 0.40  $m^2$ ; WARRISS, 1998; SCHWARTZKOPF-GENSWEIN et al., 2012).

High densities ( $< 0.40$   $m^2/100Kg$ ) are described to increase temperature and decrease air circulation inside the truck, negatively affecting its microenvironment (WARRISS et al., 1998a). They also prevent pigs from lying down resulting in increased blood creatine kinase levels (GUISE; PENNY, 1989; LEE et al., 2000; RITTER et al., 2009; CHAI et al., 2010), rectal prolapses (GUISE; WARRISS, 1989), lameness (KEPHART et al., 2010), carcass bruises (NANNI COSTA et al., 1999) and meat quality defects (BARTON-GADE; CHRISTENSEN, 1998; CHAI et al., 2010). Whereas, lower densities ( $> 0.50$   $m^2/100$  kg) result in fatigued pigs due the attempts to keep balance.

Transport floor space also has a major impact on dead and non-ambulatory pigs at the packing plant. Ritter et al. (2006, 2007) found increased mortality rate (0.08 to 0.27 %) at densities of 0.5 m<sup>2</sup>/100 kg and 0.4 m<sup>2</sup>/100 kg, respectively, and concluded that providing pigs (120 kg average weight) with 0.48 m<sup>2</sup>/pig was the most effective practice to reduce transport losses.

However, some studies (KEPHART et al., 2010; PILCHER et al., 2011) reported that floor space (< 260 kg/m<sup>2</sup> and 0.415 m<sup>2</sup>/pig, respectively) effects on the incidence of downer pigs can be exacerbated by short transportation time (< 2.5 h and < 40 min, respectively). Similar effects have been also reported on meat quality variation, with higher risk of PSE (pale, soft, exudative) pork occurring in short journeys (1 h) carried out at lower stocking densities (0.5 vs 0.25 m<sup>2</sup>/100 kg; GUARDIA et al., 2004).

### **2.13. Unloading**

On arrival at the plant, the primary recommendation is to begin unloading within 30 min and unloading to be completed within one hour to avoid heat and humidity rise inside the stationary truck and its negative consequences on animal welfare and meat quality of pigs, as previously described (AMI, 2012; DALLA COSTA; DALLA COSTA, 2014a). However, in commercial conditions, the waiting time to unload a truck after its arrival at the abattoir is very variable ranging from 5 min to 4 h (AALHUS et al., 1992; JONES, 1999). Research showed that as the waiting time increased by 30 min, the risk of pigs dying can increase by 2.2 times (HALEY et al., 2008a). Driessen and Geers (2001) also reported the greatest incidence of PSE pork after 30 min wait at 23°C ambient temperature or after 53 min wait at 11°C (70 and 45%, respectively). The presence of heat-stressed (or panting) pigs at unloading is a major risk factor for PSE pork production (VAN DER PERRE et al., 2010).

To meet this recommendation, a strict coordination of truck arrivals with the predicted number of pigs in lairage, lairage capacity and operation speed or the availability of a number of unloading docks allowing more than one truck to unload at the same time may shorten waiting times (WEYMAN, 1987).

If delay to unload is unavoidable, when pigs are waiting the truck, they must keep on moving, or sit in the shade or be cooled off using active ventilation and/or water sprinkling/misting in the truck (BENCH; SCHAEFER; FAUCITANO, 2008).

Although unloading can be considered less stressful than loading (based on heart rate and blood lactate level; CORREA et al., 2010; EDWARDS et al., 2010), poor handling resulting in carcass bruising and poor meat quality is inevitable at this stage, unless well-designed unloading facilities and trained personnel are used.

In practice the most common unloading device at the slaughter plant dock is the ramp, either combined to the truck ramp or lift. Keeping in mind that pigs have difficulties in descending a slope (BROWN et al., 2005), ramps steeper than 20° are not recommended as they result in greater pigs' heart rate, vocalization and backing up behaviour, increased handler interventions (voice and electric prods) and longer unloading time (WARRISS et al., 1991; RITTER et al., 2008; GOUMON et al., 2013; TORREY et al., 2013a, b). Higher pitched vocalization of pigs at unloading has been associated with greater risk of PSE pork production (van DER PERRE et al., 2010). The greater need for electric prod use observed on ramps may result in reduced easiness of handling (RABASTE et al., 2007) resulting in an higher incidence of fatigued pigs at unloading and lower pork quality (CORREA et al., 2013). When compared to ramps, the use of hydraulic lifts or decks reduces handling stress (lower heart rate), increases the easiness of handling and shortens off-load time (BROWN et al., 2005).

To ease handling of pigs, there should not be closed corner to negotiate, so they can move either when exiting the truck or unloading the dock. A greater heart rate and balking behaviour during unloading time have been reported in pigs facing angles or bends of 45° to 90° compared with 0° and 30° (WARRISS et al., 1992; GOUMON et al., 2013).

Handling problems due to hesitation and refusals of pigs to go forward resulting in greatest heart rate in pigs and handlers and greater number of pigs falling and slipping can also be caused by poor lighting, lack of shelter at the dock, the presence of a step (> 15 cm) or a gap between the truck deck floor and the unloading dock (SCAHAW, 2002; AMI, 2012; GOUMON et al., 2013) and the slippery ramp floor (TORREY et al., 2013b). Reducing the frequency of slips on the unloading ramp

not only prevents injuries, but also improves pork quality (ROCHA et al., 2015).

## 2.14. Lairage

Other than feeding the slaughter line, lairage is important for welfare and meat quality reasons as it provides an opportunity for animals to recover from stress and fatigue with clear benefits for meat quality (WARRISS, 1987). Lairage time, handling, facility design (pens and alleys), environment, whether pigs are mixed or not are all related to the recovery rate of pigs and carcass and meat quality variation (WARRISS, 2003; FAUCITANO, 2010).

Under commercial conditions, the lairage time applied is very variable (from < 1 to 15 h), depending on the lairage area size and livestock delivery schedule (GEVERINK et al., 1996; GISPERT et al., 2000). In Brazil, the minimum recommended lairage time is 3 h (DALLA COSTA; DALLA COSTA, 2014b).

There is evidence that a lairage time of 2 h is a sufficient time for recovery as showed by the blood cortisol levels (WARRISS et al., 1992; PÉREZ et al., 2002) and results in pigs easier to move based on the lower frequency of electric prod use (MILLIGAN et al., 1998). No or short lairage (15 - 60 min) is not recommended as it results in higher muscle temperature immediately before slaughter and higher level of lactic acid in the muscle resulting in increased incidence of PSE pork (FRAQUEZA et al., 1998; OWEN et al. 2000; SHEN et al., 2006). Whereas, longer lairage time proved to increase blood lactate and creatine kinase CK levels, reduce the risk to produce PSE pork (2% at 10 h; GUÀRDIA et al., 2005; DALLA COSTA et al., 2009), but to decrease carcass weight and increase the percentage of DFD (dark, firm, dry) pork (WARRISS et al., 1998b; GISPERT et al., 2000). Guàrdia et al. (2005) reported a 12% risk to produce DFD pork after 3 h lairage reaching a 25% risk after overnight lairage. The increase in DFD pork proportion with lairage time is the result of reduced muscle glycogen content at slaughter caused by the combined effect of fasting and fighting (MOSS; ROBB, 1978; NIELSEN, 1982; DE SMET et al., 1996; WARRISS et al., 1998b; NANNI COSTA et al., 2002; GUÀRDIA et al., 2009).

Mixing unfamiliar pigs, which is a common practice at commercial slaughter plants, inevitably causes some fighting to establish a new hierarchy within the group

(BARTON-GADE, 2008; DEEN, 2010) which results in a significant rise in body temperature (lasting up to 8 h after mixing) and greater risk of DFD pork (WARRISS; BROWN, 1985). Several studies found a more extended fighting rate in pigs kept in lairage for longer periods (overnight to > 24 h vs. 3 h) as a result of mixing and feed withdrawal (MURRAY; JONES, 1994; WARRISS, et al., 1998a; DE JONG et al. 1999; GUÀRDIA et al., 2009). Brown et al. (1999) reported a greater activity (fighting and drinking) in lairage in pigs fasted for longer time (18 vs. < 1 h). Fernandez et al. (1995) also found a greater activity in fasted pigs (24 vs. 0 h) than fed ones. To limit fighting in lairage it is either recommended to keep pigs in smaller groups (15 - 40 pigs; GEVERINK et al., 1996; RABASTE et al., 2007) or to mix very large groups (up to 200 pigs; GRANDIN 1990; CHRISTENSEN; BARTON-GADE 1997) in the pen. However, several studies reported a greater activity and number of fights during lairage in larger groups (BRYANT; EWBANK, 1972; PETHERICK, 1983; GEVERINK et al., 1996). Basically, the behaviour of pigs in both small and large groups is influenced by the perception of pigs regarding group size and on the availability of resources (food, water, etc.) in the pen (TURNER et al., 1999). Moss (1978) reported a greater incidence of fights in groups of 10 pigs kept at a low density (0.85 m<sup>2</sup>/pig) than in groups of 20 pigs stocked at 0.26 m<sup>2</sup>/pig in the pen, which highlights the greater impact of space allowance on social behaviour of pigs in the lairage pen than group size (MOSS, 1978). Based on these results and on the evidence that most fighting occurs within the first 30-60 min of lairage (MOSS, 1978; GEVERINK et al., 1996), to restrict fighting between mixed pigs stocking densities of 0.42 m<sup>2</sup>/pig for short lairage (< 3 h) and 0.66 m<sup>2</sup>/pig for long lairage (> 3 h) are recommended (WEEKS, 2008). Pigs may fight more at a low density, because larger available space allows greater opportunities to interact and fight with more individuals, while at high density pigs mostly interact with the few pigs in proximity rather than moving around the pen preventing fighting from spreading all over the pen (WEEKS, 2008). However, the densities applied in commercial conditions vary from 0.3 to 2.7 m<sup>2</sup>/pig (GISPERT et al., 2000; WEEKS, 2008).

Air temperature in lairage has an impact on pork quality variation (LAMMENS et al., 2007). Santos et al. (1997) reported that pigs have great difficulty in losing heat

and showed signs of stress, such as increased respiration rate under extreme climatic conditions ( $> 30^{\circ}\text{C}$  and  $\text{RH} > 80\%$ ).

Showering pigs in lairage is used to cool pigs ( $3^{\circ}\text{C}$  decrease in body temperature) and improve pork quality by reduce PSE incidence (LONG; TARRANT, 1990). It can also reduce aggressive behaviour during lairage and increase the ease of handling upon entrance into the stunning chute (WEEDING; GUISE; PENNY, 1993). Other benefit would be the improved electrical stunning efficiency as wetting the skin lowers its impedance, resulting in an easy and rapid achievement of unconsciousness prior to slaughter (WOTTON, 1996a). However, showering is not recommended at temperatures below  $5^{\circ}\text{C}$ , because it can result in pigs shivering and lead to DFD pork due to muscle energy depletion to maintain a constant body temperature (KNOWLES et al., 1998).

## **2.15. Handling pigs to slaughter**

Handling and facilities quality in the pre-stun area is of primary importance, given the need to handle pigs faster, so as to follow the speed of the slaughter-line, while ensuring the welfare of pigs at this stage. The combination of higher slaughter speed, poorly designed handling systems and large groups during the short period between the exit of the lairage pen and stunning may, in fact, result in greater proportion of slips and high-pitched vocalisation, and increased use of electric prods, all associated with lower pHu and higher drip loss in pork meat (van DER WAL et al., 1999; CHEVILLON, 2001; HAMBRECHT et al., 2005; RABASTE et al., 2007; DOKMANOVIĆ et al., 2014; ROCHA et al., 2015). To ease handling and keep a smooth and consistent flow of pigs to the stunner it is recommended to move pigs in groups of 6-8 to 18-20 in order to maintain slaughter throughput rates from 150 to 900 pigs/h, respectively (CHEVILLON, 2001). Moving large groups of pigs (45 pigs) through the lairage alleys may result in a greater incidence of PSE pork and blood splashing (BARTON-GADE et al., 1992).

Critical factors at this point are the entrance into the race and the "stop-start" forward motion of pigs towards the stunner, which are both observed in races feeding electrical and  $\text{CO}_2$  stunners. The entrance of pigs into a  $\text{CO}_2$  stunner has been

significantly improved by the Danish group-wise stunning system (CHRISTENSEN; BARTON-GADE, 1997). Compared to the traditional double race system, the group-wise handling system resulted in lower proportion of PSE and blood splashed pork due to reduced prodding and muscle exercise (CHRISTENSEN; BARTON-GADE, 1997; FRANCK et al., 2003).

## **2.16. Stunning methods**

Stunning before slaughter is a legal requirement according to the European legislation (EU Council Directive 93/119/EC, 1993) and the Humane Slaughter Act (1978). In farm animals, stunning is applied in order to establish an unconscious state (GREGORY et al., 2009) so as to ensure that animals do not suffer unnecessarily and are insensible to the slaughter procedure (RAJ, 2008).

Nowadays, the two most methods used for stunning pigs at commercial slaughter plants are the electrical and carbon dioxide (CO<sub>2</sub>) stunning.

### **2.16.1. Electrical stunning**

The electrical system is the most used stunning method for pigs. In this method, electricity is passed through the brain to cause instantaneous insensibility and unconsciousness by inducing a tonic/clonic epileptic fit, before any pain stimulus associated with the application itself is detected and transmitted to the central nervous system (WOTTON, 1996a). Electrical stunning must be applied properly to deliver a minimum electric current of 1.3 A in less than one second and ensure that pigs are instantly rendered unconscious, (HOENDERKEN, 1978).

Electrical stun can be either performed in three different forms. It can be done by applying an electric current across the head only (head-only stunning) or across the head followed by a second current applied to the chest, behind the position of the heart (head-to-chest/brisket stunning) or on the back (head-to-back stunning) to induce cardiac ventricular fibrillation. When compared with carbon dioxide stunning, head-only electrical stunning increases the rate of the *post-mortem* muscle glycolysis (PSE pork) in pigs due to increased post-stun convulsions and produces a 20-fold

increase in the incidence of haemorrhages in the muscle due to increased blood pressure and muscle activity (LAMBOOIJ; SYBESMA, 1988; WOTTON et al., 1992).

Compared to the head-only stunning, the head-to-chest system reduces the negative effects of electrical stunning because the application of the chest electrode inhibits the spinal nerve function reducing the post-stun clonic convulsions (GILBERT et al., 1984). However, when compared with CO<sub>2</sub> stunning, head-to-chest stunning still increases the incidence of PSE and blood splashing on the loins and hams (VELARDE et al., 2000a, 2001).

### **2.16.2. Carbon-dioxide (CO<sub>2</sub>) stunning**

Inhalation of carbon dioxide reduces the pH of blood and cerebrospinal fluid, which causes respiratory and metabolic acidosis, and consequently inducing a state of analgesia and anaesthesia in the animal (RAJ, 2008). In this system, pigs are loaded into a cradle and lowered into a concrete well pre-filled with a high gas concentration (85-90%). To avoid the presence of still conscious animals at the exit of the stunner, the minimum CO<sub>2</sub> cycle time should be longer than 130 sec. (VELARDE et al., 2000b). Pork quality is affected by gas concentration. Nowak et al. (2007) reported improvements in meat quality (higher pH<sub>u</sub> values) after stunning with 90% compared to 80% CO<sub>2</sub>.

### **2.17. Slaughter methods**

In order to limit post-electrical stun convulsions and limit the effects on meat quality variation pigs should be bled promptly (< 15 sec in case of head-only stun; WOTTON, 1996b), whilst still in the tonic phase, and in the lying position. Woltersdorf and Troeger (1987) and Faucitano et al. (1998) observed a significant decrease in the incidence of PSE pork when animals were bled out rapidly in a lying position. As head-to-chest/back stunning induces cardiac arrest and thus to animal death, the promptness of bleeding is not an issue as it is only a procedure to drain blood out of the carcass. However, to prevent blood clots in large vessels and obtain a satisfactory bleeding, a stun-to-stick interval of 3 min is recommended (GREGORY,

1998).

Vertical bleeding is usually observed in CO<sub>2</sub> stunned pigs as the extremities of pigs are relaxed (no convulsions) at the exit of the system (FAUCITANO, 2010). The stun-to-stick time for pigs exiting from a CO<sub>2</sub> system depends on the time of exposure to the gas and the gas concentration ranging from 25-35 to 40-50 sec for exposure to 80% and 90% CO<sub>2</sub>, respectively (NOWAK et al., 2007).

### 3. REFERENCES

AAFC, 1993. **Recommended code of practice for the care and handling of farm animals - pigs**. Agriculture and Agri-Food Canada Publ. 1898/E.

AALHUS, J. L. CANADA. AGRICULTURE CANADA. RESEARCH BRANCH. **Environmental conditions for swine during marketing for slaughter: A national review**. Lacombe, Alta: Research Branch, Agriculture Canada, 1992.

AMI - AMERICAN MEAT INSTITUTE. Recommended animal handling guidelines & audit guide: A systematic approach to animal welfare. **American Meat Committee**. Washington: American Meat Institute Animal Welfare Institute, 2012.

ANDERSEN, H. J. What is pork quality? **Quality of meat and fat in pigs as affected by genetics and nutrition. Proceedings of the joint session of the EAAP commissions on pig production, animal genetics and animal nutrition, Zurich, Switzerland, 25 August 1999**, n. 100, p. 15-26, 2000.

ANDERSEN, H. J.; OKSBJERG, N.; THERKILDSEN, M. Potential quality control tools in the production of fresh pork, beef and lamb demanded by the European society. **Livestock Production Science**, v. 94, n. 1, p. 105-124, 2005.

ANDERSON, D. B. Relationship of blood lactate and meat quality in market hogs. In: 63<sup>rd</sup> RECIPROCAL MEAT CONFERENCE, 2010, Lubbock. **Anais...** Lubbock, 2010. Disponível em: <http://fass.acrobat.com/p86799506/>. Acesso em: 03 de outubro de 2012.

ANDERSON, D. B., D. J. IVERS, M. E. BENJAMIN, H. W. GONYOU, D. J. JONES, K. D. MILLER, R. K. MCGUFFEY, T. A. ARMSTRONG, D. H. MOWREY, L. F. RICHARDSON, R. SENERIZ, J. R. WAGNER, L. E. WATKINS, AND A. G. ZIMMERMANN. Physiological responses of market hogs to different handling practices. In: PROCEEDING OF AMERICAN ASSOCIATION OF SWINE VETERINARIANS, 2002, Kansas City. **Anais...** Kansas City: AASV, 2002. p. 399

AREY, D. S.; EDWARDS, S. A. Factors influencing aggression between sows after mixing and the consequences for welfare and production. **Livestock Production Science**, v. 56, n. 1, p. 61-70, 1998.

AVERÓS, X.; KNOWLES, T. G.; BROWN, S. N.; WARRISS, P. D.; GONSÁLVEZ, L. F. Factors affecting the mortality of pigs being transported to slaughter. **The Veterinary Record**, v. 163, n. 13, p. 386-390, 2008.

AYO, J. O.; OLADELE, S. B.; FAYOMI, A. Stress and its adverse effects on modern swine production. **Pig News and Information**, v. 19, n. 2, p. 51N, 1998.

BARTON-GADE, P. A.; OLSEN, E. V. Experience in measuring the meat quality of stress-susceptible pigs. In: TARRANT, P. V.; EIKELNOOM, G.; MONIN, G.. **Evaluation and Control of Meat Quality in Pigs**. Dublin: Springer Netherlands, 1987. cap. 2. p. 117-128.

BARTON-GADE, P.; BLAABJERG, L.; CHRISTENSEN, L. New lairage system for slaughter pigs. Effect on behaviour and quality characteristics. In: **Proceedings of the 38<sup>th</sup> International Congress of Meat Science and Technology**, 1992, Clermont-Ferrand. **Anais...** Clermont-Ferrand: ICoMST, 1992. p. 161-164.

BARTON-GADE, P.; WARRISS, P. D.; BROWN, S. N.; LAMBOOIJ, B. Methods of improving pig welfare and meat quality by reducing stress and discomfort before slaughter - Methods of assessing meat quality. In: Proceedings of the EU-Seminar - New information on welfare and meat quality of pigs as related to handling, transport and lairage conditions, n. 166, 1996, Mariensee. **Anais...** Mariensee:, 1996. p. 57-63. Edição Especial.

BARTON-GADE, P.B.; CHRISTENSEN, L. Effect of different loading densities during transport on welfare and meat quality in Danish slaughter pigs. **Meat Science**, v. 48, n. 3, p. 237-247, 1998.

BARTON-GADE, P.; CHRISTENSEN, L.; BALTZER, M.; PETERSEN, V. J. Causes of pre-slaughter mortality in Danish slaughter pigs with special emphasis on transport. **Animal welfare**, v. 16, n. 4, p. 459-470, 2007.

BEATTIE, V.E.; BURROWS, M.S.; MOSS, B. W.; WEATHERUP, R.N. The effect of food deprivation prior to slaughter on performance, behaviour and meat quality. **Meat Science**, v. 62, n. 4, p. 413-418, 2002.

BECKER, B. A.; NIENABER, J. A.; DeSHAZER, J. A.; HAHN, G. L. Effect of transportation on cortisol concentrations and on the circadian rhythm of cortisol in gilts. **American Journal of Veterinary Research**, v. 46, n. 7, p. 1457-1459, 1985.

BENCH, C.; SCHAEFER, A.; FAUCITANO, L. The welfare of pigs during transport. In: Schaefer, A.; Faucitano, L. **Welfare of pigs: from birth to slaughter**. 1. ed. New York: Wageningen Academic, 2008. cap. 6, p. 161-180.

BENDALL, J. R. *Post-mortem* changes in muscle. In: BOURNE F. H. **The structure and function of muscle**, 2. ed. New York: Academic Press, 1973. cap. 2, p. 244-309.

BENJAMIN, M. E.; GONYOU, H. W.; IVERS, D. J.; RICHARDSON, L. F.; JONES, D. J.; WAGNER, J. R.; SENERIZ, R.; ANDERSON, D. B. Effect of animal handling method on the incidence of stress response in market swine in a model system. **Journal of Animal Science**, v. 79, n. 1, p. 279, 2001. Edição Especial.

BENNETT, M.E., BRAMBLETT, V.D., ABERLE, E.D., HARRIGTON, R.B. Muscle quality, cooking method and aging vs. palatability of pork loin chops. **Journal of Food Science**, v. 38, n. 3 p. 536-538, 1973.

BERRY, N. L.; RITTER, M.; BRUNTON, E.; STREMFERFER, W.; HOAG, B.; WOLFE, J.; FITZGERALD, N.; PORTH, M.; DELANEY, D.; WELDON, T. Effects of moving market weights pigs in different group sizes during loading on stress responses and transport losses at the packing plant. In: Proceeding Midwest

American Society of Animal Science, 2009, Des moines. **Anais...** Des moines: ASAS, 2009. p. 5.

BERRY, N. L.; JOHNSON, A. K.; HILL, J.; LONERGAN, S.; KARRIKER, L. A.; STALDER, K. J. Loading gantry versus traditional chute for the finisher pig: Effect on welfare at the time of loading and performance measures and transport losses at the harvest facility. **Journal of Animal Science**, v. 90, n. 11, p. 4028-4036, 2012.

BERTOL, T. M.; ELLIS, M.; HAMILTON, D. N.; MCKEITH, F. Effect of handling intensity on blood acid-base balance in slaughter weight pigs. **Journal of Animal Science**, v. 80, n. 2, p. 86, 2002. Edição Especial.

BERTOL, T. M.; ELLIS, M.; RITTER, M. J.; MCKEITH, F. K. Effect of feed withdrawal and handling intensity on longissimus muscle glycolytic potential and blood measurements in slaughter weight pigs. **Journal of Animal Science**, v. 83, n. 7, p. 1536-1542, 2005.

BIDNER, B. S.; ELLIS, M.; MILLER, K. D.; HEMANN, M.; CAMPION, D.; MCKEITH, F. K. Effect of the RN gene and feed withdrawal prior to slaughter on fresh longissimus quality and sensory characteristics. **Journal of Animal Science**, v. 77, n. 1, p. 49, 1999. Edição Especial.

BIDNER, B. S.; ELLIS, M.; WITTE, D. P.; CARR, S. N.; MCKEITH, F. K. Influence of dietary lysine level, pre-slaughter fasting, and rendement napole genotype on fresh pork quality. **Meat Science**, v. 68, n. 1, p. 53-60, 2004.

BLESSING, W. W. Lower brainstem pathways regulating sympathetically mediated changes in cutaneous blood flow. **Cellular and Molecular Neurobiology**, v. 23, n. 4-5, p. 527-538, 2003.

BOLANDER, F. F. Nonclassical endocrinology. In: BOLANDER, F. F. **Molecular Endocrinology**. 3. ed. San Diego: Elsevier Academic Press, 2004. cap. 3, p. 61-100.

BOLES, J. A., PARRISH, F. C., SKAGGS, C. L.; CHRISTIAN, L. L. Effect of porcine somatotropin, stress susceptibility, and final end point of cooking on the sensory, physical, and chemical properties of pork loin chops. **Journal of animal science**, v. 69, n. 7, p. 2865-2870, 1991.

BOWKER, B. C.; GRANT, A. L.; FORREST, J. C.; GERRARD, D. E. Muscle metabolism and PSE pork. **Journal of Animal Science**, v. 79, n. 1, p. 1-8, 2000.

BRADSHAW, R. H., PARROTT, R. F., GOODE, J. A., LLOYD, D. M., RODWAY, R. G., & BROOM, D. M. Behavioural and hormonal responses of pigs during transport: effect of mixing and duration of journey. **Animal Science**, v. 62, n. 03, p. 547-554, 1996.

BRANDT, P.; ROUSING, T.; HERSKIN, M. S.; AASLYNG, M. D. Identification of *post-mortem* indicators of welfare of finishing pigs on the day of slaughter. **Livestock Science**, v. 157, n. 2, p. 535-544, 2013.

BRANDT, P; AASLYNG, M. D. Welfare measurements of finishing pigs on the day of slaughter: A review. **Meat Science**, v. 103, p. 13-23, 2015.

BRASIL. 29 de março de 1952. Decreto nº 30.691, de 29/03/52 do Regulamento de Inspeção Industrial e Sanitária de Produtos de Origem Animal. Brasília, DF: Ministério da Agricultura Pecuária e Abastecimento 1952. 154 p.

BRASIL Ministério da Agricultura, Pecuária e Abastecimento. Secretária de Defesa Agropecuária. **Portaria nº 711/1995**. Normas técnicas de instalações e equipamentos para abate e industrialização de suínos. **Diário oficial da União**, Brasília, 01 nov. 1995. Seção 1, p. 17625.

BRASIL Ministério da Agricultura, Pecuária e Abastecimento. Secretária de Defesa Agropecuária. **Instrução Normativa nº 3/2000**. Regulamento técnico de métodos de insensibilização para o abate humanitário de animais de açougue. **Diário oficial da União**, Brasília, 24 jan. 2000. Seção 1, p. 14.

BREWER, M. S.; MCKEITH, F. K. Relationship among consumer quality characteristics and purchase intent of pale, soft, exudative, normal and dark, firm, dry pork at retail. **Journal of Food Science**, v. 64, n. 1, p. 1-4, 1999.

BREWER, M. S.; BIDNER, B.; MEISINGER, D.J.; MCKEITH, F.K. Measuring pork color: effects of bloom time, muscle, pH and relationship to instrumental parameters. **Meat Science**, v. 57, n. 2, p. 169-176, 2001.

BROOM, D. M. Indicators of poor welfare. **The British Veterinary Journal**, v. 142, n. 6, p. 524-526, 1986.

BROOM, D. M. Animal welfare: concepts and measurement. **Journal of Animal Science**, v. 69, n. 10, p. 4167-4175, 1991.

BROOM, D. M. Les concepts de stress et de bien-être. **Recueil de Médecine Vétérinaire**, v. 164, p. 715-722, 1998.

BROOM, D. M.; JOHNSON, K. G. **Stress and animal welfare**. London: Kluwer Academic Publishers, 1993. p. 217.

BROWN, S. N.; KNOWLES, T. G.; EDWARDS, J. E; WARRISS, P.D. Relationship between food deprivation before transport and aggression in pigs held in lairage before slaughter. **Veterinary Record**, v. 145, n. 22, p. 630-634, 1999.

BROWN, S. N.; KNOWLES, T. G.; WILKINS, L. J.; CHADD, S. A.; WARRISS, P. D. The response of pigs to being loaded or unloaded onto commercial animal transporters using three systems. **Veterinary Journal**, v. 170, n.1, p. 91-100, 2005.

BROOM D. M.; FRASER, A. F. **Domestic Animal Behaviour and Welfare**. Wallingford: CAB International, 2007. p. 438.

BROWN, J. A.; SAMARAKONE, T. S.; CROWE, T.; BERGERON, R.; WIDOWSKI, T.; CORREA, J. A.; FAUCITANO, L.; TORREY, S.; GONYOU, H. W. Temperature

and humidity conditions in trucks transporting pigs in two seasons in Eastern and Western Canada. **Transaction of the ASABE**, v. 54, p. 2311-2318, 2011.

BROWN-BRANDL, T. M.; EIGENBERG, R. A.; NIENABER, J. A.; KACHMAN, S. D. Thermoregulatory profile of a newer genetic line of pigs. **Livestock Production Science**, v. 71, n. 2, p. 253-260, 2001.

BRUNDIGE, L., OLEAS, T., DOUMIT, M., ZANELLA, A.J. Loading techniques and their effect on behavioral and physiological responses of market weight pigs. **Journal of Animal Science**, v. 76, n. 1, p. 99, 1998. Edição Especial.

BRYANT, M. J.; EWBANK, R. Some effects of stocking rate and group size upon agonistic behaviour in groups of growing pigs. **British Veterinary Journal**, v. 128, n. 2, p. 64-70, 1972.

CAMPBELL, I. Body temperature and its regulation. **Anaesthesia & Intensive Care Medicine**, v. 12, n. 6, p. 240-244, 2011.

CANNON, W. B. **Bodily changes in pain, hunger, fear and rage: an account of recent researches into the function of emotional excitement**. Nova York: Appleton, 1929. p. 334.

CARR, S. N. **Effects of seasonal environment, on-farm handling, transport, stocking density, and time in lairage on body temperature, serum cortisol levels, and pork lean quality of market hogs**. 2006. 100 f. PhD Thesis, University of Missouri, Columbia, 2006.

CARRAGHER, J. F.; MATTHEWS, L. R. Animal behaviour and stress: impacts on meat quality. In: PROCEEDINGS-NEW ZEALAND SOCIETY OF ANIMAL PRODUCTION, 56., 1996, New Zealand. **Anais...** New Zealand: New Zealand Society of Animal Production, 1996. p. 162-166.

CARSTENS, E.; MOBERG, G. P. Recognizing pain and distress in laboratory animals. **ILAR Journal**, v. 41, n. 2, p. 62-71, 2000.

CHAI, J.; XIONG, Q.; ZHANG, C. X.; MIAO, W.; LI, F. E.; ZHENG, R.; PENG, J.; JIANG, S. W. Effect of pre-slaughter transport plant on blood constituents and meat quality in halothane genotype of NN Large White x Landrace pigs. **Livestock Science**, v. 127, n. 2, p. 211-217, 2010.

CHARKOUDIAN, N. Skin blood flow in adult human thermoregulation: how it works, when it does not, and why. **Mayo Clinic Proceedings**, v. 78, p. 603-612, 2003.

CHEVILLON, P. Le contrôle des estomacs de porcs à l'abattoir: miroir de la mise à jeun en élevage. **Techni-Porc**, v. 17, n. 6, p. 47-54, 1994.

CHEVILLON, P. Pig welfare during pre-slaughter and stunning. In: PROCEEDINGS OF 1<sup>st</sup> INTERNATIONAL VIRTUAL CONFERENCE ON PORK QUALITY, WELFARE, TRANSPORT, SLAUGHTER AND CONSUMER, 2000, Concórdia. **Anais...** Concórdia: EMBRAPA:CNPSA, 2001. p. 145-158.

CHEVILLON, P. Préparation et départ de la ferme: les 24 dernières heures à la ferme. In: 26 COLLOQUE SUR LA PRODUCTION PORCINE, 25., 2005, Saint-Hyacinthe. **Anais...** Saint-Hyacinthe, 2005. p. 75-98.

CHEVILLON, P.; DUBOIS, A.; VAUTIER, A. Impact de la durée de mise à jeun, du mode d'alimentation et d'élevage sur la qualité technologique des jambons cuits. **Techni porc**, v. 29, n. 2, p. 29-38, 2006.

CHRISTENSEN, L.B. Drip loss in porcine m. *longissimus dorsi*. **Meat Science**, v. 63, n. 4, p. 469-477, 2003.

CHRISTENSEN, L.; BARTON-GADE, P. New Danish developments in pig handling at abattoirs. **Fleischwirtschaft**, v. 77, n. 7, p. 604-607, 1997.

CHRISTENSEN, L.; BARTON-GADE, P. Temperature profile in double decker transporters and some consequences for pig welfare during transport. In: FARM ANIMAL WELFARE – WHO WRITES THE RULES? **Occasional Publication of the British Society of Animal Science**, v. 23, p. 125-128, 1999.

CHRISTENSEN, L.; BARTON GADE, P.; BLAABJERG, L. O. Investigation of transport conditions in participating countries in the EC Project: PL 920262. In: PROCEEDINGS OF 40<sup>th</sup> INTERNATIONAL CONGRESS OF MEAT SCIENCE AND TECHNOLOGY, 1994, The Hague. **Anais...** The Hague: ICoMST, 1994. p. W-2.01.

CLARK, E.G. A *post-mortem* survey of transport deaths in Saskatchewan market hogs. **Western Hog Journal**, v. 1, p. 34-36, 1979.

Commision Internacional de l'Eclairage, 1976. International Commission on Illumination, Colorimetry: Official Recommendations of the International Commission on Illumination. Publication CIE N° 15 (E-1.3.1). Bureau Central de la CIE, Paris, France.

COMMITTEE ON ALLEVIATION OF DISTRESS IN LABORATORY ANIMALS. Recognition and Alleviation of Distress in Laboratory Animals, 2008. <<http://www.nap.edu/catalog/11931.html>>. Acesso em: 14 de set. 2015.

CONTE, S.; FAUCITANO, L.; BERGEON, R.; TORREY, S.; GONYOU, H. W.; CROWE, T.; TOTH TAMMINGA, E.; WIDOWSKI, T. M. Effects of season, truck type, and location within truck on gastrointestinal tract temperature of market-weight pigs during transport. **Journal of animal science**, v. 93, n. 12, p. 5840-5848, 2015.

CORNFORTH, D. P.; JAYASINGH, P. Chemical and physical characteristics of meat / Color and Pigment. In: JENSEN, W. K.; DEVINE, C.; DIKEMAN, M. **Encyclopedia of Meat Science**. 1. ed. Oxford: Elsevier Science Ltd., 2004. p. 249-256.

CORREA, J. A.; MÉTHOT, S.; FAUCITANO, L. a modified meat juice container (EZ-Driploss) procedure for a more reliable assessment of drip loss and related quality changes in pork meat. **Journal of Muscle Foods**, v. 18, n. 1, p. 67-77, 2007.

CORREA, J. A.; TORREY, S.; DEVILLERS, N.; LAFOREST, J. P.; GONYOU, H. W.; FAUCITANO, L. Effects of different moving devices at loading on stress response and meat quality in pigs. **Journal of Animal Science**, v. 88, n. 12, p. 4086-4093, 2010.

CORREA, J. A. Effect of farm handling and transport on physiological response, losses and meat quality of commercial pigs. **Advances in Pork Production**, v. 22, p. 249-256, 2011.

CORREA, J. A.; GONYOU, H. W.; TORREY, S.; WIDOWSKI, T.; BERGERON, R.; CROWE, T. G.; LAFOREST, J. P.; FAUCITANO, L. Welfare and carcass and meat quality of pigs being transported for 2 hours using two vehicle types during two seasons of the year. **Canadian Journal of Animal Science**, v. 93, n. 1, p. 43-55, 2013.

DALLA COSTA, O. A., FAUCITANO, L., COLDEBELLA, A., LUDKE, J. V., PELOSO, J. V., DALLA ROZA, D., PARANHOS DA COSTA, M. J. R. Effects of the season of the year, truck type and location on truck on skin lesions and meat quality in pigs. **Livestock Production Science**, v. 107, n. 1, p. 29-36, 2007.

DALLA COSTA, O. A.; LUDKE, J. V.; PARANHOS DA COSTA, M. J. R.; FAUCITANO, L.; COLDEBELLA, A.; KICH, J. D.; PELOSO, J. V.; DALLA ROZA, D. Fasting time at farm on hormonal status and physiological parameters of heavy-weight slaughter pigs. **Ciência Rural**, v. 38, n. 8, p. 2300-2306, 2008.

DALLA COSTA, O. A.; LUDKE, J. V.; COLDEBELLA, A.; KICH, J. D.; PARANHOS DA COSTA, M. J. R.; FAUCITANO, L.; PELOSO, J. V.; DALLA ROZA, D. Efeito do manejo pré-abate sobre alguns parâmetros fisiológicos em fêmeas suínas pesadas. **Ciência Rural**, v. 39, n. 3, p. 852-858, 2009.

DALLA COSTA, O. A.; LUDKE, J. V., PARANHOS DA COSTA, M. J. R., FAUCITANO, L., PELOSO, J. V.; DALLA ROZA, D. Efeito das condições pré-abate sobre a qualidade da carne de suínos pesados. **Archivos de Zootecnia**, v. 59, n. 227, p. 391-402, 2010.

DALLA COSTA, F. A.; DALLA COSTA, O. A. Preparo dos animais para o transporte. In: Associação Brasileira dos Criadores de Suínos (Coordenação editorial). **Manual de Industrialização dos Suínos**. Brasília: Associação Brasileira dos Criadores de Suínos, 2014a. p. 73-90.

DALLA COSTA, F. A.; DALLA COSTA, O. A. Manejo dos suínos no frigorífico. In: Associação Brasileira dos Criadores de Suínos (Coordenação editorial). **Manual de Industrialização dos Suínos**. Brasília: Associação Brasileira dos Criadores de Suínos, 2014b. p. 73-90.

DALMAU, A.; TEMPLE, D.; RODRIGUEZ, P.; LLONCH, P.; VELARDE, A. Application of the Welfare Quality (R) protocol at pig slaughterhouses. **Animal Welfare**, v. 18, n. 4, 497-505, 2009.

DAROIT, D. J.; BRANDELLI, A. Implications of skeletal muscle creatine kinase to meat quality. **Journal of Animal and Feed Sciences**, v. 17, n. 3, p. 285-294, 2008.

DAWKINS, M. S. Using behaviour to assess animal welfare. **Animal Welfare**, v. 13, p. 3-7, 2004.

DE JONG, I. C.; LAMBOOJI, E.; MECHIEL KORTE, S.; BLOCKHUIS, H. J. KOOLHAAS, J. M. Mixing induces long-term hyperthermia in growing pigs. **Animal Science**, v. 69, p. 601-605, 1999.

DE PASSILLÉ, A. M.; EHRlich, A. L.; WATKINS, L. R.; SPENCER, R. L.; MAIER, S. F.; LICINO, J.; WONG, M. L.; CHROUSOS, G. P.; WEBSTER, E.; GOLD, P. W. The impact of the nonpeptide corticotrophin-releasing hormone antagonist antalarmin on behavioral and endocrine responses to stress. **Endocrinology**, v. 149, n. 1, p. 79-86, 1995.

DE SMET, S. M.; PAUWELS, H.; DE BIE, S.; DEMEYER, D. I.; CALLEWIER, J.; EECKHOUT, W. Effect of Halothane Genotype, Breed, Feed Withdrawal, and Lairage on Pork Quality of Belgian Slaughter Pigs. **Journal of Animal Science**, v. 74, n. 8, p. 1854-1863, 1996.

DEEN, J. Pigs: Behavior and welfare assessment. In: BREED, M. D.; MOORE, J. **Encyclopedia of Animal Behavior**, Fort Collins: Academic Press, 2010. p. 731-739.

DEWEY, C.; HALEY, C.; WIDOWSKI, T.; POLJAK, Z.; FRIENDSHIP, R. Factors associated with in-transit losses of fattening pigs. **Animal Welfare**, v. 18, n. 4, p. 355-361, 2009.

DOKMANOVIC, M., VELARDE, A., TOMOVIĆ, V., GLAMOČLIJA, N., MARKOVIĆ, R., JANJIĆ, J., & BALTIĆ, M. Ž. The effects of lairage time and handling procedure prior to slaughter on stress and meat quality parameters in pigs. **Meat science**, v. 98, n. 2, p. 220-226, 2014.

DRIESSEN, B.; GEERS, R. Stress during transport and quality of pork. An European view. In: 1<sup>st</sup> INTERNATIONAL VIRTUAL CONFERENCE ON PORK QUALITY: WELFARE, TRANSPORT, SLAUGHTER AND CONSUME, 2000, Concordia. **Anais...** Concordia: EMBRAPA:CNPSA, 2001. p. 39-51.

D'SOUZA, D. N.; WARNER, R. D.; LEURY, B. J.; DUNSHEA, F. R. The effect of dietary magnesium aspartate supplementation on pork quality. **Journal of Animal Science**, v. 76, n. 1, p. 104-109, 1998.

EU COUNCIL DIRECTIVE 1993. **Council Directive (93/119/CEE) of 22 December 1993 on the protection of animals at the time of slaughter or killing**, 1993. Disponível em: <[http://ec.europa.eu/food/fs/aw/aw\\_legislation/slaughter/93-119-ec\\_en.pdf](http://ec.europa.eu/food/fs/aw/aw_legislation/slaughter/93-119-ec_en.pdf)>. Acessado em: 22 Dez. 2015.

EU COUNCIL DIRECTIVE 1995. **Animals during transport**. Official Journal European Communities, No. L340/17, p. 52-63, 1995.

EDWARDS, L. N.; GRANDIN, T.; ENGLE, T. E.; PORTER, S. P.; RITTER, M. J.; SOSNICKI, A. A.; ANDERSON, D. B. Use of exsanguination blood lactate to assess the quality of pre-slaughter pig handling. **Meat Science**, v. 86, n. 2, p. 384-390, 2010.

EUROPEAN FOOD SAFETY. Panel on Animal Health and Welfare (AHAW): Scientific Opinion on the use of animal-based measures to assess welfare in pigs. **EFSA Journal**, v. 10, n. 1:2512, p. 1-85, 2012.

EIKELENBOOM, G. In: PROCEEDINGS OF MEETING ON PIG CARCASS AND MEAT QUALITY, 1988, Bologna. **Anais...** Bologna: Universita di Bologna, 1988, p. 199-211.

EIKELENBOOM, G.; HOVING-BOLINK, A. H.; SYBESMA, W. Effect of feed withdrawal before delivery on pork quality and carcass yield. **Meat Science**, v. 29, n. 1, p. 25-30, 1991.

EKKEL, E. D.; SAVENIJE, B.; SCHOUTEN, W. G.; WIEGANT, V. M.; TIELEN, M. J.. The effects of mixing on behavior and circadian parameters of salivary cortisol in pigs. **Physiology & behavior**, v. 62, n. 1, p. 181-184, 1997.

EKESBO, I. **Farm animal behaviour: characteristics for assessment of health and welfare**. Wallingford: CABI, 2011. p. 248

ELLIS, M. Swine breeding, sex, feeding regime, and slaughter weight and their effects on carcass lean yield. In: *PROCEEDINGS OF THE WORKSHOP ON PORK CARCASS YIELD AND MEAT QUALITY, 1998, Concordia*. **Anais...** Concordia: EMBRAPA: CNPSA, 1998. p. 15-16.

ELLIS, M. Genetic and nutritional influences on pork quality. In: PROCEEDINGS OF THE WORKSHOP ON PORK CARCASS YIELD AND MEAT QUALITY, 1998, Concordia. **Anais...** Concordia: EMBRAPA: CNPSA, 1998. p. 25–54.

ELLIS, M.; MCKEITH, F.; HAMILTON, D.; BERTOL, T.; RITTER, M. Analysis of the current situation: what do downers cost the industry and what can we do about it? In: PROCEEDINGS OF THE 4<sup>th</sup> AMERICAN MEAT SCIENCE ASSOCIATION PORK QUALITY SYMPOSIUM, 2003, Columbia. **Anais...** Columbia: AMSA, 2003. p. 1-3.

ELLIS, M.; RITTER, M. J. Management practices and meat quality. In: PROCEEDING OF 59<sup>th</sup> AMERICAN MEAT ASSOCIATION – AMSA, 2006, Champaign-Urbana. **Anais...** Champaign-Urbana: AMSA, 2006. p. 29-32.

EUROPE COMISSION - E.U. Directive 2002/49/EC of the European parliament and the Council of 25 June 2002 relating to the assessment and management of environmental noise. **Official Journal of the European Communities**, v. 189, n. 12, 2002.

EUROPE COMISSION - Council Regulation 2009. Council Regulation (EC) No. 1099/2009 of 24 September 2009 on the production of animals at the time of killing.

EWBANK, R.; BRYANT, M. J. Aggressive behaviour amongst groups of domesticated pigs kept at various stocking rates. **Animal behaviour**, v. 20, n. 1, p. 21-28, 1972.

FAUCITANO, L. Causes of skin damage to pig carcasses. **Canadian Journal of Animal Science**, v. 81, n. 1, p. 39-45, 2001.

FAUCITANO, L. **PRINCIPAIS PROBLEMAS E SOLUÇÕES NO MANEJO PRÉ-ABATE DE SUÍNOS**. Concórdia. 2007. Palestra realizada no Caitá Hotel.

FAUCITANO, L. Effects of lairage and slaughter conditions on animal welfare and pork quality. **Canadian Journal of Animal Science**, v. 90, n. 4, p. 461-469, 2010.

FERNANDEZ, X.; MEUNIER-SALAUN, M-C.; ECOLAN, P.; MORMÈDE, P. Interactive effect of food deprivation and agonistic behaviour on blood parameters and muscle glycogen in pigs. **Physiology and Behaviour**, v. 58, n. 2, p. 337-345, 1995.

FAUCITANO, L.; MAQUARDT, M. S.; OLIVEIRA, M. S.; SEBASTIANY, H. S.; TERRA, N. N. The effect of two handling and slaughter systems on skin damage, meat acidification and colour in pigs. **Meat Science**, v. 50, n. 1, p. 13-19, 1998.

FAUCITANO, L.; SAUCIER, L.; CORREA, J. A.; MÉTHOT, S.; GIGUÈRE, A.; FOURY, A.; MORMÈDE, P.; BERGERON, R. Effect of feed texture, meal frequency and pre-slaughter fasting on carcass and meat quality, and urinary cortisol in pigs. **Meat Science**, v. 74, n. 4, p. 697-703, 2006.

FAUCITANO, L.; CHEVILLON, P.; ELLIS, M. Effects of feed withdrawal prior to slaughter and nutrition on stomach weight, and carcass and meat quality in pigs. **Livestock Science**, v. 127, n. 2, p. 110-114, 2010a.

FAUCITANO, L.; IELO, M. C.; STER, C.; FIEGO, D. P. L. O.; MÉTHOT, S.; SAUCIER, L. Shelf life of pork from five different quality classes. **Meat Science**, v. 84, n. 3, p. 466-469, 2010b.

FERNANDEZ, X.; TORNBERG, E. A review of the causes of variation in muscle glycogen content and ultimate pH in pigs. **Journal of Muscle Foods**, v. 2, n. 3, p. 209-235, 1991.

FERNANDEZ, X.; MEUNIER-SALAÜN, M. C.; ECOLAN, P. Glycogen depletion according to muscle and fibre types in response to dyadic encounters in pigs (*Sus scrofa domesticus*)-relationships with plasma epinephrine and aggressive behaviour. **Comparative Biochemistry and Physiology Part A: Physiology**, v. 109, n. 4, p. 869-879, 1994.

FERNANDEZ-LOPEZ, J.; PEREZ-ALVAREZ, J. A.; SAYAS-BARBERA, E. Characterization of the different states of myoglobin in pork using color parameters and reflectance ratios. **Journal of Muscle Foods**, v. 11, n. 3, p. 157-167, 2000.

FISHER, K. Improving the quality of meat for the consumer. **Animal Science Papers and reports**, v. 23, n.4, p. 269-280, 2005.

FITZGERALD, R. F.; STALDER, K. J.; MATTHEWS, J. O.; KASTER, S.; JOHNSON, A. K. Factors associated with fatigued, injured, and dead pig frequency during transport and lairage at a commercial abattoir. **Journal of Animal Science**, v. 87, n. 3, p. 1156-1166, 2009.

FORREST, J. C.; WILL, J. A.; SCHMIDT, G. R.; JUDGE, M. D.; BRISKEY, E. J. Homeostasis in animals (*Sus domesticus*) during exposure to a warm environment. **Journal of applied physiology**, v. 24, n. 1, p. 33-39, 1968.

FORTIN, A.; TONG, A. K. W.; ROBERTSON, W. M.; ZAWADSKI, S. M.; LANDRY, S. J.; ROBINSON, D. J.; LIU, T.; MOCKFORD, R. J. A novel approach to grading pork carcasses: computer vision and ultrasound. **Meat science**, v. 63, n. 4, p. 451-462, 2003.

FOX, J.; WIDOWSKI, T.; TORREY, S.; NANNO, E.; BERGERON, R.; GONYOU, H. W.; BROWN, J. A.; CROWE, T.; MAINAU, E.; FAUCITANO, L. Water sprinkling market pigs in a stationary trailer. 1. Effects on pig behaviour, gastrointestinal tract temperature and trailer micro-climate. **Livestock Science**, v. 160, p. 113-123, 2014.

FRANCK, M.; FIGWER, PH.; JOSSEL, A.; POIREL, M. T.; PASTEUR, X. Incidence of preslaughter stress on meat quality of non sensible pigs. **Revue de médecine vétérinaire**, v. 154, n. 3, p. 199-204, 2003.

FRAQUEZA, M. J.; ROSEIRO, L. C.; ALMEIDA, J.; MATIAS, E.; SANTOS, C.; RANDALL, J. M. Effects of lairage temperature and holding time on pig behaviour and on carcass and meat quality. **Applied Animal Behaviour Science**, v. 60, n. 4, p. 317-330, 1998.

FRASER, A. F.; BROOM, D. M. **Farm animal behaviour and welfare**. Cornell: CABI Publishing, 1997. p. 437.

GARCIA, A.; MCGLONE, J. J. Loading and Unloading Finishing Pigs: Effects of Bedding Types, Ramp Angle, and Bedding Moisture. **Animals**, v. 5, n. 1, p. 13-26, 2014.

GEESING L. M.; JOHNSON, A. K.; SELSBY, J. T.; FEUERBACH, C.; HILL, H.; FAGA, M.; WHILEY, A.; BAILEY, R.; STALDER, K. J.; RITTER, M. J. Effects of presorting on stress responses at loading and unloading and the impact on transport losses from market-weight pigs. **The Professional Animal Scientist**, v. 26, n. 6, p. 603-610, 2010.

GEVERINK, N. A.; ENGEL, B.; LAMBOOIJ, E.; WIEGANT, V. M. Observations on behaviour and skin damage of slaughter pigs and treatment during lairage. **Applied Animal Behaviour Science**, v. 50, n. 1, p. 1-13, 1996.

GEVERINK, A.; BUHNEMANN, A.; VAN DE BURG WAL, J. A.; LAMBOOIJ, E.; BLOKHUIS, H. J.; WIEGANT, V. M. Responses of slaughter pigs to transport and lairage sounds. **Physiology & Behavior**, v. 63, n. 4, p. 667-673, 1998.

GILBERT, K. V.; DEVINE, C. E.; HAND, R.; ELLERY, S. Electrical stunning and stillness of lambs. **Meat Science**, v. 11, n. 1, p. 45-58, 1984.

GISPERT, M.; FAUCITANO, L.; OLIVER, M. A.; GUÀRDIA, M. D.; COLL, C.; SIGGENS, K.; HARVEY, K.; DIESTRE, A. A survey of pre-slaughter conditions, halothane gene frequency, and carcass and meat quality in Five Spanish pig commercial abattoirs. **Meat Science**, v. 55, n. 1, p. 97-106, 2000.

GISPERT, M., M.; i FURNOLS, M. F; GIL, M.; VELARDE, A.; DIESTRE, A.; CARRIÓN, D.; SOSNICKI, A. A.; PLASTOW, G. S. Relationships between carcass quality parameters and genetic types. **Meat Science**, v. 77, n. 3, p. 397-404, 2007.

GOUMON, S.; FAUCITANO, L.; BERGERON, R.; CROWE, T.; CONNOR, M. L.; GONYOU, H. W. Effect of ramp configuration on easiness of handling, heart rate, and behavior of near-market weight pigs at unloading. **Journal of Animal Science**, v. 91, n. 8, p. 3889-3898, 2013.

GRANDIN, T. Design of loading facilities and holding pens. **Applied of Animal Behaviour Science**, v. 28, n. 1, p. 187-201, 1990.

GRANDIN, T. Handling and transport of agricultural animals used in research. In: PROCEEDINGS, SCIENTISTS' CENTER FOR ANIMAL WELFARE, 1991, Washington. **Anais...** Washington, 1991.

GRANDIN, T. The feasibility of using vocalization scoring as an indicator of poor welfare during cattle slaughter. **Applied Animal Behaviour Science**, v. 56, n. 2, p. 121-128, 1998.

GRANDIN, T. Handling and welfare of livestock in slaughter plants. In: GRANDIN, T. **Livestock Handling and Transport**. 2. ed. Wallingford: CAB International, 2000. cap. 20, p. 409-439.

GRANDIN, T. Welfare of cattle during slaughter and the prevention of nonambulatory (downer) cattle. **Journal of the American Veterinary Medical Association**, v. 219, n. 10, p. 1377, 2001.

GRANDIN, T. Engineering and design of holding yards, loading ramps and handling facilities for land and sea transport of livestock. **Veterinaria italiana**, v. 44, n. 1, p. 235-245, 2008.

GRANDIN, Temple. Welfare of Pigs During Transport (Updated November 2014). Acesso em: 19/01/2016. <<http://www.grandin.com/welfare.pigs.during.transport.html>>

GREGORY, N.G. **Animal Welfare and Meat Science**. New York: CABI Publishing, 1998. p. 298.

GREGORY, N.G. **Physiology and Behaviour of Animal Suffering**. UK: Blackwell Publishing, 2004. p. 268.

GREGORY, N. G.; VON WENZLAWOWICZ, M.; VON HOLLEBEN, K. Blood in the respiratory tract during slaughter with and without stunning in cattle. **Meat science**, v. 82, n. 1, p. 13-16, 2009.

GRUNERT, K. G. Food quality and safety: consumer perception and demand. **European Review of Agricultural Economics**, v. 32, n. 3, p. 369-391, 2005.

GUÀRDIA, M. D.; GISPERT, M.; DIESTRE, A. Mortality rates during transport and lairage in pigs for slaughter. **Meat Focus International**, v. 10, p. 362–366, 1996.

GUÀRDIA, M. D.; ESTANY, J.; BALASH, S.; OLIVER, M. A.; GISPERT, M.; DIESTRE, A. Risk assessment of PSE condition due to pre-slaughter conditions and RYR1 gene in pigs. **Meat Science**, v. 67, n. 3, p. 471-478, 2004.

GUÀRDIA, M. D.; ESTANY, J.; BALASH, S.; OLIVER, M. A.; GISPERT, M.; DIESTRE, A. Risk assessment of DFD condition due to pre-slaughter conditions and RYR1 gene in pigs. **Meat Science**, v. 70, n. 3, p. 709–716, 2005.

GUÀRDIA, M. D.; ESTANY, J.; BALASCH, S.; OLIVER, M. A.; GISPERT, M.; DIESTRE, A. Risk assessment of skin damage due to pre-slaughter conditions and RYR1 gene in pigs. **Meat Science**, v. 81, n. 4, p. 745-751, 2009.

GUISE, H. J.; PENNY, R. H. C. Factors influencing the welfare and carcass and meat quality of pigs. 1. The effects of stocking density in transport and the use of electric goads. **Animal Production**, v. 49, n. 3, p. 511-515, 1989.

HALEY, C.; DEWEY, C. E.; WIDOWSKI, T.; FRIENDSHIP, R. Association between in-transit losses, internal trailer temperature, and distance travelled by Ontario market hogs. **Canadian Journal of Veterinary Research**, v. 72, n. 5, p. 385-389, 2008a.

HALEY, C.; DEWEY, C. E.; WIDOWSKI, T.; POLJAK, Z.; FRIENDSHIP, R. Factors associated with in-transit losses of market hogs in Ontario in 2001. **Canadian Journal of Veterinary Research**, v. 72, n. 5, p. 377, 2008b.

HAMBRECHT, E.; EISSEN, J. J.; NEWMAN, D. J.; SMITS, C. H. M.; VERSTEGEN, M. W. A.; DEN HARTOG, L. A. Preslaughter handling effects on pork quality and glycolytic potential in two muscles differing in fiber type composition. **Journal of Animal Science**, v. 83, n. 4, p. 900–907, 2005.

HAMILTON, D. N.; ELLIS, M.; BRESSNER, G. E.; WOLTER, B. F.; JONES, D. J.; WATKINS, L. E. Relationships between environmental conditions on trucks and losses during transport to slaughter in finishing pigs. **Journal of Animal Science**, v. 81, n. 2, p. 53, 2003. Edição Suplementar.

HAMILTON, D. N.; ELLIS, M.; BERTOL, T. M.; MILLER, K. D. Effects of handling intensity and live weight on blood acid-base status in finishing pigs. **Journal of Animal Science**, v. 82, n. 8, p. 2405-2409, 2004.

HARRIS, R. A. Glycolysis, Overview. In: LENNARZ, W. J.; LANE, M. D. **Encyclopedia of Biological Chemistry**. Indiana: University School of Medicine, 2004. p. 266-271.

HAY, M.; MEUNIER-SALAÜN, M. C.; BRULAUD, F.; MONNIER, M.; MORMÈDE, P. Assessment of hypothalamic-pituitary-adrenal axis and sympathetic nervous system activity in pregnant sows through the measurement of glucocorticoids and catecholamines in urine. **Journal of Animal Science**, v. 78, n.2, p. 420-428, 2000.

HELD, S.; COOPER, J. J.; MENDEL, M.T. Advances on the study of cognition, behavioural priorities and emotions. In: MARCHANT-FORDE, J. N. **Animal Welfare**. 2. ed. Wallingford: CABI Publishing, 2009. cap. 3, p. 47-94.

HEMSWORTH, P. H.; MELLOR, D. J.; CRONIN, G. M.; TILBROOK, A. J. Scientific assessment of animal welfare. **New Zealand Veterinary Journal**, v. 63, n.1, p. 24-30, 2015.

HOENDERKEN, R. **Electrical stunning of slaughter pigs**. 1978. XXX f. Ph.D. thesis - University of Utrecht, the Netherlands, 1978.

HOLMER, S. F.; MCKEITH, R. O.; BOLER, D. D.; DILGER, A. C.; EGGERT, J. M.; PETRY, D. B.; MCKEITH, F. K.; JONES, K. L.; KILLEFER, J. The effect of pH on shelf-life of pork during aging and simulated retail display. **Meat Science**, v. 82, n. 1, p. 86-93, 2009.

HONIKEL, K. O. Reference methods for the assessment of physical characteristics of meat. **Meat Science**, v. 49, n. 4, p. 447-457, 1998.

HONIKEL, K. O. Water-holding capacity of meat. In: Te PAS, M.F.W.; Everts, M.E.; Haagsman, H.P. **Muscle development of livestock animals**. Wallingford: CABI Publishing, 2004. p. 389-399.

HOUPT, K. A.; BALDWIN, B. A.; HOUPT, R.; HILLS, F. Humoral and cardiovascular responses to feeding in pigs. **American Journal of Physiology-Regulatory, Integrative and Comparative Physiology**, v. 244, n. 2, p. 279-284, 1983.

HUFF-LONERGAN, E.; LONERGAN, S. M. Mechanisms of water-holding capacity of meat: The role of *post-mortem* biochemical and structural changes. **Meat Science**, v. 71, 194-204, 2005.

HUGHES, B.O. Behaviour as Index of Welfare. In: PROCEEDINGS V<sup>th</sup> EUROPEAN POULTRY CONFERENCE, 1976, Malta. **Anais...** Malta, 1976. p. 1005-1018.

HUMANE METHODS OF SLAUGHTER ACT. **United States Congress**. PL 95-445, 1978.

INGRAM, D. L. Evaporative cooling in the pig. **Nature**, v. 207, p. 415-416, 1965.

ITP. 1996. Notation des hématomes sur couenne - porcs vivant ou carcasse. Institute Technique du Porc, Rennes, France.

JENSEN, P. Observations on the maternal behaviour of free-ranging domestic pigs. **Applied Animal Behaviour Science**, v. 16, n. 2, p. 131-142, 1986.

JENNEN, D. G. J.; BRINGS, A. D.; LIU, G.; JUENGST, H.; THOLEN, E.; JONAS, E.; TEFAYE, D.; SCHELLANDER, K.; PHATSARA, C. Genetic aspects concerning drip loss and water holding capacity of porcine meat. **Journal of Animal Breeding and Genetics**, v. 124, n. 1, p. 2-11, 2007.

JOHNSON, J. M.; PROPPE, D. W. Cardiovascular Adjustments to Heat Stress. In: JOHNSON J. M.; PROPPE, D. W. **Handbook of Physiology, Environmental Physiology**. 14. ed. New York: Oxford University Press, 1996. p. 215-243.

JOHNSON, A. K.; GESING, L. M.; ELLIS, M.; MCGLONE, J. J.; BERG, E.; LONERGAN, S. M.; FITZGERALD, R.; KARRIKER, L. A.; RAMIREZ, A.; STALDER, K. J.; SAPKOTA, A.; KEPHART, R.; SELSBY, J. T.; SADLER, L. J.; RITTER, M. J.. 2011 AND 2012 EARLY CAREERS ACHIEVEMENT AWARDS: Farm and pig factors affecting welfare during the marketing process. **Journal of Animal Science**, v. 91, n. 6, p. 2481-2491, 2013.

JOO, S. T.; KAUFFMAN, R. G.; KIM, B. C.; PARK, G. B. The relationship of sarcoplasmic and myofibrillar protein solubility to colour and water-holding capacity in porcine longissimus muscle. **Meat science**, v. 52, n. 3, p. 291-297, 1999.

JONG, I. C.; SGOIFO, A.; LAMBOOIJ, E.; KORTE, S. M.; BLOKHUIS, H. J.; KOOLHAAS, J. M. Effects of social stress on heart rate and heart rate variability in growing pigs. **Canadian Journal of Animal Science**, v. 80, n. 2, p. 273-280, 2000.

JONES, S. D. M., ROMPALA, C.R.; HAWORTH, C. R. Effects of fasting and water restriction on carcass shrink and pork quality. **Canadian Journal of Animal Science**, v. 65, n. 3, p. 613-618, 1985

JONES, S. D. M.; CLIPLEF, R. L.; FORTIN, A.; MCKAY, R. M.; MURRAY, A. C.; POMMIER, S. A.; SATHER, A. P.; SCHAEFER, A. L. Production and *antemortem* factors influencing pork quality. **Pig News Info**, v. 15, n. 1, p. 15-18, 1994.

JONES, T. A. Improved handling systems for pigs at slaughter. 1999. Ph.D. thesis, Royal Veterinary College, University of London, London, 1999.

JONGMAN, E. C.; BARNETT, J. L.; HEMSWORTH, P. H. The aversiveness of carbon dioxide stunning in pigs and a comparison of the CO<sub>2</sub> stunner crate vs. the V-restrainer. **Applied Animal Behaviour Science**, v. 67, n. 1, p. 67-76, 2000.

KAUFFMAN, R. G.; EIKELNBOOM, G.; VAN DERWAL, P. G.; MERKUS, G.; ZAAR, M. The use of filter paper to estimate drip loss in porcine musculature. **Meat Science**, v. 18, n. 3, p. 191-200, 1986.

KAUFFMAN, R. G.; WARNER, R. D. Evaluating pork carcasses for composition and quality. In: HOLLIS, G. R. **Gowth of the pig**. 1. ed. Wallingford: Cab International, 1993. p. 256.

KEPHART, K. B.; MILLS, E. W. Effect of withholding feed from swine before slaughter on carcass and viscera weights and meat quality. **Journal of Animal Science**, v. 83, n. 3, p. 715-721, 2005.

KEPHART, K. B.; HARPER, M. T.; RAINES, C. R. Observations of market pigs following transport to a packing plant. **Journal of Animal Science**, v. 88, n. 6, p. 2199-2203, 2010.

KNOWLES, T. G.; BROWN, S. N.; EDWARDS, J. E.; WARRISS, P. D. Ambient temperature below which pigs should not be continuously showered in lairage. **The Veterinary Record**, v. 143, n. 21, p. 575-578, 1998.

KÖHLER, R. G. **Estudo da redução do tempo de permanência na pocilga do frigorífico sobre a qualidade da carne suína**. 2001. 77 f. Dissertação (Mestrado em Tecnologia de Alimentos - Setor de Tecnologia), Universidade Federal do Pará, Curitiba, 2001.

KÜCHENMEISTER, U.; KUHN, G.; ENDER, K. Preslaughter handling of pigs and the effect on heart rate, meat quality, including tenderness, and sarcoplasmic reticulum Ca<sup>2+</sup> transport. **Meat Science**, v. 71, n. 4, p. 690-695, 2005.

LAMBOOIJ, E.; SYBESMA, W. The effect of environmental factors such as preslaughter treatment and electrical stunning on the occurrence of haemorrhages in the shoulder of slaughter pigs. In: Proceedings of the 34<sup>th</sup> International Congress of Meat Science and Technology, 1998, Brisbane. **Anais...** Brisbane: ICoMST, 1988. p. 1-3.

LAMMENS, V., E. PEETERS, H. DE MAERE, E. DE MEY, H. PAELINCK, J. LEYTEN, AND R. GEERS. A survey of pork quality in relation to pre-slaughter conditions, slaughterhouse facilities, and quality assurance. **Meat science**, v. 75, n. 3, p. 381-387, 2007.

LATORRE, M. A.; LÁZARO, R.; VALENCIA, D. G.; MEDEL, P.; MATEOS, G. G. The effects of gender and slaughter weight on the growth performance, carcass traits, and meat quality characteristics of heavy pigs. **Journal of Animal Science**, v. 82, n. 2, p. 526-533, 2004.

LATORRE, M. A.; LÁZARO, R.; GRACIA, M. I.; NIETO, M.; MATEOS, G. G. Effect of sex and terminal sire genotype on performance, carcass characteristics, and meat quality of pigs slaughtered at 117 kg body weight. **Meat science**, v. 65, n. 4, p. 1369-1377, 2003.

LAWRIE, R. E. Metabolic stress which affect muscle. In: **BRISKEY, E. J.; CASSENS, R. G.; TRAUTMAN, J. C.** The Physiology and Biochemistry of Muscle as a Food. Madison: University of Wisconsin Press, 1966. p. 437.

- LEE, J. R.; KIM, D. H.; HUR, T. Y.; LEE, J. I.; JOO, S. T.; PARK, G. B. The effect of stocking density in transit on the meat quality and blood profile of slaughter pig. **Korean Journal of Animal Science**, v. 42, n. 5, p. 669-676, 2000.
- LEHESKA, J. M.; WULF, D. M.; MADDOCK, R. J. Effects of fasting and transportation on pork quality development and extent of *post-mortem* metabolism. **Journal of Animal Science**, v. 80, n. 12, p. 3194-3202, 2003.
- LEWIS, C. R. G.; McGLONE, J. J. Moving finishing pigs in different group sizes: Cardiovascular responses, time and ease of handling. **Livestock Science**, v. 107, n. 1, p. 86-90, 2007.
- LIDFORS, L.; BERG, C.; ALGERS, B. Integration of natural behavior in housing systems. **AMBIO: A Journal of the Human Environment**, v. 34, n. 4, p. 325-330, 2005.
- LO FIEGO, D.P., Minelli, G and Santoro, P. Effect of pre-slaughter fasting on the meat quality traits of crossbred pigs. In: PROCEEDINGS OF SOCIETA' ITALIANA DELLE SCIENZE VETERINARIE, 1997, Bologna. **Anais...** Bologna: SISVET, 1997, p. 699-700.
- LONG, V. P.; TARRANT, P. V. The effect of pre-slaughter showering and post-slaughter rapid chilling on meat quality in intact pork sides. **Meat Science**, v. 27, n. 3, p. 181-195, 1990.
- LØVENDAHL, P., DAMGAARD, L.H., NIELSEN, B.L., THODBERG, K., SU, G., RYDHMER, L. Aggressive behaviour of sows at mixing and maternal behaviour are heritable and genetically correlated traits. **Livestock Production Science**, v. 93, p. 73-85, 2005.
- LYNCH, P. B.; LAWLOR, P. G.; DAVIS, D.; KERRY, J. P.; BUCKLEY, D. J.; WALSH, L. **Studies on pre-slaughter handling of pigs and its relationship to meat quality**. Moorepark Food Research Centre, 1998.
- MANCINI, R. A.; HUNT, M. C. Current research in meat color. **Meat Science**, v. 71, p. 100-121, n. 1, 2005.
- MATTERI, R. L., CARROLL, J. A., DYER, C. J. Neuroendocrine responses to stress. In: MOBERG, G.; MENCH, J. A. **The biology of animal stress: basic principles and implications for animal welfare**. 1. ed. Davis: University of California, 2000. cap. 3, p.43-76.
- McGLONE, J. J.; MCPHERSON, R. L.; ANDERSON, D. L. Case study: moving devices for finishing pigs: efficacy of electric prod, board, paddle, or flag. **The Professional Animal Scientist**, v. 20, n. 6, p. 518-523, 2004.
- MEESE, G. B.; EWBANK, R. A note on instability of the dominance hierarchy and variations in level of aggression within groups of fattening pigs. **Animal Production**, v. 14, n. 03, p. 359-362, 1972.

MEESE, G. B.; EWBANK, R. The establishment and nature of the dominance hierarchy in the domesticated pig. **Animal Behaviour**, v. 21, n. 2, p. 326-334, 1973.

MELLOR, D. J.; MURRAY, L. Effects of tail docking and castration on behaviour and plasma cortisol concentrations in young lambs. **Research in Veterinary Science**, v. 46, n. 3, p. 387-391, 1989.

MILLS, D. S.; MARCHANT-FORDE, J. N.; MCGREENY, P. D.; MORON, D. B.; NICOL, C. J.; PHILLIPS, D. J. C.; SANDØE, P.; SWAISGOUD, R. R. **The Encyclopedia of Applied Animal Behaviour and Welfare**. Wallingford: CABI, 2010. p. 685

MILLER, M. F.; CARR, M. A.; BAWCOM, D. B.; RAMSEY, C. B.; THOMPSON, L. D. Microbiology of pork carcasses from pigs with differing origins and feed withdrawal times. **Journal of Food Protection**, v. 60, n. 3, p. 242-245, 1997.

MILLIGAN, S. D.; RAMSEY, C. B.; MILLER, M. F.; KASTER, C. S.; THOMPSON, L. D. Resting of pigs and hot-fat trimming and accelerated chilling of carcasses to improve pork quality. **Journal of Animal Science**, v. 76, n. 1, p. 74-86, 1998.

MITCHELL, M. A. Thermal imaging in physiology: theoretical and practical considerations. In: LUZI, F.; MITCHELL, M. A.; NANNI COSTA, L.; REDAELLI, V. **Thermography: Current Status and Advances in Livestock Animals and in Veterinary Medicine**. Italy: Fondazione Iniziative Zooprofilattiche e Zootecniche, Brescia, 2013. p. 47-65.

MLC. 1985. Concern at rindside damage in pigs. **Meat and Marketing Technical Notes**, n. 4, pp 14-16, Milton Keynes, Meat and Livestock Commission, Bletchley, UK.

MOBERG, G. P. Problems in defining stress and distress in animals. **Journal of the American Veterinary Medical Association**, v. 191, n. 10, p. 1207-1211, 1987.

MOBERG, G. P. Biological response to stress: Implications for animal welfare. In: MOBERG, G. P.; MENCH, J. A. **The biology of animal stress – Basic principles and implications for animal welfare**. 1. ed. Wallingford: CABI Publishing, 2000. cap. 1. p 1-22.

MORMÈDE, P. 2008. Assessment of pig welfare. In: FAUCITANO, L.; SCHAEFER, A. L. **The welfare of Pigs: from Birth to Slaughter**. The Netherlands: Wageningen Academic Publishing, 2008. cap. 2, p. 33-64.

MOSS, B. W. Some observations on the activity and aggressive behaviour of pigs when penned prior to slaughter. **Applied Animal Ethology**, v. 4, n. 4, p. 323-339, 1978.

MOSS, B. W.; ROBB, J. D. The effect of preslaughter lairage on serum thyroxine and cortisol levels at slaughter, and meat quality of boars, hogs and gilts. **Journal of the Science of Food and Agriculture**, v. 29, n. 8, p. 689-696, 1978.

MOUNT, L. E. Environmental physiology in relation to pig production. In: COLE, D. J. A. **Pig Production**. London: Butterworths Publishing, 1972.

MURRAY, C.; JONES, S. D.; SATHER, A. P. The effect of preslaughter feed restriction and genotype for stress susceptibility on pork lean quality and composition. **Canadian Journal of Animal Science**, v. 69, n. 1, p. 83-91, 1989.

MURRAY, A. C.; JONES, S. D. M. The effect of mixing, feed restriction and genotype with respect to stress susceptibility on pork carcass and meat quality. **Canadian Journal of Animal Science**, v. 74, n. 4, p. 587-594, 1994.

MURRAY, A. C.; JOHNSON, C. P. Impact of the halothane gene on muscle quality and pre-slaughter deaths in Western Canadian pigs. **Canadian Journal of Animal Science**, v. 78, n. 4, p. 543-548, 1998.

MURRAY, A.; ROBERTSON, W.; NATTRESS, F.; FORTIN, A. Effect of pre-slaughter overnight feed withdrawal on pig carcass and muscle quality. **Canadian Journal of Animal Science**, v. 81, n. 1, p. 89-97, 2001.

NANNI COSTA, L.; LO FIEGO, D. P.; DALL'OLIO, S.; DAVOLI, R.; RUSSO, V. Influence of loading method and stocking density during transport on meat and dry-cured ham quality in pigs with halothane genotypes. **Meat Science**, v. 51, n. 4, p. 391-399, 1999.

NANI COSTA, L.; FIEGO, D. L.; DALL'OLIO, S.; DAVOLI, R.; RUSSO, V. Combined effects of pre-slaughter treatments and lairage time on carcass and meat quality in pigs of different halothane genotype. **Meat Science**, v. 61, n. 1, p. 41-47, 2002.

NANNONI, E.; WIDOWSKI, T.; TORREY, S.; FOX, J.; ROCHA, L. M.; GONYOU, H.; Weschenfelder, A. V.; CROWE, T.; MARTELLI, G.; FAUCITANO, L. Water sprinkling market pigs in a stationary trailer. 2. Effects on selected exsanguination blood parameters and carcass and meat quality variation. **Livestock Science**, v. 160, p. 124-131, 2014.

NAKAI, H.; SAITO, F.; IKEDA, T.; ANDO, S.; KOMATSU, A. Standards models of pork color. **Bulletin of National Institute of Animal Industry**, v. 30, p. 69-74, 1975.

NICOLAIEWSKY, S. Fatores produtivos que afetam a qualidade da carne suína. In: SIMPÓSIO SOBRE RENDIMENTO E QUALIDADE DA CARNE SUÍNA, 1998, Concórdia. **Anais...** Concórdia: EMBRAPA Suínos e Aves, 1998. p. 1-4. (EMBRAPA suínos e aves. Documentos, 51).

NORMAN, A. W.; LITWACK, G. Adrenal Corticoids. In: NORMAN, A. W.; LITWACK, G. **Hormones**. 2. ed. San Diego: Elsevier Academic Press, 1997. cap. 10, p. 281-339.

NOWAK, B.; MUEFFLING, T. V.; HARTUNG, J. Effect of different carbon dioxide concentrations and exposure times in stunning of slaughter pigs: impact on animal welfare and meat quality. **Meat Science**, v. 75, n. 2, p. 290-298, 2007.

NIELSEN, N. J. Recent results from investigations of transportation of pigs for slaughter. In: MOSS, R. **Transport of animals intended for breeding, production and slaughter**. 18. ed. The Hague: Springer Netherlands, 1982. cap. 3, p. 115-124.

NYBERG, L.; LUNDSTROM, K.; EDFORS LILJA, L.; RUNDGREN, M. Effects of transport stress on concentrations of cortisol, corticosteroid-binding globulin and glucocorticoid receptors in pigs with different halothane genotypes. **Journal of Animal Science**, v. 66, n. 5, p. 1202-1211, 1988.

O'CONNELL, N. E.; BEATTIE, V. E. Influence of environmental enrichment on aggressive behaviour and dominance relationships in growing pigs. **Animal Welfare**, v. 8, n. 3, p. 269-279, 1999.

OFFER, G.; COUSINS, T. The mechanism of drip production: formation of two compartments of extracellular space in muscle *post-mortem*. **Journal of the Science of Food and Agriculture**, v. 58, n. 1, p. 107-116, 1992.

OLSSON, I. A. S.; WÜRBEL, H.; MENCH, J. A. Behaviour. In: APPLEBY, M. C.; MENCH, J. A.; OLSSON, I. A. S. **Animal Welfare**. 2. ed. Wallingford: CABI, 2011. p. 138-154.

OWEN, B. L.; MONTGOMERY, J. L.; RAMSEY, C. B.; MILLER, M. F. Preslaughter resting and hot-fat trimming effects on the incidence of pale, soft and exudative (PSE) pork and ham processing characteristics. **Meat science**, v. 54, n. 3, p. 221-229, 2000.

PANELLA-RIERA, N.; GISPERT, M.; GIL, M.; SOLER, J.; TIBAU, J.; OLIVER, M. A.; VELARDE, A.; FÀBREGA, E. Effect of feed deprivation and lairage time on carcass and meat quality traits on pigs under minimal stressful conditions. **Livestock Science**, v. 146, n. 1, p. 29-37, 2012.

PARANHOS DA COSTA, M. J. R.; OLIVEIRA, D.; RUEDA, P. M. Conceitos de bem-estar animal e implicações práticas no manejo reprodutivo. In: OLIVEIRA, M. E. F. Oliveira; Teixeira, P. P. A.; Vicente, W. R. R. **Biotécnicas Reprodutivas em Ovinos e Caprinos**. 1. ed. São Paulo: Editora MedVet, 2013. v. 1, p. 293-305.

PARROTT, R. F.; MISSON, B. H. Changes in pig salivary cortisol in response to transport simulation, food and water deprivation, and mixing. **British Veterinary Journal**, v. 145, n. 6, p. 501-505, 1989.

PATIENCE, J. F.; UMBOH, T. J. F.; CHAPLIN, R. K.; NYACHOTI, C.M.; PARK, B.Y.; LEE, C.E.; KIM, I.S.; CHO, S.H.; KIM, Y.G.; LEE, J.M.; YOON, S.K. Nutritional and physiological responses of growing pigs exposed to a diurnal pattern of heat stress. **Livestock Production Science**, v. 96, p. 205-214, 2003.

PEETERS, E.; DRIESSEN, B.; STEEGMANS, R.; HENOT, D.; GEERS, R. Effect of supplemental tryptophan, vitamin E, and a herbal product on responses by pigs to vibration. **Journal of animal science**, v. 82, n. 8, p. 2410-2420, 2004.

PELOSO, J. V. Influência do jejum pré-abate sobre a condição muscular em suínos e seus efeitos na qualidade final da carne para industrialização. In: CONFERÊNCIA VIRTUAL INTERNACIONAL SOBRE QUALIDADE DA CARNE SUÍNA, v.2. 2001, Concórdia. **Anais...** Concórdia: EMBRAPA Suínos e Aves, 2002. p. 385-392.

PERSON, R. C.; MCKENNA, D. R.; ELLEBRACHT, J. W.; GRIFFIN, D. B.; MCKEITH, F. K.; SCANGA, J. A.; BELK, K. E.; SMITH, G. C.; SAVELL, J. W. Benchmarking value in the pork supply chain: Processing and consumer characteristics of hams manufactured from different quality raw materials. **Meat Science**, v. 70, n. 1, p. 91-97, 2005.

PETHERICK, J. C. A biological basis for the design of space in livestock housing. In: BAXTER, S. H.; BAXTER M. R.; MacCORMACK, J. A. D. **Farm animal housing and welfare**. 1. ed. The Netherlands: Kluwer Academic Publishers Group, 1983. p. 103-120.

PILCHER, C. M.; ELLIS, M.; ROJO-GÓMEZ, A.; CURTIS, S. E.; WOLTER, B. F.; PETERSON, C.; RITTER, M. J.; BRINKMANN, J. Effects of floor space during transport and journey time on indicators of stress and transport losses of market-weight pigs. **Journal of Animal Science**, v. 89, n. 11, p. 3809-3818, 2011.

PITTS A. D.; WEARY, D. M.; PAJOR E. A.; FRASER D. Mixing at young ages reduces fighting in unacquainted domestic pigs. **Applied Animal Behaviour Science**, v. 68, n. 3, p. 191-197, 2000.

POSO, A. R.; PUOLANNE, E. Carbohydrate metabolism in meat animals. **Meat Science**, v. 70, n. 3, p. 423-434, 2005.

POMAR, C.; MARCOUX, M.; GISPERT, M.; FONTI FURNOLS, M.; DAUMAS, G. Determining the lean content of pork carcasses. In: KERRY, J. P.; LEDWARD, D. **Improving the sensory and nutritional quality of fresh meat**. 1. ed. Cambridge: Woodhead Publishing, 2008. cap. 21, p. 493-513.

RABASTE, C.; FAUCITANO, L.; SAUCIER, L.; MORMÈDE, P.; CORREA, J. A.; GIGUÈRE, A.; BERGERON, R. The effects of handling and group size on welfare of pigs in lairage and their influence on stomach weight, carcass microbial contamination and meat quality. **Canadian Journal of Animal Science**, v.87, n. 1, p. 3-12, 2007.

RAJ, A. B. M. Welfare of pigs during stunning and slaughter. In: FAUCITANO, L.; SCHAEFER, A. L. **The welfare of pigs: From birth to slaughter**. 1. ed. The Netherlands: Academic Publisher, 2008. cap. 8, p. 225-240.

RHIM, S. J.; SON, S. H.; HWANG, H. S.; LEE, J. K.; HONG, J. K. Effects of Mixing on the Aggressive Behavior of Commercially Housed Pigs. **Asian Australasian Journal of Animal Science**, v. 28, n. 7, p. 1038-1043, 2015.

RODRIGUEZ, A.; DE MERCADO, E.; GÓMEZ, E.; ALEVIA, A.; CHICO, M.; FLORES, L.; SANZ, E. Influencia de la duración del ayuno y de la espera antes del sacrificio sobre los rendimientos productivos y características de la canal de cerdos de cebo. **Avances en tecnología porcina**, v. 5, n. 9, p. 32-42, 2008.

RITTER, M. J. **Effects of animal handling and transportation factors on the welfare, stress responses, and incidences of transport losses in market weight pigs at the packing plant**. 2007. Dissertação de Mestrado - University of Illinois, Urbana, Champaign, 2007.

RITTER, M. J.; ELLIS, M.; BRINKMANN, J.; DEDECKER, J. M.; KOCHER, M. E.; KEFFABER, K. K.; PETERSON, B. A.; SCHLIPF, J. M.; WOLTER, B. F. Effect of floor space during transport of market weight pigs on incidence of transport losses (dead and non-ambulatory pigs) at the packing plant and relationships between transport conditions and losses. **Journal of animal science**, v. 84, n. 10, p. 2856-2864, 2006.

RITTER, M. J.; ELLIS, M.; BERTELSEN, C.R.; BOWMAN, R.; BRINKMANN, J.; DEDECKER, J. M.; KEFFABER, K. K.; MURPHY, C. M.; PETERSON, B. A.; SCHLIPF, J. M.; WOLTER, B. F. Effects of distance moved during loading and floor space on the trailer during transport on losses of market weight pigs on arrival at the packing plant. **Journal of animal science**, v. 85, n. 12, p. 3454-3461, 2007.

RITTER, M. J.; ELLIS, M.; BOWMAN, R.; BRINKMANN, J.; CURTIS, S. E.; DEDECKER, J. M.; MENDOZA, O.; MURPHY, C. M.; ORELLANA, D. G.; PETERSON, B. A.; ROJO, A.; SCHLIPF, J. M.; WOLTER, B. F. Effects of season and distance moved during loading on transport losses of market-weight pigs in two commercially available types of trailer. **Journal of Animal Science**, v. 86, n. 11, p. 3137-3145, 2008.

RITTER, M. J.; ELLIS, M.; BERRY, N. L.; CURTIS, S. E.; ANIL, L.; BERG, E.; BENJAMIN, M.; BUTLER, D.; DEWEY, C.; DRIESSEN, B.; DUBOIS, P.; HILL, J. D.; MARCHANT-FORDE, J. N.; MATZAT, P.; MCGLONE, J. J.; MORMEDE, P.; MOYER, T.; PFALZGRAF, K.; SALAK-JOHNSON, J.; SIEMENS, M.; STERLE, J.; STULL, C.; WHITING, T.; WOLTER, B.; NIEKAMP, S. R.; JOHNSON, A. K. Transport losses in market weight pigs: A review of definitions, incidence, and economic impact. **The Professional Animal Scientist**, v. 25, n. 4, p. 404-414, 2009.

ROBIN, S.; COURDEROT-MASUYER, C.; TAUZIN, H.; HARBON, S.; CHAVAGNAC-BONNEVILLE, M.; CADARS, B.; JOURDAN, E.; TROMPEZINSKI, S.; HUMBERT, P. Use of a model of a blood-induced bruise for the evaluation of formulations on bruising. **Journal of Cosmetics, Dermatological Sciences and Applications**, v. 5, n. 1, p. 7-14, 2015.

ROCHA, L. M.; DIONNE, A.; SAUCIER, L.; NANNONI, E.; FAUCITANO, L. Hand-held lactate analyzer as a tool for the real-time measurement of physical fatigue before slaughter and pork quality prediction. **Animal**, v. 9, n. 4, p. 707-714, 2015.

RUIS, M. A. W.; TE BRAKE, J. H. A.; ENGEL, B.; EKKELE, E. D.; BUIST, W. G.; BLOKHUIS, H. J.; KOOLHAAS, J. M. The Circadian Rhythm of Salivary Cortisol in Growing Pigs: Effects of Age, Gender, and Stress. **Physiology & Behavior**, v. 62, n. 3, p. 623–630, 1997.

RUIS, M. A.; TE BRAKE, J. H.; ENGEL, B.; BUIST, W. G.; BLOKHUIS, H. J.; KOOLHAAS, J. M. Adaptation to social isolation: acute and long-term stress responses of growing gilts with different coping characteristics. **Physiology & Behavior**, v. 73, n. 4, p. 541-551, 2001.

RYU, Y. C.; KIM, B. C. The relationship between muscle fiber characteristics, *post-mortem* metabolic rate, and meat quality of pig longissimus dorsi muscle. **Meat Science**, v. 71, n. 2, p. 351-357, 2005.

SALAK-JOHNSON, J., McGLONE, J. Making sense of apparently conflicting data: Stress and immunity in swine and cattle. **Journal of Animal Science**, v. 85, n. 13, p. 81-88, 2007.

SANTOS, C.; ALMEIDA, J. M.; MATIAS, E. C.; FRAQUEZA, M. J.; ROSEIRO, C.; SARDINA, L. Influence of lairage environmental conditions and resting time on meat quality in pigs. **Meat Science**, v. 45, n. 2, p. 253-262, 1997.

SAUCIER, L., BERNIER, D., BERGERON, R., GIGUERE, A., METHOT, S., FAUCITANO, L. Effect of feed texture, meal frequency and pre-slaughter fasting on behaviour, stomach content and carcass microbial quality in pigs. **Canadian Journal of Animal Science**, v. 87, p. 479-487, 2007.

SAVAGE, A. W. J; WARRISS, P. D.; JOLLEY, P. D. The amount and composition of the proteins in drip from stored pig meat. **Meat Science**, v. 27, n. 4, p. 289-303, 1990.

SCHAEFER, A. L.; MATTHEWS, L. R.; COOK, N. J.; WEBSTER, J.; SCOTT, S. L. Novel non-invasive measures of animal welfare. In: JOINT NAWAC/ISAE CONFERENCE OF ANIMAL WELFARE AND BEHAVIOUR: FROM SCIENCE TO SOLUTION, 2002, Hamilton. **Anais...** Hamilton: ISAE, 2002.

SCAHAW. **The Welfare of Animals During Transport (details for Horses, Pigs, Sheep and Cattle)**. Report of the Scientific Committee on Animal Health and Animal Welfare: European Commission, 2002. 130 p.

SCHWARTZKOPF-GENSWEIN, K. S. G.; FAUCITANO, L.; DADGAR, S.; SHAND, P.; GONZÁLEZ, L. A.; CROWE, T. G. Road transport of cattle, swine and poultry in North America and its impact on animal welfare, carcass and meat quality: A review. **Meat Science**, v. 92, n. 3, p. 227-243, 2012.

SCHRAMA, J. W.; van der HER, W.; GORSSSEN, J.; HENKEN, A. M.; VERSTEGEN, M. W. A.; NOORDHUIZEN, J. P. T. M. Required thermal thresholds during transport of animals. **Veterinary Quarterly**, v. 18, n. 3, p. 90-95, 1996.

SCROGGS, L. V.; KATTESH, H. G.; MORROW, J. L.; STALDER, K. J.; DAILEY, J. W.; ROBERTS, M. P.; SCHNEIDER, J. F.; SAXTON, A. M. The effects of split marketing on the physiology, behavior, and performance of finishing swine. **Journal of animal science**, v. 80, n. 2, p. 338-345, 2002.

SELLIER, P. Genetics of meat and carcass traits. In: ROTHSCHILD, M. F.; RUVINSKI, A. **The genetics of the pig**. 2. ed. Cambridge: CABI Publishing, 1998. cap. 15, p. 355-389.

SELLIER, P.; G. MONIN. Genetics of pig meat quality: a review. **Journal Muscle Foods**, v. 5, n. 2, p. 187- 219, 1994.

SHEN, Q. W.; MEANS, W. J.; THOMPSON, S. A.; UNDERWOOD, K. R.; ZHU, M. J.; McCORMICK, R. J.; FORD, S. P.; DU, M. Pre-slaughter transport, AMP-activated protein kinase, glycolysis, and quality of pork loin. **Meat Science**, v. 74, n. 2, p. 388-395, 2006.

SMITH, W. W. **Marketing: Market classes and grades**. New York: The MacMillan Company, 1937. p. 450

STEWART, G.; RITTER, M.; CULBERTSON, M.; MANN, G.; WOFFORD, R. Effects of previous handling and feed withdrawal prior to loading on transport losses in market weight pigs. In: PROCEEDINGS OF AMERICAN ASSOCIATION OF SWINE VETERINARIANS, 2008, San Diego. **Anais...** San Diego: AASV, 2008. p. 359-362.

STOOKEY, J. M.; GONYOU, H. W. The effects of regrouping on behavioral and production parameters in finishing swine. **Journal of animal science**, v. 72, n. 11, p. 2804-2811, 1994.

SUTHERLAND, M.A.; McDONALD, A.; McGLONE, J. J. Effects of variations in the environment, length of journey and type of trailer on the mortality and morbidity of pigs being transported to slaughter. **Veterinary Records**, v. 165, n. 1, 13-18, 2009.

SUTHERLAND, M. A.; ERLANDSON, K.; CONNOR, J. F.; SALAK-JOHNSON, J. L.; MATZAT, P.; SMITH, J. F.; McGLONE, J. J. Health of non-ambulatory, non-injured pigs at processing. **Livestock Science**, v. 116, n. 1, p. 237-245, 2008.

TARRANT, P. V. Transportation of cattle by road. **Applied Animal Behaviour Science**, v. 28, n. 1, p. 153-170, 1990.

TARRANT, P. V. The effects of handling, transport, slaughter and shilling on meat quality and yield in pigs – a review. **Irish Journal of Food Science and Technology**, v.13, n. 2, p.70–107, 1989.

TARRANT, P. J. V.; MCLOUGHLIN, J. V.; HARRINGTON, M. G. Anaerobic glycolysis in biopsy and *post-mortem* porcine *longissimus dorsi* muscle. In: PROCEEDINGS OF THE ROYAL IRISH ACADEMY. SECTION B: BIOLOGICAL, GEOLOGICAL, AND CHEMICAL SCIENCE, 72., 1972, .Dublin. **Anais...** Dubling: Royal Irish Academy, 1972. p. 55-73.

TERLOUW, C.; ASTRUC, T.; PORCHER, J.; RYBARCZYK, P.; FERNANDEZ, X. Transport and other potential stress factors at slaughter: effects of genetic and rearing background. In: INTERNATIONAL SOCIETY FOR ANIMAL HYGIÈNE, 2004, Saint-Malo. **Anais...** Saint-Malo: ISAH, 2004. p. 35-36. Acessado em: 19/01/2015. <<http://www.isah-soc.org/documents/2004/Terlouw.pdf>> Accessed on April 25, 2012.

TERRA, N. N. **Apontamentos de tecnologia de carnes**. São Leopoldo: Unisinos, 1998. p. 216

TORREY, S.; BERGERON, R.; WIDOWSKI, T.; LEWIS, N.; CROWE, T.; CORREA, J. A.; BROWN, J.; GONYOU, H. W.; FAUCITANO, L. Transportation of market-weight pigs: I. Effect of season and truck type on behavior with a 2h transport. **Journal of Animal Science**, v. 91, n. 6, p. 2863-2871, 2013a.

TORREY, S.; BERGERON, R.; FAUCITANO, L.; WIDOWSKI, T.; LEWIS, N.; CROWE, T.; CORREA, J. A.; BROWN, J.; HAYNE, S.; GONYOU, H. W. Transportation of market-weight pigs: II. Effect of season and location within truck on behavior with an eight-hour transport. **Journal of animal science**, v. 91, n. 6, p. 2872-2878, 2013b.

TERRIEN, J.; PERRET, M.; AUJARD, F. Behavioral thermoregulation in mammals: a review. **Frontiers in Bioscience**, n. 16, p. 1428-1444, 2011.

TOLDRA, F. Muscle foods: water, structure and functionality. **Food science and technology international**, v. 9, n. 3, p. 173-177, 2003.

TURGEON, M. J. **Évaluation de différents scénarios de mise à jeun avant l'abattage sur les performances zootechniques, le comportement et la qualité de la viande**. 2006. Dissertação de Mestrado (Mestrado em Zootecnia) - Université Laval, Laval Sainte-Foy, 2003.

TURNER, S. P.; EDWARDS, S. A.; BLAND, V. C. The influence of drinker allocation and group size on the drinking behaviour, welfare and production of growing pigs. **Animal Science**, v. 68, p. 617-624, 1999.

TURNER, A. I., KEATING C, TILBROOK, A. J. Sex differences and the role of sex steroids in sympatho-adrenal medullary system the hypothalamo-pituitary adrenal axis responses to stress. In: KAHN S. M. **Sex Steroids**. 1. ed. Rijeka: Tech Publishing, 2012. cap. 6, p 115-36.

ULRICH-LAI, Y. M.; HERMAN, J. P. Neural regulation of endocrine and autonomic stress responses. **Nature Reviews Neuroscience**, v. 10, n. 6, p. 397-409, 2009.

VAN DE PERRE, V.; PERMENTIER, L.; DE BIE, S.; VERBECKE, G.; GEERS, R. Effect of unloading, lairage, pig handling, stunning and season on pH of pork. **Meat Science**, v. 86, n. 4, p. 931-937, 2010.

VAN DER WAL, P. G.; ENGEL, B.; REIMERT, H. G. M. The effect of stress, applied immediately before stunning, on pork quality. **Meat Science**, v. 53, n. 2, p. 101-106, 1999.

VAN LAACK, R. L. J. M.; KAUFFMAN, R. G.; SYBESMA, W.; SMULDERS, F. J. M.; EIKELENBOOM, G.; PINHEIRO, J. C. Is colour brightness (L-value) a reliable indicator of water-holding capacity in porcine muscle? **Meat Science**, v. 38, n. 2, p. 193-201, 1994.

VAN PUTTEN, G.; ELSHOF, W. J. Observations on the effect of transport on the well-being and lean quality of slaughter pigs. **Animal Regulation Studies**, v. 1, n. 3, p. 247-271, 1978.

VECEREK, V.; MALENA, M.; MALENA JR., M.; VOŠLAROVA, E; CHLOUPEK, P. The impact of the transport distance and season on losses of fattened pigs during transport to the slaughterhouse in the Czech Republic in the period from 1997 to 2004. **Veterinarni Medicina**, v. 51, n. 1, p. 21, 2006.

VELARDE, A.; GISPERT, M.; FAUCITANO, L.; ALONSO, P.; MANTECA, X.; DIESTRE, A. Effects of the stunning procedure and the halothane genotype on meat quality and incidence of haemorrhages in pigs. **Meat Science**, v. 58, n. 3, p. 313-319, 2001.

VELARDE, A.; GISPERT, M.; FAUCITANO, L.; DIESTRE, A.; MANTECA, X. Survey of the effectiveness of stunning procedures used in Spanish pig abattoirs. **Veterinary Record**, v. 146, n. 3, p. 65–68, 2000a.

VELARDE, A.; GISPERT, M.; FAUCITANO, L.; MANTECA, X.; DIESTRE, A. The effect of stunning method on the incidence of PSE meat and haemorrhages in pork carcasses. **Meat Science**, v. 55, n. 3, p. 309-314, 2000b.

VIAU, J.; CHAMPAGNE, D. La mise à jeun chez le porc avant l'abattage. **Report of Centre de développement du Porc du Québec**, Quebec City, Canada, 1998.

VOET, D.; VOET, J.G. **Biochemistry**. New York: Headington Hill Hall, 1995. p. 104

Von BORELL, E.; SCHAFFER, D. Legal requirements and assessment of stress and welfare during transportation and pre-slaughter handling of pigs. **Livestock Production Science**, v. 97, n. 2, p. 81-87, 2005.

Von HOLST, D. The concept of stress and its relevance for animal behaviour. In: MOLLER, A. P.; MILINSKI, M.; Slater, P. J. B. **Stress and Behavior**. 1. ed. California: Academic Press, 1998. vol. 27. p. 1-132.

Von MICKWITZ, G. Various transport conditions and their influence on physiological reactions. In: MOSS, R. **Transport of animals intended for breeding, production and slaughter**. 1. ed. Netherlands: Kluwer Academic Publishers, 1982. v. 18 p. 45-56.

WAIBLINGER, S.; BOIVIN, X.; PEDERSEN, V.; TOSI, M. V.; JANCZAK, A. M.; VISSER, E. K.; JONES, R. B. Assessing the human–animal relationship in farmed species: a critical review. **Applied Animal Behaviour Science**, v. 101, n. 3, p. 185-242, 2006.

WALLIMANN, T.; WYSS, M.; BRDICZKA, D.; NICOLAY, K.; EPPENBERGER, H. M. Intracellular compartmentation, structure and function of creatine kinase isoenzymes in tissues with high and fluctuating energy demands: the "phosphocreatine circuit" for cellular energy homeostasis. **Biochemical Journal**, v. 281, n. 1, p. 21-40, 1992.

WARNER, R.D. **Physical properties of porcine musculature in relation to post-mortem biochemical changes in muscle proteins**. 1994. PhD Thesis, University of Wisconsin – Madison, 1994.

WARNER, R. D.; KAUFFMAN, R. G.; GREASER, M. L. Muscle protein changes *post mortem* in relation to pork quality traits. **Meat science**, v. 45, n. 3, p. 339-352, 1997.

WARNER, R. D.; KAUFFMAN, R. G.; RUSSELL, R. L. Quality attributes of major porcine muscles: A comparison with the *Longissimus lumborum*. **Meat Science**, v. 33, n. 3, p. 359-372, 1993.

WARRISS, P. D. Loss of carcass weight, liver weight and liver glycogen, and the effects on muscle glycogen and ultimate pH in pigs fasted pre-slaughter. **Journal of the Science of Food and Agriculture**, v. 33, n. 9, p. 840-846, 1982.

WARRISS, P. D. Marketing losses caused by fasting and transport during the preslaughter handling of pigs. **Pig News and Information**, v. 6, p. 155-157, 1985.

WARRISS, P. D. The effect of time and conditions of transport and lairage on pig meat quality. In: TARRANT, P. V.; EIKELENBOOM, G.; MONIN, G. **Evaluation and control of meat quality in pigs**. 1. ed. Boston: Martinus Nijhoff Publisher, 1987. p. 245-264.

WARRISS, P. D. Handling pigs before slaughter and the consequences for welfare and product quality. **Meat Focus International**, v.1, p.135-138, 1992.

WARRISS, P. D. The consequences of fighting between mixed groups of unfamiliar pigs before slaughter. **Meat Focus International**, p.89-92, 1996.

WARRISS, P. D. The welfare of slaughter pigs during transport. **Animal Welfare**, v. 7, n. 4, p. 365-381, 1998.

WARRISS, P. D. Optimal lairage times and conditions for slaughter pigs: a review. **Veterinary Records**, v. 153, n. 6, p. 170-176, 2003.

WARRISS, P. D. The growth and body composition of animals. In: WARRISS, P. D. **Meat Science: an introductory text**. 2. ed. Massachusetts: CABI, 2010a. cap. 2, p. 9-25.

WARRISS, P. D. Animal welfare. In: WARRISS, P. D. **Meat Science: an introductory text**. 2. ed. Massachusetts: CABI, 2010b. cap. 10, p. 209-228.

WARRISS, P. D.; BROWN, S. N. The physiological responses of fighting between in pigs and the consequences for meat quality. **Journal of the Science and Food Agriculture**, v. 86, n. 2, p. 87-92, 1985.

WARRISS, P. D.; BROWN, S. N. The relationships between initial pH, reflectance and exudation in pig muscle. **Meat Science**, v. 20, n. 1, p. 65-74, 1987.

WARRISS, P. D.; BEVIS, E. A.; EDWARDS, J. E.; BROWN, S. N.; KNOWLES, T. G. Effect of angle of slope on the ease with which pigs negotiate loading ramps. **Veterinary Records**, v. 128, n. 18, p. 419-421, 1991.

WARRISS, P. D.; BROWN, S. N.; EDWARDS, J. E.; ANIL, M. H.; FORDHAM, D. P. Time in lairage needed by pigs to recover from the stress of transport. **Veterinary Record**, v. 131, n. 9, p. 194-196, 1992.

WARRISS, P. D.; BROWN, S. N.; ADAMS, S. J. M.; CORLETT, I. K. Relationships between subjective and objective assessments of stress at slaughter and meat quality in pigs. **Meat Science**, v. 38, n. 2, p. 329-340, 1994.

WARRISS, P. D.; BROWN, S. N. A survey of mortality in slaughter pigs during transport and lairage. **The Veterinary Record**, v. 134, n. 20, p. 513-515, 1994.

WARRISS, P. D.; BROWN, S. N.; KNOWLES, T. G.; EDWARDS, J. E.; KETTLEWELL, P. J.; GUISE, H. J. The effect of stocking density in transit on the carcass quality and welfare of slaughter pigs: 2. Results from the analysis of blood and meat samples. **Meat Science**, v. 50, n. 4, p. 447-456, 1998a.

WARRISS, P. D.; BROWN, S. N.; EDWARDS, J. E.; KNOWLES, T. G. Effect of lairage time on levels of stress and meat quality in pigs. **Animal Science**, v. 66, n. 1, p. 255-261, 1998b.

WEBB, P. A. U. L. The physiology of heat regulation. **American Journal of Physiology-Regulatory, Integrative and Comparative Physiology**, v. 268, n. 4, p. 838-850, 1995.

WEEDING, C. M.; GUISE, H. J.; PENNY, R. H. C. Factors influencing the welfare and carcass and meat quality of pigs: the use of water sprays in lairage. **Animal Production**, v. 56, n. 3, p. 393-397, 1993.

WEEKS, C. A. A review of welfare in cattle, sheep, and pig lairages, with emphasis on stocking rates, ventilation and noise. **Animal Welfare**, v. 17, n. 3, p. 275-284, 2008.

WELFARE QUALITY (2009). Assessment protocol for pig. Disponível em: <http://www.welfarequality.net>. Acesso em: 26 de outubro de 2015.

WERNER, C.; REINERS, K.; WICKE, M. Mortality rates during transport of slaughter pigs. **Fleischwirtschaft**, v. 85, n. 9, p. 133-136, 2005.

WESCHENFELDER, A. V. **Animal welfare and meat quality in pigs as affected by trailer type, travel distance and genotype**. 2013. 207 f. PhD Thesis - Université Laval, Laval Sainte-Foy, 2013.

WESCHENFELDER, A. V.; TORREY, S.; DEVILLERS, N.; CROWE, T.; BASSOLS, A.; SACO, Y.; PIÑEIRO, M.; SAUCIER, L.; FAUCITANO, L. Effects of trailer design on animal welfare parameters and carcass and meat quality of three Pietrain crosses being transported over a long distance. **Journal of Animal Science**, v. 90, p. 3220-3231, 2012.

WESCHENFELDER, A. V.; TORREY, S.; DEVILLERS, N.; CROWE, T.; BASSOLS, A.; SACO, Y.; PIÑEIRO, M.; SAUCIER, L.; FAUCITANO, L. Effects of trailer design on animal welfare parameters and carcass and meat quality of three Pietrain crosses being transported over a short distance. **Livestock Science**, v. 157, n. 1, p. 234-244, 2013.

WESCHENFELDER, A. V.; SAUCIER, L.; MALDAGUE, X.; ROCHA, L. M.; SCHAEFER, A. L.; FAUCITANO, L. Use of infrared ocular thermography to assess physiological conditions of pigs prior to slaughter and predict pork quality variation. **Meat science**, v. 95, n. 3, p. 616-620, 2013.

WEYMAN, G. **Unloading and loading facilities at livestock markets**. Council of National and Academic Awards, Environmental Studies, Hatfield Polytechnic, Hatfield, 1987.

WILLIAMS, R. B JR. Neuroendocrine response patterns and stress: Biobehavioral mechanisms of disease. In: WILLIAMS, R. B. **Perspectives on Behavioral Medicine: Neuroendocrine Controls and Behavior**. 2 ed. New York: Academic press, 1983. cap. 5, p. 71-101.

WOLTERS DORF, W.; TROEGER, K. Slaughter technique to reduce proportion of PSE pigs. **Fleischwirtschaft**, v. 10, p. 1248-1251, 1987.

WOTTON, S. B.; ANIL, M. H.; WHITTINGTON, P. E.; MCKINSTY, J. L. Pig slaughtering procedures: head-to-back stunning. **Meat Science**, v. 32, n. 3, p. 245-255, 1992.

WOTTON, S. B. New advances in stunning techniques for slaughter animals. **Meat Focus International**, v. 12, p. 461-465, 1996a.

WOTTON, S. B. Sticking techniques and exsanguination in pigs, sheep and calves. **Meat Focus International**, v. 7, p. 234-237, 1996b.

WYSS M.; SMEITINK J.; WEVERS R. A.; WALLIMANN T. Mitochondrial creatine kinase: a keyenzyme of aerobic energy metabolism. **Biochimica et Biophysica Acta**, v. 1102, n. 2, p. 119-166, 1992.

YOUNG, O. A.; WEST, J. Meat Color. In: HUI, Y. H.; NIP, W. K., ROGERS, W. R.; YOUNG, O. A. **Meat Science and Applications**. 1. ed. New York: Marcel Dekker, 2001. cap. 3, p. 39-70.

ZHANG, W., KUHLLERS, D.L.; REMPEL, W.E. Halothane gene and swine performance. **Journal of Animal Science**, v. 70, p. 1307-1313, 1992.

**EFFECTS OF APPLYING PRESLAUGHTER FEED WITHDRAWAL AT THE  
ABATTOIR ON BEHAVIOUR, BLOOD PARAMETERS AND MEAT QUALITY IN  
PIGS**

Artigo submetido para a revista: Meat Science.

ISSN: 0309-1740

Normas para publicação disponíveis em:

[https://www.elsevier.com/wps/find/journaldescription.cws\\_home/405866?generationpdf=true](https://www.elsevier.com/wps/find/journaldescription.cws_home/405866?generationpdf=true)

## CHAPTER 2 - EFFECTS OF APPLYING PRESLAUGHTER FEED WITHDRAWAL AT THE ABATTOIR ON BEHAVIOUR, BLOOD PARAMETERS AND MEAT QUALITY IN PIGS

F. A. Dalla Costa<sup>a,b</sup>, N. Devillers<sup>c</sup>, M. J. R. Paranhos da Costa<sup>a,b</sup> and L. Faucitano<sup>c,\*</sup>

<sup>a</sup> *Programa de Pós-Graduação em Zootecnia, Faculdade de Ciências Agrárias e Veterinárias, UNESP, Jaboticabal, SP, Brazil*

<sup>b</sup> *Departamento de Zootecnia, Faculdade de Ciências Agrárias e Veterinárias, UNESP, 14884-900 Jaboticabal, SP, Brazil*

<sup>c</sup> *Agriculture and Agri-Food Canada, 2000 College Street, J1M 0C8 Sherbrooke, Quebec, Canada*

\*Corresponding author:

Tel.: +1 819 7807237; fax: +1 819 5645507

E-mail address: luigi.faucitano@agr.gc.ca (L. Faucitano).

### ABSTRACT

To study the effects of starting time of feed withdrawal on behaviour during loading and lairage, and meat quality (pH, colour and drip loss) in pigs, a total of 700 pigs was distributed into two groups. One group (FARM) was fasted for 18 h at the farm before transport and for 6 h before slaughter (including 2 h transport and 4 h lairage), while for the other group (PLANT) fasting time started at the departure from the farm (2 h transport plus 22 h lairage). Total fasting time was 24 h before slaughter for both groups. Fasting treatments neither influenced transport losses nor blood parameters. PLANT pigs had longer fights ( $P < 0.05$ ) than FARM pigs during lairage and produced darker ( $P = 0.02$ ) and drier ( $P = 0.03$ ) loins. Based on the results of this study, it

may be preferable to apply the most part of feed withdrawal time at the farm.

*Keywords:* Behaviour, fasting, meat quality, pigs, stress

## 1. Introduction

Feed withdrawal is a recommended practice for on-farm preparation of pigs before slaughter in order to prevent animal losses or travel sickness during transport (Averós, Knowles, Brown, Warriss, & Gonsálvez, 2008; Bradshaw et al., 1996; National Farm Animal Care Council, 2014), to reduce carcass contamination due to lower risk of gut contents spillage during carcass evisceration (Saucier et al., 2007) and to improve pork quality (Faucitano, Chevillon, & Ellis, 2010). A feed withdrawal time between 16 and 24 h appears to be an acceptable compromise to obtain optimal carcass yield and pork quality and safety (Eikelenboom, Hoving-Bolink, & Sybesma, 1991; Faucitano et al., 2010). The application of this on farm fasting interval has been also reported to decrease the proportion of dead-on-arrival (DOA; Guàrdia, Gispert, & Diestre, 1996) and total transport losses (DOA and non-ambulatory pigs; Stewart, Ritter, Culbertson, Mann, & Wofford, 2008).

Despite these potential advantages, however, fasting is sometimes only applied from the departure from the farm to the abattoir in Canadian swine farms (Aalhus et al., 1992; Viau, & Champagne, 1998). The reason for this misapplication is the producers' concern about the risk of slower growth rate of pigs remaining in the pen and the lack of a shipping room where the heaviest pigs can be transferred to ahead of loading time and feed be withdrawn at the same time (Viau & Champagne, 1998). If, on one hand, this practice may still prevent the risk of full stomachs at slaughter, provided the 16-24 h fasting time is respected by imposing longer lairage time at the abattoir, on the other hand, it may reduce the welfare of pigs during transport and handling (Eikelenboom et al., 1991; Guàrdia et al., 1996; Stewart et al., 2008). Furthermore, longer lairage time is known to result in higher risk of fighting in mixed groups of pigs leading to higher incidence of dark, firm and dry (DFD) pork (Faucitano et al., 2010; Guàrdia et al., 2005, 2009).

The objectives of this study were to study the effects of starting time of fasting preslaughter,

either on-farm or at the departure from the farm, on behaviour during loading and lairage, blood parameters and carcass and meat quality in pigs.

## **2. Material and methods**

All experimental procedures performed in this study were approved by the institutional animal care committee on the basis of the current guidelines of the Canadian Council on Animal Care (2009).

### *2.1. Animals and treatments*

In this study, a total of 700 crossbred pigs raised at a commercial swine growing-finishing farm and fed a standard commercial diet were randomly selected, tagged, weighed and distributed into two groups of 350 pigs each (BW:  $123.7 \pm 4.0$  kg and  $123.2 \pm 3.9$  kg;  $P = 0.25$ ) on the day before the departure from the farm to the abattoir. The first group (FARM) had feed withdrawn for 18 h at the farm before the transport (feeder locked at 1 pm), loaded at 7 am, transported for 2 h and kept in lairage at the abattoir for 4 h before slaughter, whereas the second group (PLANT) had access to the feeder until start of loading (1 pm), was transported for 2 h and kept in lairage at the abattoir for 22 h. Overall, a 24 h feed withdrawal time prior to slaughter was applied for both groups. At the time of loading for each transport (7 trips in terms of one per week) one batch of 50 pigs per fasting treatment was formed by collecting pigs randomly (or mixed) from 11 pens within each treatment based on the slaughter weight. Pigs were thus shipped to the abattoir over 7 slaughter days (weeks or replicates; 50 pigs x 2 treatments x 7 replicates). Pigs were loaded onto two similar trucks (one truck/treatment) in groups of 5 pigs by a trained handler using paddles and boards and transported at a loading density of 0.43 m<sup>2</sup>/pig. The handler at the farm, and the two drivers and trucks were the same throughout the 7 weeks. A randomization of truck loading order and drivers was done at each load to avoid the confounding effect of the truck and driver on the response of the pigs to

transport, as reported in previous studies (Peeters et al., 2008; Schwartzkopf-Genswein et al., 2012).

On arrival at the abattoir, pigs were unloaded using a whip only and driven to separate lairage pens on the basis of the treatment group as they were in the truck (25 pigs; no mixing). In each lairage pen, pigs were kept at a stocking density of 0.58 m<sup>2</sup>/pig. At the end of the lairage period, pigs were driven to a CO<sub>2</sub> stunner and exsanguinated in the prone position.

## *2.2. Behavioural observations*

### *2.2.1. Behaviour at loading*

Behaviour of pigs during loading was recorded using three digital camcorders (DCR-HC48; Sony, Sony of Canada Ltd., Toronto, Canada), two installed in the alley to the loading ramp and one in the loading ramp. All occurrences of pig behaviours and handler interventions (Table 1) were counted for each loading group of 5 pigs from a predetermined start gate in the farm alley up to the trailer gate for an overall distance of 12 m. Video recordings were analysed by two trained observers using a handheld Psion Workabout computer (HC-110, Psion Inc., Mississauga, Canada). Inter-observer agreement was 91.8%. The total time taken to load the treatment pigs from the starting gate to the trailer gate averaged 25.6±3 min.

### *2.2.2. Behaviour during lairage*

Behaviour during lairage was recorded using video cameras (WV-BP50, Panasonic Canada Inc., Mississauga, Canada) installed over the pens and connected to a digital encoder (Nextiva S5712e, Verint, Melville, NY). Images were captured by the Omnicast system (version 4.0; Genetec Inc., Montreal, Canada) at a frequency of 5 to 7 images/sec during two periods of lairage which excluded the time of showering (first 20 min and last 40 min of lairage time) to

avoid the confounding effect of this practice on pig behavior (Weeding, Guise, & Penny, 1993). The first period was of 2 and 3 h for FARM and PLANT pigs, respectively, and started from the end of the first shower, while the second period covered the last 2 h of lairage before the start of the last shower. Scan sampling was used at 2 min intervals to determine the number of pigs lying down. Number and duration of fights were determined at the group level using continuous sampling. A fight was considered when two or more animals performed a sequence of the following behaviours for more than 3 sec.: biting, head knocking, pushing and shoving each other, with no greater intervals than 10 sec. between behaviour occurrences (D'Eath, 2002; Pitts, Weary, Pajor, & Fraser, 2000). Proportion of pigs lying down, and number and duration of fights were calculated per period of 30 min. in two ways: forward from the beginning of lairage and backward prior to slaughter. To assess the recovery rate of the pigs after transport, the time necessary for 75% of the pigs from the same pen to rest (latency to rest) was also determined by the percentage of pigs lying down. Video recordings were analysed by two trained observers and inter-observer agreement was 94.2%.

### *2.3. Blood parameters*

Blood samples were collected from the bleeding wound of 210 pigs (15 pigs/treatments/replicates) in a plastic cup and lactate values were immediately assessed in duplicate with the Lactate Scout Analyzer (Lactate Scout, EKF Diagnostic GmbH, Magdeburg, Germany) by dipping two test strips (two strips/animal) in the collected sample. Later on, 10 mL of blood were collected in a tube (BD Vacutainers; VWR International Ltd., Montreal, Canada) to extract the serum for creatine kinase (CK) analysis. Serum samples were kept at room temperature (~23°C) for 1 h before refrigeration at 4°C. The next day, serum samples were centrifuged at 4°C for 12 min at 1,400 g, transferred to Eppendorf tubes

(1.5 mL), and stored at  $-80^{\circ}\text{C}$  until analysis. CK concentrations were measured with a creatine kinase-sl kit (Creatine Kinase-SL Assay of Chemicals Diagnostic Limited, Vancouver, Canada). The intra-assay coefficient of variation for CK was 5.49 %.

#### 2.4. Carcass and meat quality measures

After slaughter, carcasses were eviscerated, split and chilled according to standard commercial practices. Hot carcass weight (HCW) was recorded and used to calculate carcass yield. Lean yield was obtained by measuring carcass fat and lean depth at the 3<sup>rd</sup>/4<sup>th</sup> last rib level by a Destron optical probe.

Meat quality was assessed in the carcass of the same pigs that were blood sampled (210 or 15 carcasses/treatments/replicates) by measuring pH at 1 h (pH<sub>1</sub>) and 24 h (pH<sub>24</sub>) *post-mortem* with a pH meter (Oakton Instruments Model pH 100 Series, Nilis, IL) fitted with a spear type electrode (Cole Palmer Instrument Company, Vernon Hills, IL) and an automatic temperature compensation probe by insertion in the *Longissimus thoracis* (LT) between the 3<sup>rd</sup>/4<sup>th</sup> last rib (Canadian grading site), *Semimembranosus* (SM) and in the *Adductor* (AD) muscles 24 h *post-mortem* the following measurements were conducted in the LT muscle in the carcass grading site region: subjective color score using the Japanese Color Standards (JCS) ranging from 1 = pale to 6 = dark color (Nakai, Saito, Ikeda, Ando, & Komatsu, 1975), instrumental colour (L\*, a\* and b\* values) using a Minolta Chromameter (CR-300; Minolta Canada Inc., Mississauga, Canada) equipped with a 25-mm aperture, 0° viewing angle, and D65 illuminant, and drip loss using the modified EZ-driploss method as describe by Correa, Méthot, & Faucitano (2007). Briefly, 25 mm diameter cores were removed from the center of 2.5 cm thick LT muscle cross-section, weighed and placed into plastic drip loss containers (Christensen Aps Industrivaengetand, Hilleroed, Denmark), before being stored for 48h at 4°C.

At the end of the 48 h storage period, muscle cores were removed from their container, surface moisture was carefully dabbed, cores were reweighed, and drip loss percentage was calculated by dividing the difference between initial and final core weights by the initial core weight.

### *2.5. Statistical analysis*

Statistical analyses were carried out according to a generalized randomized complete block design to check any effects of treatments by analysis of variance in one-way ANOVA. The MIXED procedure of SAS (SAS Inst. Inc., Cary, NC) was applied to analyze the physiological data using the group as the experimental unit and meat quality data using the individual animal as the experimental unit. Values of blood lactate and CK concentration were log-transformed ( $\log_{10}$ ) for data normalization before analysis. Loading behaviour data were analyzed with the FREQ procedure using the loading group as the experimental unit. As behaviours, such as back-up, 180° turn, go backward, slip/fall, underlapping, vocalizing, squeeze and noise, had low frequencies, they were transformed to binary variables and were analyzed with the LOGISTIC procedure of SAS. Lairage behaviour data were analyzed as repeated measures over time (periods of 30 min.) after transformation (angular, square root and logarithmic transformations for percentage of pigs lying down, number of fights and total duration of fights, respectively). The model included treatment, time period and their interaction as fixed effects, while week was used as a repetition. Results from lairage behaviour are presented as back-transformed least square means and confidence interval. A probability level of  $P < 0.05$  was chosen as the limit for statistical significance in all tests. Whereas, probability levels of  $P \leq 0.10$  were considered as a tendency.

### 3. Results and Discussion

#### 3.1. Behavioural response

##### 3.1.1. Behaviour at loading

The fasting condition of pigs at the time of loading affected their behaviour while being moved through the farm alley to the truck gate (Table 2). Compared with FARM pigs, PLANT pigs showed less behaviours of going backward and 180° turn ( $P = 0.03$  for both), and vocalized less ( $P = 0.05$ ) during loading. However, these behaviours did not result in different frequency of interventions of handler between the two groups ( $P > 0.10$ ). The use of a trained handler and the small group size in this study may have compensated for these fear behaviours and ease of handling. Overall, these observations do not confirm the lower easiness of handling of pigs moved on full stomach reported by Eikelenboom et al. (1991). The behaviour showed by FARM pigs at loading may be explained by the increased frustration and excitement caused by hunger and feed restriction (Arnone, & Dantzer, 1980; Lewis, 1999; Munksgaard, & Simonsen, 1996).

##### 3.1.2. Behaviour in lairage

Pigs fasted at the farm tended to have a shorter latency to lie down than pigs fasted at the abattoir (35.7 vs.  $46.7 \pm 5.6$  min.;  $P = 0.06$ ). During the first 2 h of lairage, a greater ( $P < 0.01$ ) proportion of FARM pigs were lying (Fig. 1a), while PLANT pigs showed a higher ( $P < 0.05$ ) number of fights and a longer ( $P < 0.01$ ) total duration of fights for several periods (Fig. 1b, c). Moreover, fighting behaviour of PLANT pigs continued with the same intensity during the third hour of lairage (Fig. 1b, c). However, during the last 2 h of lairage, FARM pigs laid down less and fought more than PLANT pigs ( $P < 0.05$ ; Fig. 2a, b, c). The shorter latency to rest in FARM pigs on arrival at the pen may indicate their need to recover

from physical exhaustion caused by the additive effect of feed restriction, handling and transportation (Brandt, & Aaslyng, 2015). In this metabolic condition, in fact, pigs prefer settling down to rest rather than doing other activities (i.e. exploring the pen or fighting with the new pen mates). These results are in contrast with Brown, Knowles, Edwards, and Warriss (1999) who reported a greater activity (fighting and drinking) on arrival at the lairage pen in pigs fasted for longer time (18 vs. < 1 h). Fernandez, Meunier-Salaun, Ecolan, and Mormède (1995) also found a greater activity in fasted pigs (24 vs. 0 h) than fed ones that spent more time lying down. In contrast, in the present study, PLANT pigs started fighting intensively with unfamiliar pen-mates from the beginning of lairage and kept fighting constantly up to 3 h of lairage (limit of the observation time). Several studies already reported a more extended fighting rate in pigs kept in lairage for longer periods (overnight to > 24 h vs. 3 h) as a result of mixing (Guàrdia et al., 2009; Murray, & Jones, 1994; Warriss, Brown, Edwards, & Knowles, 1998). Because of the longer time spent in lairage resulting in more extended fighting PLANT pigs got fatigued over time and were mostly resting and recovering at the end of lairage time. Brown et al. (1999) also reported a significant reduction in fights between pigs after 1 h of lairage. Possibly, FARM pigs, which fought less during the first 2 h of lairage, recovered from the fatigue of transport and handling during this period in contrast to PLANT pigs, but this translated into greater fighting just before slaughter in FARM pigs.

### *3.2. Physiological Response*

*The variation of both lactate and CK levels in blood indicates the pig response to short and long-term physical stress, respectively, resulting from muscle fatigue or damage (Broom, 1995; Edwards et al., 2010). In this study, despite the greater activity observed for PLANT pigs in lairage, no difference in lactate and CK concentration was found in exsanguination*

blood between the two fasting groups ( $P = 0.76$  and  $P = 0.96$ , respectively; Table 3). The lack of effect of treatments on whole blood lactate may indicate that thanks to the adequate lairage conditions (sufficient space allowance to lie down) PLANT pigs had sufficient time to rest and recover from the vigorous initial fighting activity. It is known that if pigs are allowed to rest after physical exercise blood lactate returns to basal levels within 2 h (Edwards et al., 2010b). The greater fighting activity of FARM pigs during the last hours of lairage may have increased their blood lactate concentrations eventually levelling those of PLANT pigs. Overall, in both groups, the blood lactate levels at exsanguination were greater than 4 mM, which is the resting level of blood lactate reported for market-weight pigs (Edwards et al., 2011), indicating a general status of fatigue in all pigs in this study. Similarly, the lagged expression of fighting behaviour between the two groups during lairage may explain the lack of difference in blood CK level at slaughter, which achieves the maximum concentration peak at 6 h after the stress and does not return to basal levels until 48 h (Anderson, 2010 cited by Correa et al., 2013). Thus, it may appear that at the time of slaughter blood CK levels were increasing in FARM pigs and descending in PLANT pigs, which once again explains the lack of difference in the physiological response between the two fasting groups likely resulting from the confounding effect of the different time spent in lairage on these pigs' recovery rate.

#### 3.4. Carcass quality traits

When compared to FARM pigs, PLANT pigs neither differed in HCW (85.7 vs. 85.3 kg, SEM: 0.3;  $P = 0.28$ ) nor in carcass yield (69.3 % for both groups, SEM: 0.2;  $P = 0.99$ ) in this study. This result is not surprising as only fasting time in excess of 24 h preslaughter showed to result in liveweight losses, in terms of 100 g weight loss per hour, and consequently in carcass yield losses in pigs in previous studies (Faucitano et al., 2010). Finally, these results

*show that, no matter where a 24 h fasting time is applied through the preslaughter period, this duration has no impact on carcass weight and yield losses.*

### *3.5. Meat quality traits*

*In this study, the applied fasting time strategies only influenced the JCS score and drip loss percent of the LT muscle, with PLANT loins being darker (higher JCS scores;  $P = 0.02$ ) and drier (lower drip loss percent;  $P = 0.03$ ) than those from FARM pigs (Table 4). Indeed, when PLANT loins were classified in darker and drier or drier-only quality classes based on the JCS score ( $> 3.5$ ) and drip loss percent value ( $< 2\%$ ), a greater number of darker and drier or drier loins were found in PLANT carcasses than in FARM carcasses (17 vs. 12 and 81 vs. 72, respectively). However, this difference was only numerical ( $P = 0.43$  and  $P = 0.20$ , respectively). The variation in pork colour and drip loss, but not in the pHu values, may be indicative of a mild fatigue metabolic condition of PLANT pigs at slaughter that did not result in a sufficient depletion of muscle glycogen content to increase pHu, even in locomotor muscles, such as SM and AD muscles, that are more prone to glycogen exhaustion after feed restriction and physical exercise (Rocha, Dionne, Saucier, Nannoni, & Faucitano, 2015; Wittman, Ecolan, Levasseur, & Fernandez, 2004). According to Henckel, Karlsson, Jensen, Oksbjerg, and Petersen (2002), pHu values only increase when muscle glycogen content at slaughter is lower than  $53 \mu\text{mol/g}$ . The only mild fatigue status of these pigs at slaughter is also confirmed by the above-mentioned lack of variation in blood CK and lactate levels at exsanguination, which again may show the difference in recovery rate between pigs likely due to the different lairage time applied. Differently from the visual colour score, the Minolta  $L^*$  value, indicator of colour lightness, was not influenced by the fasting strategy (including lairage time) applied in this study. This result may be explained by the confounding effect of*

*the light reflectance of marbling fat, which was not evaluated in this study. High marbling fat scores have been, in fact, associated with higher  $L^*$  values due to fat colour reflectance, while they have no influence on visual colour scores (Jones, Tong, Campbell, & Dyckl, 1994; Van der Wal, Olsman, Garssen, & Engel, 1992).*

#### *4. Conclusions*

*Within the pork chain, it is generally agreed that a feed withdrawal time of 24 h preslaughter is the best compromise to obtain optimal carcass yield and pork safety. However, under split-marketing conditions the management of this practice, in terms of the timing of its application (at the farm or at the abattoir), is at the discretion of the producer or abattoir manager. Based on the results of this study, the choice of applying most of the fasting time at the abattoir rather than at the farm may not be appropriate as it prevents pigs from resting before slaughter and increases the risk for DFD-like pork production due to increased activity (fighting) in mixed groups of pigs that have to be kept in lairage for long time.*

#### *Acknowledgements*

*We appreciate the assistance of S. Horth, M.J. St.-Louis, C. Cavaleiro, S. Conte, A. de Castro, D. Morissette, M. Gil and R. Somavilla for the behaviour data collection and analysis and meat quality measurements at the abattoir. Special thanks go to S. Méthot for his support with the statistical analysis. The authors are grateful to Lucyporc Inc. for the financial support and the National Research Council of Brazil (CNPq) for the scholarship granted to F. Dalla Costa to conduct his graduate studies.*

#### *References*

- Aalhus, J. L., Murray, A. C., Jones, S. D. M., Tong, A. K. W., Raymond, D. P., & Weir, L. (1992). *Environmental conditions for swine during the marketing for slaughter: A national review*. Agriculture Canada, Research Branch, Ottawa, ON. Technical Bulletin 1992–6E.
- Arnone, M., & Dantzer, R. (1980). Does frustration induce aggression in pigs? *Applied Animal Ethology*, 6, 351–362.
- Averós, X., Knowles, T. G., Brown, S. N., Warriss, P. D., & Gonsálvez, L. F. (2008). Factors affecting the mortality of pigs being transported to slaughter. *Veterinary Record*, 163, 386–390.
- Bradshaw, R. H., Parrott, R. F., Goode, J. A., Lloyd, D. M., Rodway, R. G., & Broom, D. M. (1996). Behavioural and hormonal responses of pigs during transport: effect of mixing and duration of journey. *Animal Science*, 62, 547–554.
- Brandt, P., & Aaslyng, M. D. (2015). Welfare measurements of finishing pigs on the day of slaughter: a review. *Meat Science*, 103, 13–23.
- Brandt, P., Rousing, T., Herskin, M. S., & Aaslyng, M. D. (2013). Identification of post-mortem indicators of welfare of finishing pigs on the day of slaughter. *Livestock Science*, 157, 535–544.
- Broom, D. M. (1995). *Quantifying pig welfare during transport using physiological measures*. Proceedings EU Seminar “New Information on Welfare and Meat Quality of Pigs as Related to Handling, Transport and Lairage Conditions”. Mariensee, Germany (pp. 3–10).
- Brown, S. N., Knowles, T. G., Edwards, J. E., & Warriss, P. D. (1999). Relationship between food deprivation before transport and aggression in pigs held in lairage before slaughter. *Veterinary Record*, 145, 630–634.
- Correa, J. A., Méthot, S., & Faucitano, L. (2007). A modified meat juice container (EZ-DripLoss) procedure for a more reliable assessment of drip loss and related quality changes

*in pork meat. Journal of Muscle Foods, 18, 67–77.*

Correa, J. A., Gonyou, H. W., Torrey, S., Widowski, T. M., Crowe, T. G., LaForest, J.-P., & Faucitano, L. (2013). *Welfare and carcass and meat quality of pigs being transported for 2 hours using two vehicle types during two seasons of the year. Canadian Journal of Animal Science, 93, 43–55.*

D'Eath, R. B. (2002). *Individual aggressiveness measured in a resident-intruder test predicts the persistence of aggressive behaviour and weight gain of young pigs after mixing. Applied Animal Behaviour Science, 77, 267–283.*

Edwards, L. N., Grandin, T., Engle, T. E., Porter, S. P., Ritter, M. J., Sosnicki, A. A., & Anderson, D. B. (2010a). *Use of exsanguination blood lactate to assess the quality of pre-slaughter pig handling, Meat Science, 86, 384–390.*

Edwards, L. N., Grandin, T., Engle, T. E., Porter, S. P., Ritter, M. J., Sosnicki, A. A., Paradis, M. A., Correa, J. A., & Anderson, D. B. (2010b). *Relationship of blood lactate and meat quality in market hogs. Presentation at the Reciprocal Meat Conference, Lubbock, TX. [https://www.researchgate.net/publication/260983276\\_Relationship\\_of\\_blood\\_lactate\\_and\\_meat\\_quality\\_in\\_market\\_hogs](https://www.researchgate.net/publication/260983276_Relationship_of_blood_lactate_and_meat_quality_in_market_hogs) (accessed on January 25, 2016).*

Edwards, L. N., Engle, T. E., Grandin, T., Ritter, M. J., Sosnicki, A., Carlson, B. A., & Anderson, D. B. (2011). *The effects of distance traveled during loading, lairage time prior to slaughter, and distance traveled to the stunning area on blood lactate concentration of pigs in a commercial packing plant. Professional Animal Scientist, 27, 485-491.*

Eikelenboom, G., Hoving-Bolink, A. H., & Sybesma, W. (1991). *Effect of feed withdrawal before delivery on pork quality and carcass yield. Meat Science, 29, 25–30.*

Faucitano, L., Chevillon, P., & Ellis, M. (2010). *Effects of feed withdrawal prior to slaughter and nutrition on stomach weight, and carcass and meat quality in pigs. Livestock Science,*

127, 110–114.

Fernandez, X., Meunier-Salaun, M-C., Ecolan, P., & Mormède, P. (1995). Interactive effect of food deprivation and agonistic behaviour on blood parameters and muscle glycogen in pigs. *Physiology and Behaviour*, 58, 337–345

Guàrdia, M. D., Gispert, M., & Diestre, A. (1996). Mortality rates during transport and lairage in pigs for slaughter. *Meat Focus International*, 10, 362–366.

Guàrdia, M. D., Estany, J., Balash, S., Oliver, M. A., Gispert, M., & Diestre, A. (2005). Risk assessment of DFD condition due to pre-slaughter conditions and RYR1 gene in pigs. *Meat Science*, 70, 709–716.

Guàrdia, M. D., Estany, J., Balasch, S., Oliver, M. A., Gispert, M., & Diestre, A. (2009). Risk assessment of skin damage due to pre-slaughter conditions and RYR1 gene in pigs. *Meat Science*, 81, 745–751.

Henckel, P., Karlsson, A., Jensen, M. T., Oksbjerg, N., & Petersen, J. S. (2002). Metabolic conditions in porcine longissimus muscle immediately pre-slaughter and its influence on peri- and post mortem energy metabolism. *Meat Science*, 62, 145-155.

Jones, S. D. M., Tong, A. K. W., Campbell, C., & Dyck, R. (1994). The effects of fat thickness and degree of marbling on pork colour and structure. *Canadian Journal of Animal Science*, 74, 155–157.

Lewis, N. J. (1999). Frustration of goal-directed behaviour in swine. *Applied Animal Behaviour Science*, 64, 19–29.

Munksgaard, L. & Simonsen, H. B. (1996). Behavioural and pituitary adrenal-axis responses of dairy cows to social isolation and deprivation of lying down. *Journal of Animal Science*, 74, 769–778.

Murray, A. C., & Jones, S. D. M. (1994). The effect of mixing, feed restriction and genotype

*with respect to stress susceptibility on pork carcass and meat quality. Canadian Journal of Animal Science, 74, 587–594.*

*Nakai, H., F. Saito, T. Ikeda, S. Ando, & A. Komatsu (1975). Standards models of pork color. Bulletin of National Institute of Animal Industry 30,69–74.*

*National Farm Animal Care Council (2014). Codes of practice for the care and handling of pigs. <https://www.nfacc.ca/codes-of-practice/pigs> (accessed on January 25, 2016).*

*Peeters, E., Deprez, K., Beckers, F., De Baerdemaeker, J., Aubert, A. E., & Geers, R. (2008). Effect of driver and driving style on the stress responses of pigs during a short journey by trailer. Animal Welfare, 17, 189–196.*

*Pitts, A. D., Weary, D. M., Pajor, E. A., & Fraser, D. (2000). Mixing at young ages reduces fighting in unacquainted domestic pigs. Applied Animal Behaviour Science, 68, 191–197.*

*Rocha, L. M., Dionne, A., Saucier, L., Nannoni, E., & Faucitano L. (2015). Hand-held lactate analyzer as a tool for the real-time measurement of physical fatigue before slaughter and pork quality prediction. Animal, 9, 707–714.*

*Saucier, L., Bernier, D., Bergeron, R., Giguere, A., Methot, S., Faucitano, L. (2007). Effect of feed texture, meal frequency and pre-slaughter fasting on behaviour, stomach content and carcass microbial quality in pigs. Canadian Journal of Animal Science, 87, 479–487.*

*Schwartzkopf-Genswein, K. S., Faucitano, L., Dadgar, S., Shand, P., González, L. A., & Crowe, T. G. (2012). Road transport of cattle, swine and poultry in North America and its impact on animal welfare, carcass and meat quality: A review. Meat Science, 92, 227–243.*

*SAS (2002). Statistical analysis system. Release 9.1. SAS Institute Inc., Cary, NC, USA.*

*Stewart, G., Ritter, M. Culbertson, G. Mann, G., & Wofford, R. (2008). Effects of previous handling and feed withdrawal prior to loading on transport losses in market weight pigs. Proceedings of American Association of Swine Veterinarians. San Diego, CA (pp. 359–362).*

- Van der Wal, P. G., Olsman, W. J., Garssen, G. J., & Engel, B. (1992). Marbling, intramuscular fat and meat colour of Dutch pork. *Meat Science*, 32, 351–355.
- Viau, J., & Champagne, D. (1998). *La mise à jeun chez le porc avant l'abattage. Report of Centre de développement du Porc du Québec, Quebec City, Canada.*
- Warriss, P. D., Brown, S. N., Edwards, J. E., & Knowles, T. G. (1998). Effect of lairage time on levels of stress and meat quality in pigs. *Animal Science*, 66, 255–261.
- Weeding, C. M., Guise, H. J., & Penny, R. H. C. (1993). Factors influencing the welfare and carcass and meat quality of pigs: the use of water sprays in lairage. *Animal Production* 56, 393-397.
- Weschenfelder, A. V., Torrey, S. Devillers, N. Crowe, T., Bassols, A., Saco, Y., Piñeiro, M., Saucier, L., & Faucitano L. (2012). Effects of trailer design on animal welfare parameters and carcass and meat quality of three Pietrain crosses being transported over a long distance. *Journal of Animal Science*, 90, 3220–3231.

**Table 1.** Description of pig behaviors and interventions of the handler during loading  
(adapted from Weschenfelder et al., 2012)

	Description
<i>Pig behavior</i>	
Slip/fall	Leg of pig splits away from the other legs or pig falls down (at least 2 legs buckled under)
Overlap	Pig mounts another pig, with its 2 front legs on the back of the other pig
180° turn	Pig makes a 180° turn, ending with its rear extended in the direction of intended movement
Back up	Pig moves at least 2 steps rearward, opposite the direction of intended motion
Backward	Pig moves in the intended direction with its body oriented in the opposite direction
Underlap	Head of pig goes under the body of another pig
Vocalize	Pig vocalizes
Balk	Pig stops for more than 2 sec.
Squeeze	Pig is squeezed at the door, corridor, or the exit of the ramp (or at the trailer door, when unloading)
<i>Intervention of handler</i>	
Vocal sound	Handler uses his voice to encourage forward movement of 1 or a group of pigs
Rattle noise	Handler uses the paddle to produce noise (one time)
Physical contact	Handler uses his hands, paddle, or board (or whip at unloading) to push and encourage forward movement of 1 or a group of pigs

**Table 2.** Effects of feed withdrawal as mostly applied on-farm (FARM) or at slaughter plant (PLANT) on the frequency of pig behaviours and interventions of handlers at loading

Item	Feed withdrawal treatment		SEM	<i>P</i>
	FARM	PLANT		
<i>Pig behaviour</i>				
Slip/fall <sup>1</sup>	21.7	16.7		0.81
Overlap <sup>1</sup>	3.3	3.3		1.00
180° turn <sup>1</sup>	30.0	13.3		0.03
Back-up <sup>1</sup>	41.7	40.0		0.85
Backward <sup>1</sup>	20.0	6.7		0.03
Underlap <sup>1</sup>	10.0	11.7		0.77
Vocalize <sup>1</sup>	10.0	1.7		0.05
Squeeze <sup>1</sup>	21.7	10.0		0.08
Balk <sup>2</sup>	8.5	8.8	0.55	0.75
<i>Intervention of handler</i>				
Rattle noise <sup>1</sup>	6.7	8.3		0.73
Vocal sound <sup>2</sup>	14.32	15.18	1.04	0.56
Physical contact <sup>2</sup>	50.62	48.93	2.20	0.59

<sup>1</sup>Percentage of groups where the behaviour was observed

<sup>2</sup>Average frequency per group of 5 pigs

**Table 3.** Effects of feed withdrawal as mostly applied on-farm (FARM) or at slaughter plant (PLANT) on exsanguination blood parameters<sup>a</sup>

Parameters	Feed withdrawal treatment		<i>P</i>
	FARM (n = 105)	PLANT (n = 105)	
Lactate (mMol)	9.98 [8.88-11.28]	9.76 [8.64-11.03]	0.76
CK (IU/L)	3,172.65 [2,711-3,712]	3,187.18 [2,723-3,729]	0.96

<sup>a</sup>The values between brackets are lower and upper value of confidence interval

**Table 4.** Effects of feed withdrawal as applied on-farm or at the slaughter plant on meat quality traits as assessed in the *Longissimus thoracis* (LT), *Semimembranosus* (SM) and *Adductor* (AD) muscles

Variable	Feed withdrawal treatment		SEM	P
	FARM (n = 105)	PLANT (n = 105)		
<i>LT muscle</i>				
pH <sub>1</sub>	6.42	6.47	0.05	0.34
pHu	5.77	5.76	0.04	0.76
L*	52.67	52.08	0.58	0.19
a*	9.50	9.75	0.23	0.21
b*	5.99	5.91	0.16	0.61
JCS <sup>1</sup>	2.6	2.8	0.09	0.02
Drip loss, %	1.8	1.3	0.14	0.03
<i>SM muscle</i>				
pH <sub>1</sub>	6.39	6.43	0.08	0.48
pHu	5.84	5.80	0.05	0.10
<i>AD muscle</i>				
pHu	6.00	5.95	0.05	0.29

<sup>1</sup>JCS: Japanese colour scales (1 = pale to 6 = dark; Nakai et al., 1976).

## FIGURE CAPTIONS

Figure 1. Behaviour (back transformed least square means and confidence interval) at the beginning of lairage (after 20 min shower) of pigs being fasted for 24 h starting at the farm or the slaughter plant. a) percentage of pigs lying down, b) number of fights, c) total duration of fights. †P < 0.10, \*P < 0.05, \*\*P < 0.01 difference between treatments within lairage periods. For PLANT pigs only, black bars lacking a common superscript letter differ (P < 0.05)

Figure 2. Behaviour (back transformed least square means and confidence interval) at the end of lairage (from 120 min to the beginning of the last shower) of pigs being fasted for 24 h starting at the farm or the slaughter plant. a) percentage of pigs lying down, b) number of fights, c) total duration of fights. \*P < 0.05, \*\*\*P < 0.001 difference between treatments within lairage periods

Fig. 1





Fig. 2



### CHAPTER 3 – FINAL CONSIDERATIONS AND IMPLICATIONS

Feed withdrawal on-farm is recommended due to several benefits, such as feed saving, better pigs' welfare, reduction in transport losses and carcass contamination, and improvements in pork quality. Usually, a period of 24 h between the last meal and slaughter is recommended, which on-farm fasting of 12 hours should be respected for the guarantee of all benefits.

Reasons for fasting misapplication include concerns with slower growth rate of pigs remaining in the pen that were also withdrawn of feed and the lack of shipping rooms to transfer the heaviest pigs before loading time. However, fasting mainly done at slaughter plant than on-farm impairs welfare of pigs and results in lower pork quality.

Fights during preslaughter increase incidence of skin lesion, which is detrimental to pigs' welfare and produce higher incidence of DFD pork (GUÀRDIA et al., 2009). Unfortunately, because of the slaughter line conditions, it was not possible to measure skin lesions in this study. However, the results of darker and/or drier pork reflect the more fatigued conditions of pigs fasted at slaughter plant caused by fighting, which can be associated to psychological stress. Probably, if skin lesions were measured they would confirm these results. The lairage period should allow pigs to rest and recover from previous stress loading, transport and unloading. However, when pigs are not fasted before loading, they are prevented to lie down during lairage, which is demonstrated by the longer latency to rest. Indeed, the greater number of fights increases stress during preslaughter handling.

Lairage is also a reserve of pigs for feed the slaughter line and prevent from a lack of animals due to variations in the delivery schedule from farm to slaughter plant. When good conditions are provided, lairage leads to better pigs' welfare and pork quality (WARRISS, 2003). A lairage period of 1-3 h has been suggested as enough time to recover pigs from previous stresses during preslaughter handling and not causing problems of long-term food deprivation, skin damages and pork quality (DE SMET et al., 1996; WARRISS, 2003; YOUNG et al., 2009). However, the decision of increase lairage time in order to complete feed withdrawal of pigs have considerable impacts, not only for pigs' welfare and meat quality, but also for environment and

slaughter plant. When pigs are kept for longer times at lairage, it is necessary to have larger lairage area and greater number of pens at slaughter plant. The longer lairage time also increase the amount of fecal content which has to managed daily at the plant and can be a problem to process. Indeed, a greater amount of water is needed to clean the facilities and animals. Nowadays, the manner how companies treat natural resources are not only a legislation issue, but also an interest of costumers that impairs pork production. Therefore, problems with environment should be avoided in order to keep the production and a good image of pork products.

A possible alternative to apply the correct feed withdrawal on-farm avoiding loss of weight gain and the need of shipping rooms is based on the pigs' behaviour according to feed intake pattern. It allows keeping the weight selected pigs in their own finishing pen without closing the feeders. Finishing pigs have diurnal feeding behaviour (FAUCITANO; CHEVILLON, 2010). Some studies (HYUN et al., 1997; HYUN; ELLIS, 2001, 2002) reported that consume 70-75% reaches it daily feed intake between 06 a.m. and 06 p.m. hours for growing and finishing pigs (FAUCITANO; CHEVILLON, 2010). As well as the duration, it has implications for the starting time of feed withdrawal period (FAUCITANO; CHEVILLON, 2010). So, pigs that started feed withdrawal at 06 a.m. compared to 06 p.m. would have different stomach weight. Lewis and McGlone (2008), based on the feed intake behaviour, suggested that the combination of load pigs between 04 a.m. and 08 a.m., and transport for 4 h or more could be used to reduce the incidence of full stomachs at slaughter. Another option is anticipate the last meal in the finishing pen and ship pigs early in the morning.

The lack of effects found on transport losses was removed from the manuscript due to the inadequacy of the experimental design to get representative data for such measure in this study. Power tests performed in recent transportation trials (Lewis et al., 2010. *Animal Production Science*, 50, 1–9) showed, in fact, that a minimum of 250 transports (or loads) are needed to yield significant results in this type of study. The rate of transport losses was only collected as complementary information in this study.

Therefore, results from this study showed that on-farm feed withdrawal combined to proper lairage time is recommended to improve pigs' welfare and pork

quality. Indeed, for a correct application of this practice, communication between producer and slaughter plant is necessary to plan on-farm feed withdrawal before loading and total feed withdrawal period.

## REFERENCES

- DE SMET, S. M.; PAUWELS, H.; DE BIE, S.; DEMEYER, D. I.; CALLEWIER, J.; EECKHOUT, W. Effect of halothane genotype, breed, feed withdrawal, and lairage on pork quality of Belgian slaughter pigs. **Journal of Animal Science**, v. 74, p. 1854-1863, 1996.
- FAUCITANO, L.; CHEVILLON, P.; ELLIS, M. Effects of feed withdrawal prior to slaughter and nutrition on stomach weight, and carcass and meat quality in pigs. **Livestock Science**, v. 127, n. 2, p. 110-114, 2010.
- GUÀRDIA, M. D.; ESTANY, J.; BALASCH, S.; OLIVER, M. A.; GISPERT, M.; DIESTRE, A. Risk assessment of skin damage due to pre-slaughter conditions and RYR1 gene in pigs. **Meat Science**, v. 81, n. 4, p. 745-751, 2009.
- HYUN, Y.; ELLIS, M. Effect of group size and feeder type on growth performance and feeding patterns in growing pigs. **Journal of animal science**, v. 79, n. 4, p. 803-810, 2001.
- HYUN, Y.; ELLIS, M. Effect of group size and feeder type on growth performance and feeding patterns in finishing pigs. **Journal of Animal Science**, v. 80, n. 3, p. 568-574, 2002.
- HYUN, Y.; ELLIS, M.; McKEITH, F. K.; WILSON, E. R. Feed intake pattern of group-housed growing-finishing pigs monitored using a computerized feed intake recording system. **Journal of Animal Science**, v. 75, n. 6, p. 1443-1451, 1997.
- LEWIS, C. R. G.; MCGLONE, J. J. Modelling feeding behaviour, rate of feed passage and daily feeding cycles, as possible causes of fatigued pigs. **Animals**, v. 2, p. 600-605. 2008.
- WARRISS, P. D. Optimal lairage times and conditions for slaughter pigs: a review. **The Veterinary Record**, v. 153, n. 6, p. 170-176, 2003.
- YOUNG, J. F.; BERTRAM, H. C.; OKSBJERG, N. Rest before slaughter ameliorates pre-slaughter stress-induced increased drip loss but not stress-induced increase in the toughness of pork. **Meat Science**, v. 83, n. 4, p. 634-641, 2009.