



Domain wall description of superconductivity



F.A. Brito^{a,*}, M.L.F. Freire^b, J.C. Mota-Silva^{a,c}

^a Departamento de Física, Universidade Federal de Campina Grande, Caixa Postal 10071, 58109-970 Campina Grande, Paraíba, Brazil

^b Departamento de Física, Universidade Estadual da Paraíba, 58109-753 Campina Grande, Paraíba, Brazil

^c Departamento de Física, Universidade Federal da Paraíba, Caixa Postal 5008, 58051-970 João Pessoa, Paraíba, Brazil

ARTICLE INFO

Article history:

Received 21 November 2013

Received in revised form 21 November 2013

Accepted 26 November 2013

Available online 3 December 2013

Editor: M. Cvetič

ABSTRACT

In the present work we shall address the issue of electrical conductivity in superconductors in the perspective of superconducting domain wall solutions in the realm of field theory. We take our set up made out of a dynamical complex scalar field coupled to gauge field to be responsible for superconductivity and an extra scalar real field that plays the role of superconducting domain walls. The temperature of the system is interpreted through the fact that the soliton following accelerating orbits is a Rindler observer experiencing a thermal bath.

© 2013 The Authors. Published by Elsevier B.V. Open access under CC BY license.

1. Introduction

In this Letter we present an alternative description of superconductivity in field theory. We take advantage of the rich and well established features involving classical soliton solutions, mainly those formulated in multi-scalar fields [1–4]. The reason for using such solutions in our purpose is twofold. Firstly, because they can develop internal structures such as condensates which are fundamental for studying superconductivity. Alternatively, this opens the possibility of studying *superconducting solitons*. The first example of such objects were the superconducting strings [1,2] – other developments in domain walls with internal structures were also considered in [3,4]. Secondly, because these solutions can follow non-trivial orbits in the field space [4]. They mostly force the solitons to move into accelerated trajectories. As such, we can identify these solitons as non-relativistic Rindler observers experiencing a thermal bath. As we shall see, this will be fundamental to introduce temperature in the system in a very natural way and identify several important quantities such as the condensate and resistivity as a function of the temperature. We believe this alternative can open a new window for investigating superconductivity in field theory through superconducting solitons since there exist many types of soliton solutions in many well established field theories such a way superconducting solitons can also be identified. This new perspective may complement and shed some new light on earlier studies of high- T_c superconductivity in field theory focused on ‘particle excitations’ only [5].

In the present study, we take our set up made out of a dynamical complex scalar field coupled to the abelian gauge field to be responsible for superconductivity and an extra real scalar field that plays the role of superconducting domain walls. They are domain wall backgrounds that develop a condensate in their core. Furthermore, the domain wall model is a good approximation for superconductors presenting, for instance, a layer-type perovskite-like structure [6,7]. The quantum field theory can explain some effects of superconductivity. The results can be obtained with an appropriate classical regime of a quantum field theory inspired by the Ginzburg–Landau (GL) theory [8]. Though initially proposed as a phenomenological theory, the GL theory can be shown to be a limiting case of a microscopic theory [9] such as the Bardeen–Cooper–Schrieffer (BCS) [10] theory of superconductivity. The domain of validity of the GL theory is shown to be restricted to temperatures sufficiently near the critical temperature and to spatially slow varying fields [7].

The Letter is organized as follows. In Section 2 we address the issue of type II domain wall solutions in field theory coupled to gauge field. We consider such solutions as background fields to solve a Schrodinger-like equation for the electromagnetic field. In Section 3 we compute the condensate at finite temperature. The temperature of the system is interpreted through the fact that the soliton following accelerating orbits is a Rindler observer experiencing a thermal bath. In Section 4 we calculate the optical conductivity in terms of the frequency and temperature. In the limit of very low frequencies and temperatures the superconductor develops infinite DC conductivity, as expected. Furthermore, in this regime we can easily read off the binding energy Δ of a Cooper pair from the real part of the optical conductivity. At some temperature around the critical temperature the AC resistivity develops

* Corresponding author.

an effect commonly seen in high- T_c superconductors [6]. Finally, in Section 5 we make our final comments.

2. Superconducting type II domain walls solution

To obtain superconducting domain walls is needed a complex scalar field with charge q that must couple to a real scalar field that produces the domain walls. By introducing a coupling between the complex scalar field and the electromagnetic field, the former develops a *condensate* inside the domain wall which becomes superconducting and develops almost all the properties of a superconducting material. The superconducting domain wall developing a condensate can be generated by the following Lagrangian with the $Z_2 \times U(1)$ symmetry:

$$\mathcal{L} = \frac{1}{2} \partial_\mu \phi \partial^\mu \phi + (\partial^\mu \chi + iq A^\mu \chi) (\partial_\mu \chi^* - iq A_\mu \chi^*) - V(\phi, \chi, \chi^*) - \frac{1}{4} F_{\mu\nu} F^{\mu\nu}, \quad (1)$$

where $\mu, \nu = 0, 1, 2, \dots, d$ are bulk indices for arbitrary $(d-2)$ -dimensional domain walls. We shall focus on two-dimensional domain walls with bulk indices running as $\mu, \nu = t, x, y, r$. The potential $V(\phi, \chi, \chi^*)$ is chosen appropriately so that the domain wall becomes a superconductor

$$V(\phi, \chi, \chi^*) = \frac{1}{2} \lambda^2 (\phi^2 - a^2)^2 + \lambda \mu (\phi^2 - a^2) |\chi|^2 + \frac{1}{2} \mu^2 |\chi|^4 + \mu^2 \phi^2 |\chi|^2. \quad (2)$$

The real scalar field ϕ develops Z_2 symmetry and is responsible to form the domain wall, whereas the charged scalar field χ develops a condensate inside the domain wall and is also responsible to produce type II domain walls. The equations of motion for the real and complex scalar fields coupled to electromagnetic field are given by

$$\square \phi + \frac{\partial V}{\partial \phi} = 0, \quad (3)$$

$$\square \chi + \frac{\partial V}{\partial \chi^*} + 2iq A_\mu \partial^\mu \chi - q^2 A_\mu A^\mu \chi = 0, \quad \text{c.c.}, \quad (4)$$

$$\square A_\mu + iq(\chi^* \partial_\mu \chi - \chi \partial_\mu \chi^*) + 2q^2 A_\mu |\chi|^2 = 0. \quad (5)$$

For $A_\mu = 0$ the scalar real sector produces domain wall solutions whose kink profiles are the following well-known BPS *static* solutions obtained in terms of first order formalism and a specific superpotential [4]. The type I solution

$$\phi(r) = -a \tanh(\lambda r),$$

$$\chi = 0, \quad (6)$$

and the type II solution

$$\phi(r) = -a \tanh(2\mu ar),$$

$$\chi(r) = \pm a \sqrt{\frac{\lambda}{\mu} - 2} \operatorname{sech}(2\mu ar), \quad (7)$$

where r is the spatial coordinate transverse to the domain walls. These solutions correspond to straight and elliptic orbit, respectively [4] – see Fig. 1. They have the same Bogomol'nyi energy. Note that for sufficiently large λ/μ the elliptic orbit

$$\phi^2 + \left(\frac{\lambda}{\mu} - 2\right)^{-1} \chi^2 = a^2, \quad (8)$$




Fig. 1. Elliptic orbits followed by the type II kink. The temperature becomes high from the top ($T = 0$) to bottom ($T = T_c$) orbits.

passes through the ‘supersymmetric vacuum’ $\phi = 0$ and $\chi = \pm a \sqrt{\lambda/\mu}$ – global minima of the scalar potential written in terms of first derivatives of superpotential [4]. As we shall discuss later, this will correspond to zero temperature, which agrees with the formation of a condensate inside the superconducting domain wall.

Now considering these solutions as background fields we solve the equation of motion for electromagnetic field (5) on these backgrounds. By introducing $A_\mu(t, r) = A_\mu(r) e^{-i\omega t}$ and $\chi(t, r) = \chi(r) e^{-i\theta t}$ we obtain the Schrodinger-like equation for A_x (or A_y) as follows

$$-A_x'' + \frac{1}{4} \ell^2 \operatorname{sech}^2(\alpha r) A_x = \omega^2 A_x, \quad (9)$$

where $\ell = 2\sqrt{2}qa\sqrt{\frac{\lambda}{\mu} - 2}$ and $\alpha = 2\mu a$. This is a well-known Schrodinger problem with a sech-type barrier potential whose solution is given by

$$A_x(\omega, \alpha, \ell, r) = (\operatorname{sech}(\alpha r))^{-\frac{i\omega}{\alpha}} {}_2F_1\left[a1, a2; a3; \frac{1}{2}(1 - \tanh(\alpha r))\right], \quad (10)$$

where ${}_2F_1$ is a hypergeometric function with the parameters defined as

$$\begin{aligned} a1 &= \frac{1}{2} \frac{-2i\omega + \alpha + \sqrt{-\ell^2 + \alpha^2}}{\alpha}, \\ a2 &= -\frac{1}{2} \frac{2i\omega - \alpha + \sqrt{-\ell^2 + \alpha^2}}{\alpha}, \\ a3 &= -\frac{i\omega - \alpha}{\alpha}. \end{aligned} \quad (11)$$

3. The condensate at finite temperature

The usual manner to introduce finite temperature in quantum field theory is through path integral with Euclidean time whose period is related with the temperature of the system. In this sense the path integral turns out to be the partition function from which one can obtain Bose–Einstein and Fermi–Dirac statistics for bosonic and fermionic fields, respectively. Furthermore, in curved spacetime or equivalently for accelerated observers such statistics tell us that the temperature is related to surface gravity or acceleration – see [11] for further details. In our system although the superconducting domain wall is living in a flat spacetime its related soliton (i.e., its BPS kink profile (7)) is accelerated in the field space as a non-relativistic Rindler observer – see below. As we have previously mentioned we shall take advantage of this fact to identify




Fig. 2. Hyperbolic worldlines (orbits) followed by the type II solitons in 'Lorentzian' signature of the (ϕ, χ) -space.

the temperature of the system as in the following. We shall first attempt to convince the reader that the parameter α can be indeed related to the temperature of the system. The orbits of Fig. 1 followed by the BPS solutions into the (ϕ, χ) -plane make the type II kink solutions to experiment constant *accelerations*. One can think of such soliton solutions following accelerating trajectories (or orbits). We can also see these trajectories in a 'Lorentzian' signature in (ϕ, χ) -plane by making the change $\phi \rightarrow i\phi$ – see Fig. 2 – and $r \rightarrow i\tau$ in the BPS solutions (7).

The soliton in this sense is a Rindler observer experiencing a thermal bath of bosonic and fermionic modes that are distributed, respectively, according to Bose–Einstein and Fermi–Dirac statistics (this is the Unruh effect [11])

$$\langle n_\Omega \rangle = \frac{1}{\exp\left(\frac{2\pi\Omega}{a}\right) \mp 1}, \quad (12)$$

where the constant acceleration a is promptly identified with the Unruh temperature $T = \frac{a}{2\pi}$ and Ω are the frequencies of the *thermal radiation*. In the following we show that the solution (7) indeed describes an accelerated trajectory. Firstly, we redefine the fields ϕ , χ in terms of coordinates of space and time as follows $\phi = \alpha a t(\tau)$ and $\chi = \alpha a (\frac{\lambda}{\mu} - 2)^{1/2} z(\tau)$ such that

$$\begin{aligned} t(\tau) &= \frac{1}{\alpha} \tanh \alpha \tau, \\ z(\tau) &= \frac{1}{\alpha} \operatorname{sech} \alpha \tau, \end{aligned} \quad (13)$$

where $\tau \equiv x$ is identified with (Euclidean) proper time. Now using the definition of the acceleration $a^\mu = \frac{d^2 x(\tau)^\mu}{d\tau^2}$ we find that

$$\begin{aligned} a \equiv \sqrt{a_\mu a^\mu} &= \alpha - \frac{1}{2} \alpha (\alpha \tau)^2 + \dots \\ &\simeq \alpha. \end{aligned} \quad (14)$$

In the last step we have assumed the regime of very slow velocities $v = a\tau \ll 1$ (non-relativistic Rindler observer). Thus, using the Unruh temperature we find $\alpha \simeq 2\pi T$ that we shall simply assume throughout the Letter the useful correspondence

$$\alpha \equiv T. \quad (15)$$

On the other hand, let us now compute the analog of acceleration in the (ϕ, χ) -plane. Actually we will be indeed calculating the inverse of acceleration since the coordinates (ϕ, χ) have dimension of energy instead of length as in the usual sense. We can now define the acceleration of the system as

$$a(r_0)^{-1} = \frac{d^2 \chi}{d\phi^2} \Big|_{r_0}, \quad (16)$$

where r_0 is some point on the bulk. The r.h.s. of Eq. (16) can be written in terms of the superpotential [4] and type II kink solutions in the form

$$\begin{aligned} \frac{d^2 \chi}{d\phi^2} \Big|_{r_0} &= \frac{d}{d\phi} \left(\frac{d\chi}{d\phi} \right) \Big|_{r_0} = \frac{d}{d\phi} \left(\frac{W_\chi}{W_\phi} \right) \Big|_{r_0} = \frac{W_{\phi\chi}}{W_\phi} - \frac{W_{\phi\phi} W_\chi}{W_\phi^2} \Big|_{r_0} \\ &= -\frac{\sqrt{\frac{\lambda}{\mu} - 2}}{a \operatorname{sech}(2\mu a r_0)} \left(1 + \frac{\lambda}{\mu} \tanh^2(2\mu a r_0) \right), \end{aligned} \quad (17)$$

where the minus sign simply reflects the concavity of the orbits. The temperature can be now defined in the general form

$$\beta = \left| \frac{d^2 \chi}{d\phi^2} \right|_{r_0}, \quad \beta = \frac{1}{T}. \quad (18)$$

In order to identify the temperature in terms of the parameters of the theory let us investigate the regime near the core of the domain wall (i.e., in the supersymmetric vacuum $\phi = 0$ and $\chi = \pm a\sqrt{\lambda/\mu}$), that is $r_0 \approx 0$ and $\lambda/\mu \gg 1$. This means that the temperature is now defined through the following equation

$$\frac{1}{T} = \frac{\sqrt{\frac{\lambda}{\mu} - 2}}{a} \approx \frac{\sqrt{\frac{\lambda}{\mu}}}{a}. \quad (19)$$

Eqs. (15) and (19) together with the fact that $\lambda a \equiv T_c$ (that by construction we assume to be the regime of critical temperature, since for $T < T_c$ we have shown that $2\mu a = T$), are easily satisfied by the dimensionless (λ, μ) and dimensionful (a) parameters, up to numeric irrelevant pre-factors, as follows

$$\lambda \sim \frac{T_c^{1/2}}{T^{1/2}}, \quad \mu \sim \frac{1}{2} \frac{T^{1/2}}{T_c^{1/2}}, \quad a \sim T_c^{1/2} T^{1/2}. \quad (20)$$

Let us now consider a region out of the core, that is $2\mu a r_0 \ll 1$ and $\lambda/\mu \gg 1$. Now substituting these assumptions into (17) we obtain the temperature

$$T = \frac{a(\lambda/\mu)^{-3/2}}{(2\mu a r_0)^2}. \quad (21)$$

Now the temperature has a dependence with four parameters. However, we shall keep using the correspondence $\alpha = 2\mu a \equiv T$ and $\lambda a \equiv T_c$ as in the previous analysis. Thus, substituting again the parameters (20) into the temperature (21) we readily find a relation between the critical temperature T_c and the scale r_0 given by

$$T = \frac{1}{2^{3/2} T_c r_0^2}. \quad (22)$$

Since by definition the temperature T does not depend on r_0 then

$$T_c r_0^2 = \text{const.} \quad (23)$$

We are still tempted to write this formula in terms of atomic mass A in the bulk since the lattice parameter can be well approximated by $r_0 \simeq A^{1/3}$ fm. Thus, we arrive to the *isotopic mass formula* [7]




Fig. 3. The effective condensate as a function of temperature for charges $q = 1, 2$, and 3 from bottom to top.

$$T_c A^{2/3} = \text{const.} \quad (24)$$

The condensate can be easily isolated by power expanding the scalar solution $\chi(r)$ in Eq. (7) around the core of the type II domain walls at $r \approx 0$ as

$$\chi(r) = m - \frac{1}{2} m \alpha^2 r^2 + \dots, \quad (25)$$

with the condensate given by $\langle \chi \rangle \simeq m$. We have written the original type II solution as $\phi(r) = -a \tanh \alpha r$, $\chi(r) = m \text{sech } \alpha r$. To find the explicit dependence of the condensate with the temperature we recall the original form of the parameters m, α , that is

$$m = a \sqrt{\frac{\lambda}{\mu} - 2}, \quad \alpha = 2\mu a. \quad (26)$$

Now substituting (20) into (26) allows us to obtain

$$m = \sqrt{2} T_c \sqrt{1 - \frac{T}{T_c}}, \quad (27)$$

which implies that the condensate has precisely the desired form $\langle \chi \rangle \simeq m = \sqrt{2} T_c \sqrt{1 - T/T_c}$. We also note that from the equation of motion of electromagnetic field A_x the *effective condensate* ‘seen’ by the electromagnetic field is given in terms of the charge q , i.e., $\langle \chi \rangle_{eff} \simeq \ell = 2\sqrt{2} q m$ or simply $\langle \chi \rangle_{eff} \simeq 4q T_c \sqrt{1 - T/T_c}$, where ℓ is defined just below Eq. (9) – see Fig. 3 for the explicit behavior of the condensate $\langle \chi \rangle_{eff}$ with the temperature.

4. Conductivity

As well known, from the Ohm’s law we can readily obtain the conductivity along a direction, say x -direction along the domain walls, in the form

$$\sigma_x(x, y) = \frac{J_x}{E_x} = \frac{A'_x(0)}{i\omega A_x(0)}, \quad (28)$$

where in the last step we used $E_x = -\partial_t A_x = i\omega A_x$, with $A_x(t, r) = A_x(r)e^{-i\omega t}$ and defined the current as $J_x = A'_x(0)$. This can be justified by using the boundary conditions for the electromagnetic field on an interface at $r = 0$ (or in other position as we shall consider later) that corresponds to a plane along the superconducting domain wall. More specifically, the boundary conditions for the magnetic field at an interface is

$$\hat{n} \times \vec{B} = \vec{J}, \quad \text{at } r = 0, \quad (29)$$

where \hat{n} is a normal vector to the surface of the domain wall and \vec{J} is a surface current. For $\hat{n} = (0, 0, 1)$ and $\vec{A} = (A_x, A_y, 0)$ the boundary condition (29) simply becomes




Fig. 4. The real part of the conductivity as a function of the frequency normalized by the effective condensate. We use the charges $q = 8, 20$, and 32 from top to bottom; $\delta = 0.01$ and $\ell \simeq \langle \chi \rangle_{eff} = 4$.

$$-\partial_r A_x(r) = J_x, \quad \text{at } r = 0, \quad (30)$$

which is nothing but our anticipated assumption $J_x = A'_x(0)$.

Now using the solution (10) for the electromagnetic field around a generic point $r \approx \delta$ (same as r_0) we are able to write the explicit form of the conductivity $\sigma_x = \sigma_y \equiv \sigma$ as follows

$$\begin{aligned} \sigma(\omega, \alpha, \ell, \delta) &= \frac{\frac{1}{8}(4\omega^2 + 4i\omega\alpha - \ell^2) {}_2F_1[b1, b2; b3; \frac{1}{2}(1 - \tanh(\alpha\delta))] \text{sech}^2(\alpha\delta)}{\omega(i\omega - \alpha) {}_2F_1[a1, a2; a3; \frac{1}{2}(1 - \tanh(\alpha\delta))]} + \tanh(\alpha\delta), \end{aligned} \quad (31)$$

with the parameters $b1, b2$ and $b3$ defined as

$$\begin{aligned} b1 &= -\frac{1}{2} \frac{2i\omega - 3\alpha + \sqrt{-\ell^2 + \alpha^2}}{\alpha}, \\ b2 &= \frac{1}{2} \frac{-2i\omega + 3\alpha + \sqrt{-\ell^2 + \alpha^2}}{\alpha}, \\ b3 &= -\frac{i\omega - 2\alpha}{\alpha}. \end{aligned} \quad (32)$$

Recall we have previously defined the temperature $\alpha \equiv T$ and the condensate $m \simeq \langle \chi \rangle$. We now consider the conductivity normalized by the ‘effective condensate’ $\ell \rightarrow q\ell$, such that we define $\alpha = q^{-1}q\ell$ and $\omega = \omega_r q\ell$ into σ . We can still write $\frac{\alpha}{q\langle \chi \rangle_{eff}} = q^{-1}$ and $\frac{\omega}{q\langle \chi \rangle_{eff}} = \omega_r$ (reduced frequency). Finally we substitute all over this into (31)–(32). The results have shown that for $\delta \approx 0$ the optical conductivity – see Fig. 4 – is essentially the same as the one computed at $r = 0$, i.e., at the core of the domain wall. On the other hand, as we shall see later, the conductivity (or AC resistivity) as a function of the temperature is more sensitive to the values of δ . In the following we focus on further characteristics of the optical conductivity by simply assuming $\alpha\delta = 0$.

As $\omega \rightarrow 0$ and $T \rightarrow 0$ the conductivity (31) approaches a delta function $\delta(\omega)$. This is because for $T \rightarrow 0$, we have $\ell \sim T_c$. Thus, in this limit, $\alpha^2 \ll \ell^2$ and $\omega^2 \ll \ell^2$ implies that the real and imaginary parts of the conductivity, up to a prefactor $\sim 2\alpha/\ell$ from the hypergeometric functions, can be written as

$$\begin{aligned} \text{Re } \sigma(\omega) &\propto \frac{(\ell/\alpha)^2}{(\omega/\alpha)^2 + 1} \rightarrow \delta(\omega), \\ \text{Im } \sigma(\omega) &\propto \frac{(\ell/\alpha)^2}{(\omega/\alpha)^3 + \omega/\alpha} \rightarrow \frac{\ell^2}{\alpha} \frac{1}{\omega}. \end{aligned} \quad (33)$$

Note that the distribution in real part of (33) tends to a delta function, whereas the imaginary part presents a pole. This in accord with the Kramers–Kronig relations and Drude model of conductor in the limit of the relaxation time due to scattering $\tau \rightarrow \infty$ (superconductor). We conclude from such discussion that in the limit $\omega \rightarrow 0$ at $T \rightarrow 0$ our model presents an infinite DC conductivity as expected for a superconductor.

Now let us consider the conductivity as a function of the temperature. Repeating the previous analysis for $\alpha\delta \rightarrow \infty$ the argument in the hypergeometric functions goes to zero as $e^{-2\alpha\delta}$. In this regime the ratio of the hypergeometric functions in the conductivity formula can be well approximated by a series of a few terms. By keeping only leading and next-to-leading terms we find

$$\text{Re } \sigma(\omega, \alpha) \propto \delta(\omega) \left(1 - \frac{1}{8} \frac{\ell^2}{\alpha^2} e^{-2\alpha\delta} + \dots \right) \simeq \delta(\omega) e^{-\frac{1}{8}(\frac{\Delta}{\alpha})^2}, \quad (34)$$

where

$$\Delta = \ell e^{-\alpha\delta}, \quad (35)$$

precisely defines the binding energy of a Cooper pair as long as we identify $\ell = 2\omega_D$ as the Debye temperature and $\delta\alpha = 1/VN_F$, being $V > 0$ the binding potential and N_F the density of orbitals with Fermi's energy. Note that the limit $\delta\alpha \rightarrow \infty$ corresponds to $VN_F \rightarrow 0$ that is the limit of weak coupling which is in accord with the BCS theory. On the other hand, the limit $\delta\alpha \rightarrow 0$ corresponds to $VN_F \rightarrow \infty$ that is the limit of strong coupling.

In order to go further into our analysis we extend the power expansion of the condensate in (25) to the expansion around a plane $r \approx \delta$, parallel to the domain wall near its center given by

$$\chi(r) = m \text{sech}(\alpha\delta) - m \text{sech}(\alpha\delta) \tanh(\alpha\delta) \alpha(r - \delta) + \dots, \quad (36)$$

which allows us to redefine the usual condensate previously studied in the form $\langle \chi \rangle \simeq m \text{sech}(\alpha\delta) = \sqrt{2}T_c \sqrt{1 - T/T_c} \text{sech}(\alpha\delta)$ and the effective condensate as $\langle \chi \rangle_{\text{eff}} \simeq \ell \text{sech}(\alpha\delta) = 2\sqrt{2}qm \text{sech}(\alpha\delta)$ or simply $\langle \chi \rangle_{\text{eff}} \simeq 4qT_c \sqrt{1 - T/T_c} \text{sech}(\alpha\delta)$. In the regime $\alpha\delta \rightarrow \infty$, particularly, we write the effective condensate given by

$$\langle \chi \rangle_{\text{eff}} \simeq 2\ell e^{-\alpha\delta}. \quad (37)$$

Eqs. (35) and (37) now allow us to write the important relation

$$\frac{2\Delta}{T_c} = \frac{\langle \chi \rangle_{\text{eff}}}{T_c}. \quad (38)$$

Recall the examples in Fig. 3 for the effective condensate for three distinct charges in the regime $\alpha\delta \rightarrow 0$ previously considered. Using Eq. (38) we identify the relations between binding energy and the critical temperature given by $2\Delta \simeq 4T_c$, $2\Delta \simeq 8T_c$ and $2\Delta \simeq 12T_c$.

This seems to point out a behavior of high- T_c superconductors. For the sake of comparison we know that BCS superconductors have a typical relation $2\Delta \simeq 3.5T_c$, whereas the high- T_c superconductors normally enjoy the relation $2\Delta \simeq 5T_c$ to $2\Delta \simeq 8T_c$.

In the following we present the plot that describes the behavior of the real part of the low frequency AC resistivity $\rho = 1/\sigma$ as a function of the temperature — see Fig. 5. Notice that for the resistivity sufficiently above the critical temperature T_c decreases almost linearly with the temperature. Moreover, as the system approaches the critical temperature, the resistivity tends to locally increases, but decreases very quickly below the critical temperature until achieve the resistivity very close to zero. This should be




Fig. 5. The real part of the AC resistivity at low frequencies as a function of temperature. We use $\delta = 0.40, 0.45$, and 0.55 , from bottom to top; $T_c = 3$, $\omega = 0.8$ and $q = 1$.

compared with the resistivity versus temperature for three high- T_c superconductor samples of La–Ba–Cu–O with $T_c = 35$ K by Bednorz and Müller [6]. This result confirms, at least qualitatively, that our superconducting domain walls model agrees with some properties of cuprates which are well-known layer-type perovskite-like structures. We still note that for values of δ larger than 0.40 the system tends to reduce its critical temperature — see Fig. 5, the plots for $\delta = 0.45$ and $\delta = 0.55$.

Finally, as a last comment we can even improve our above discussions by considering more general orbits. The original idea of trial orbit method [4] enables us extending the aforementioned elliptic orbit to general orbits

$$\phi^2 + \left(\frac{\lambda}{\mu} - 2 \right)^{-1} \chi^n = a^2. \quad (39)$$

Although much harder to deal with, this system will produce more general type I and type II solutions in such a way that the condensate may have a more general power

$$\langle \chi \rangle \simeq T_c \left(1 - \frac{T}{T_c} \right)^{1/n}. \quad (40)$$

This extension may reveal an even more realistic analysis for considering domain wall description of superconductivity.

5. Conclusions

In this Letter, by considering domain wall description of superconductivity, we have identified a relationship between the binding energy of Cooper pairs and the effective condensate depending on the temperature and the electrical charge. For charges large enough we get a typical ratio of high- T_c superconductors. We calculate the optical conductivity and show that in the regime of low temperatures and frequencies we get an infinite DC conductivity. We conclude that the low frequency AC resistivity as a function of temperature is similar to what happens in high- T_c superconductors. The critical temperature tends to be reduced when we

move far from the condensate via the deviation position δ . As future prospects, we intend to attack the problem by investigating other quantities such as the London penetration depth and upper critical field H_{c2} as a function of temperature and effects of anisotropy within the domain walls.

Acknowledgements

We would like to thank CNPq, CAPES, PNPd/PROCAD-CAPES for partial financial support.

References

- [1] E. Witten, Nucl. Phys. B 249 (1985) 557.
- [2] A. Vilenkin, E.P.S. Shellard, Cosmic Strings and Other Topological Defects, Cambridge University Press, 1994.
- [3] R. MacKenzie, Nucl. Phys. B 303 (1988) 149;
J.R. Morris, Int. J. Mod. Phys. A 13 (1998) 1115, arXiv:hep-ph/9707519;
F.A. Brito, D. Bazeia, Phys. Rev. D 56 (1997) 7869, arXiv:hep-th/9706139;
J.D. Edelstein, M.L. Trobo, F.A. Brito, D. Bazeia, Phys. Rev. D 57 (1998) 7561, arXiv:hep-th/9707016.
- [4] R. Rajaraman, Solitons and Instantons: An Introduction to Solitons in Quantum Field Theory, North-Holland, Amsterdam, 1982;
D. Bazeia, M.J. dos Santos, R.F. Ribeiro, Phys. Lett. A 208 (1995) 84, arXiv:hep-th/0311265;
D. Bazeia, H. Boschi-Filho, F.A. Brito, J. High Energy Phys. 9904 (1999) 028, arXiv:hep-th/9811084;
D. Bazeia, F.A. Brito, Phys. Rev. D 61 (2000) 105019, arXiv:hep-th/9912015;
D. Bazeia, F.A. Brito, Phys. Rev. D 62 (2000) 101701, arXiv:hep-th/0005045;
F.A. Brito, D. Bazeia, Phys. Rev. D 64 (2001) 065022, arXiv:hep-th/0105296.
- [5] P.W. Anderson, et al., Phys. Rev. Lett. 58 (1987) 2790;
E.C. Marino, M.B. Silva Neto, Phys. Rev. B 64 (2001) 092511, arXiv:cond-mat/0008186.
- [6] J.G. Bednorz, K.A. Müller, Z. Phys. B 64 (1986) 189.
- [7] M. Tinkham, Introduction to Superconductivity, 2nd ed., Dover Publications, Inc., Mineola, NY, 1996.
- [8] V.L. Ginzburg, L.D. Landau, Zh. Èksp. Teor. Fiz. 20 (1950) 1064.
- [9] L.P. Gor'kov, Zh. Èksp. Teor. Fiz. 36 (1959) 1918, Sov. Phys. JETP 9 (1959) 1364.
- [10] J. Bardeen, L.N. Cooper, J.R. Schrieffer, Phys. Rev. 106 (1957) 162.
- [11] W.G. Unruh, Phys. Rev. D 14 (1976) 870;
N.D. Birrell, P.C.W. Davies, Quantum Fields in Curved Space, Cambridge Monographs on Mathematical Physics, Cambridge University Press, 1984;
L.C.B. Crispino, A. Higuchi, G.E.A. Matsas, Rev. Mod. Phys. 80 (2008) 787, arXiv:0710.5373 [gr-qc];
D. Roy, Phys. Lett. B 681 (2009) 185, arXiv:0908.3149 [hep-th].