

**UNIVERSIDADE ESTADUAL PAULISTA “JÚLIO DE MESQUITA FILHO”
FACULDADE DE MEDICINA
CAMPUS DE BOTUCATU**

**EFEITO DA PRÓPOLIS SOBRE A PRODUÇÃO DE CITOCINAS
POR CAMUNDONGOS SUBMETIDOS A ESTRESSE CRÔNICO
E DESAFIADOS COM MELANOMA**

FABIANE MISSIMA

Tese apresentada ao Programa de Pós-Graduação em Patologia da Faculdade de Medicina de Botucatu, Universidade Estadual Paulista – UNESP para obtenção do título de Doutor.

BOTUCATU – SP

2009

**UNIVERSIDADE ESTADUAL PAULISTA “JÚLIO DE MESQUITA FILHO”
FACULDADE DE MEDICINA
CAMPUS DE BOTUCATU**

**EFEITO DA PRÓPOLIS SOBRE A PRODUÇÃO DE CITOCINAS
POR CAMUNDONGOS SUBMETIDOS A ESTRESSE CRÔNICO
E DESAFIADOS COM MELANOMA**

Doutoranda: FABIANE MISSIMA

Orientador: PROF. ADJUNTO JOSÉ MAURÍCIO SFORCIN

Tese apresentada ao Programa de Pós-Graduação em Patologia da Faculdade de Medicina de Botucatu, Universidade Estadual Paulista – UNESP para obtenção do título de Doutor.

BOTUCATU – SP

2009

FICHA CATALOGRÁFICA ELABORADA PELA SEÇÃO TÉCNICA DE AQUISIÇÃO E TRATAMENTO
DA INFORMAÇÃO
DIVISÃO TÉCNICA DE BIBLIOTECA E DOCUMENTAÇÃO - CAMPUS DE BOTUCATU - UNESP

Bibliotecária responsável: Selma Maria de Jesus

Missima, Fabiane.

Efeito da própolis sobre a produção de citocinas por camundongos submetidos a estresse crônico e desafiados com melanoma / Fabiane Missima. – Botucatu : [s.n.], 2009.

Tese (doutorado) – Faculdade de Medicina de Botucatu, Universidade Estadual Paulista, 2009.

Orientador: José Maurício Sforcin
Assunto CAPES: 40102009

1. Câncer - Tratamento - Estudos experimentais 2. Câncer - Aspectos imunológicos 3. Estresse 4. Própolis - Uso terapêutico

CDD 616.992

Palavras chave: Citocinas; Corticosterona; Estresse; Melanoma; Própolis

“Cada homem tem seu lugar no mundo e no tempo que lhe é concedido. Sua tarefa nunca é maior que sua capacidade para poder cumpri-la. Ela consiste em preencher seu lugar, em servir a verdade e aos homens.”

(João Guimarães Rosa)

SUMÁRIO

Revisão Bibliográfica	01
Objetivos	10
Referências	11
Manuscrito 1	18
<i>Abstract</i>	20
<i>1. Introduction</i>	21
<i>2. Methods</i>	22
2.1 Animals.....	22
2.2 Melanoma cells.....	22
2.3 Propolis sample.....	23
2.4 Experimental groups and stress procedure.....	23
2.5 Corticosterone determination.....	24
2.6 Spleen cells cultures and cytokines determination.....	25
2.7 Cytokines gene expression.....	25
2.7.1 mRNA extraction and cDNA synthesis.....	25
2.7.2 Primers.....	26
2.7.3 Real-Time PCR.....	27
2.8 Data Analysis.....	27
<i>3. Results</i>	27
3.1 Th1 cytokines' expression and production.....	28
3.2 Th2 cytokines' expression and production.....	29
<i>4. Discussion</i>	30
Acknowledgements.....	34
References.....	34
Graphs.....	38
Manuscrito 2	44
<i>Summary</i>	46
<i>Introduction</i>	47
<i>Materials and Methods</i>	48
Propolis sample.....	48

Melanoma cells.....	49
Animals, experimental groups and stress procedure.....	49
Corticosterone determination.....	50
Spleen cells cultures and cytokines determination.....	50
<i>Statistical analysis</i>	51
<i>Results</i>	51
<i>Discussion</i>	52
Acknowledgements.....	55
References.....	55
Graphs.....	59
Anexo 1	63
Anexo 2	72

Revisão Bibliográfica

1. Revisão Bibliográfica

Em virtude de suas inúmeras propriedades biológicas, dentre elas, a imunomoduladora (SFORCIN, 2007), a própolis foi empiricamente utilizada por séculos.

A própolis é um produto natural, elaborado pelas abelhas a partir de material coletado em botões, gemas e córtex vegetais, sendo um conjunto complexo de substâncias (BANKOVA, 2005). A própolis apresenta, de forma genérica, em sua composição química *in natura* 55% de resinas e bálsamos, 30% de cera, 10% de óleos voláteis e 5% de pólen, além de impurezas mecânicas. GREENAWAY *et al.* (1991) identificaram as estruturas químicas de muitos componentes da própolis: flavonóides, cumarinas, aldeídos aromáticos, ácidos aromáticos, ácidos orgânicos, minerais, vitaminas, aminoácidos, polissacarídeos. O método de preparo dos extratos de própolis pode influenciar sua atividade, uma vez que os diferentes solventes solubilizam e extraem compostos diferentes. Os extratos mais utilizados experimentalmente são os etanólicos, em diferentes concentrações, o metanólico e o aquoso (CUNHA *et al.*, 2004).

A composição química da própolis por nós coletada ao longo das quatro estações na região de Botucatu, SP, foi analisada, verificando-se que diferenças sazonais não são significantes, mas predominantemente quantitativas (BOUDOUROVA-KRASTEVA *et al.*, 1997; BANKOVA *et al.*, 1998). Ademais, as fontes vegetais da própolis obtida no apiário de nossa Universidade foram investigadas, sendo *Baccharis dracunculifolia* a principal fonte, seguida de *Araucaria angustifolia* e *Eucalyptus citriodora* (BANKOVA *et al.*, 1999).

SFORCIN *et al.* (2002b) demonstraram que a administração de própolis a ratos, 2 vezes ao dia, durante 3 dias consecutivos, não induziu alterações na concentração sérica de componentes bioquímicos de significado clínico. Neste trabalho, foi observado também a ausência de efeito da sazonalidade sobre a atividade da própolis em variáveis bioquímica. Dando continuidade a este trabalho, MANI *et al.* (2006) avaliaram o efeito de diferentes concentrações de própolis administradas a ratos, o efeito de diferentes extratos (aquoso e etanólico), e o efeito do tratamento a longo prazo (30, 90 e 150 dias) sobre a concentração sérica de variáveis bioquímicas. Os resultados não evidenciaram diferenças entre o grupo controle e os grupos tratados com própolis, sugerindo a ausência de efeitos colaterais após tratamento com este produto apícola.

Em estudos toxicológicos *in vitro*, TAVARES *et al.* (2006) sugeriram que baixa concentração de própolis (12.5 µg/mL) apresenta efeito antimutagênico, enquanto que alta concentração (100 µg/mL) parece ser mutagênica. PEREIRA *et al.* (2008) avaliaram o efeito mutagênico da própolis *in vivo* em eritrócitos de camundongos tratados com 1000, 1500 e 2000 mg/Kg, concluindo que estas concentrações podem acarretar efeitos mutagênicos.

Atualmente, a própolis tem despertado a atenção de pesquisadores, no intuito de elucidar cientificamente suas propriedades biológicas (BANSKOTA *et al.*, 2001). Nesse sentido, a ação antibiótica da própolis tem sido investigada por nosso grupo, evidenciando-se sua ação antibacteriana, antiviral, antifúngica e anti-parasitária (SFORCIN *et al.*, 2000; SFORCIN *et al.*, 2001; BÚFALO, 2006; FREITAS *et al.*, 2006; ORSI *et al.*, 2006).

Com relação ao sistema imune e seus mecanismos efetores, SCHELLER *et al.* (1988) verificaram que a própolis estimula a formação de anticorpos por células esplênicas de camundongos BALB/c imunizados com hemácias de carneiro. Avaliando

o efeito deste apiterápico sobre a ativação de células *natural killer* (NK) contra células tumorais, observamos que a administração de própolis a ratos, durante 3 dias, induziu aumento na atividade lítica de células NK, quando comparada com o do grupo controle (SFORCIN *et al.*, 2002a). O tratamento com própolis durante 3 dias também induziu aumento na produção de anticorpos em ratos imunizados com albumina sérica bovina (SFORCIN *et al.*, 2005). Estes achados reforçam a afirmação prévia de SCHELLER *et al.* (1988), os quais sugeriram que este apiterápico atua sobre o sistema imune em curto prazo, após sua administração, e que a atividade imunoestimulante da própolis pode estar associada com a ativação de macrófagos e aumento de sua capacidade fagocítica. TATEFUJI *et al.* (1996) estudaram o efeito de seis componentes presentes na própolis e observaram que, dentro das propriedades biológicas apresentadas por este apiterápico, houve um aumento no espriamento de macrófagos e na sua mobilidade.

Trabalhos experimentais de nosso laboratório revelaram também que macrófagos peritoneais de camundongos estimulados com própolis apresentaram aumento na atividade fungicida contra *Paracoccidioides brasiliensis* (MURAD *et al.*, 2002) e na atividade bactericida contra *Salmonella Typhimurium* (ORSI *et al.*, 2005). Foi evidenciado também que a própolis induz elevação na geração de peróxido de hidrogênio (H_2O_2) por macrófagos peritoneais de camundongos e inibição na liberação de óxido nítrico (NO) por estas células, de forma dose-dependente (ORSI *et al.*, 2000).

LOPES *et al.* (2003) avaliaram a ação de extratos das plantas fontes de própolis em nossa região sobre a produção de H_2O_2 e NO por macrófagos, verificando a ausência de efeito sobre a produção de tais metabólitos. Estes resultados sugeriram que a ação da própolis é uma consequência dos produtos derivados das plantas e adição de substâncias secretadas pelas abelhas, atuando possivelmente de forma sinérgica. Considerando que *Baccharis dracunculifolia* DC é a principal fonte de própolis em

nosso apiário e dando continuidade a estes ensaios, avaliamos o efeito do extrato total das partes aéreas, extrato do lavado glandular, extrato das folhas, extrato das raízes e óleo essencial desta planta sobre o estado de ativação de macrófagos, bem como de substâncias purificadas. Os dados revelaram que o extrato do lavado glandular, extrato das folhas e extrato das raízes induziram elevação na geração de H_2O_2 por macrófagos. Com relação aos compostos isolados, o óxido de *Baccharis* e o friedelanol estimularam a produção de H_2O_2 . Estes resultados sugerem uma ação ativadora de extratos e compostos isolados de *B. dracunculifolia* sobre macrófagos (MISSIMA *et al.*, 2007).

Vários pesquisadores têm relatado a propriedade antitumoral da própolis (CHIAO *et al.*, 1995; RAO *et al.*, 1995). Trabalhos de nosso grupo evidenciaram que o extrato hidroalcoólico de própolis possui papel protetor no processo de carcinogênese do cólon, suprimindo o desenvolvimento de lesões pré-neoplásicas (BAZO *et al.*, 2002). Utilizando extrato aquoso de própolis, verificamos que este reduziu os danos induzidos pela dimetilhidrazina no DNA de células do cólon de ratos (ALVES DE LIMA *et al.*, 2005). Em ensaios *in vitro*, verificamos sua eficiente ação citotóxica contra células de tumor venéreo transmissível de cães (BASSANI-SILVA *et al.*, no prelo) e contra células de carcinoma epidermóide de laringe (BÚFALO *et al.*, no prelo), de forma dose- e tempo-dependente.

ORSOLIC *et al.* (2004), avaliando o efeito da própolis, bem como de compostos polifenólicos isolados da mesma, sobre o crescimento e potencial metastático de carcinoma mamário transplantável em camundongos, relataram que a atividade antitumoral está associada à propriedade imunomoduladora dos compostos, à citotoxicidade em relação às células tumorais, indução de apoptose e necrose.

Em virtude de sua ação imunomoduladora, ensaios têm sido desenvolvidos em nossos laboratórios para investigar o papel da própolis durante o estresse. O conceito de

estresse foi apresentado por Hans Selye em 1936, tendo havido, posteriormente, um crescente interesse dos pesquisadores quanto à identificação dos agentes estressores bem como dos eventos fisiológicos envolvidos na resposta ao estresse.

O estresse é comumente definido como uma condição ou estado em que a homeostase do organismo é perturbada, como resultado de estímulos estressores. O estresse é uma constelação de eventos, envolvendo a participação de diferentes sistemas do organismo em resposta a agentes estressores, como fatores climáticos, hiperpopulação, infecções, exercício físico intenso, desnutrição, ruído, odor, entre muitos outros (KIOUKIA-FOUGIA *et al.*, 2002).

A crescente complexidade social também tem contribuído com o aparecimento de tensão psicológica e doenças relacionadas ao estresse, como câncer, depressão, ansiedade, pânico, dermatite atópica, colite ulcerativa, entre outras. Dentre os agentes estressogênicos da vida moderna, temos a competição profissional, desemprego, violência, desagregação familiar, relações interpessoais instáveis, solidão, preocupação financeira persistente, pressão da globalização e tecnologia, entre outras causas de maior impacto, como morte, doença grave, acidentes, entre outros (OLFF, 1999).

Há três fases importantes junto à “Síndrome Geral de Adaptação” ao estresse: a reação de alarme, na qual o organismo percebe o estímulo estressante; o estágio de resistência - uma tentativa de adaptação do organismo frente ao agente estressor; e fase de exaustão, quando o organismo perde a capacidade de adaptação, aparecendo lesões ou doenças (SELYE, 1978).

A resposta ao estresse envolve a ativação do sistema nervoso autônomo (SNA), bem como a ativação do eixo hipotálamo-pituitária-adrenal (HPA), com conseqüente liberação de catecolaminas e glicocorticóides, respectivamente. Em resposta à estimulação simpática, a epinefrina e norepinefrina são rapidamente secretadas para a

corrente sanguínea como uma resposta adaptativa, capacitando o organismo a responder à condição estressante, apresentando efeito pronunciado no sistema cardiovascular e na mobilização nas reservas de energia (LUNDBERG, 2005).

O estresse origina impulsos nervosos, que são transmitidos desde a periferia até o hipotálamo. O hipotálamo, então, secreta o fator liberador de corticotropina (CRH), que passa pelo sistema porta hipotálamo-hipofisário chegando até a hipófise anterior. Neste local, o CRH induz a secreção de hormônio adrenocorticotrópico (ACTH), que flui pelo sangue até o córtex da adrenal, induzindo a secreção de glicocorticóides, como cortisol, corticosterona e aldosterona (SELYE, 1978).

O estresse é uma resposta adaptativa fundamental à sobrevivência, presente não só em mamíferos, mas também em outros vertebrados. Os mecanismos básicos e moléculas envolvidas na resposta ao estresse são similares e bem preservados ao longo da evolução das espécies (OTTAVIANI & FRANCESCHI, 1996).

O estresse agudo pode ser entendido como uma ameaça imediata, a curto prazo, comumente conhecido como resposta à luta ou fuga. A “ameaça” pode ser qualquer situação experienciada como perigo, e, após o evento, que dura minutos a horas, há uma resposta de relaxamento (DHABHAR, 2002). Por outro lado, o estresse é denominado crônico quando persiste por várias horas a dias, semanas ou meses. Como exemplo, temos modelos experimentais com imobilização, exposição à temperatura adversa, exercício físico intenso, entre outros. Modelos experimentais de estresse crônico são apropriados para a investigação da relação estresse-depressão (BAUER *et al.*, 2000).

O modelo de imobilização é um procedimento que não causa dor e o animal não é comprimido fisicamente, mas a sensação de confinamento induz estresse psicológico, afetando o SNA e o eixo HPA, resultando em níveis elevados de catecolaminas e corticosterona (BAUER *et al.*, 2001).

Há vários indicadores de estresse encontrados na literatura pertinente, como concentração de corticosterona, glicemia, atividade da desidrogenase láctica (LDH), aparecimento de úlcera, determinação de hormônio adrenocorticotrófico (ACTH), entre outros. A determinação de cortisol é utilizada como indicador de estresse em seres humanos, e a de corticosterona em roedores (BESEDOVSKY & DEL REY, 1996; LUNDBERG, 2005).

Muitas evidências têm sugerido que o estresse modula a resposta imune (BESEDOVSKY & DEL REY, 1996). A Imunoneuroendocrinologia ou Psiconeuroimunologia tem despertado a atenção dos pesquisadores quanto ao estudo da relação entre os sistemas imune, nervoso e endócrino, os quais compartilham receptores para citocinas, neurotransmissores e hormônios. O comportamento doente reflete esta íntima relação, visto que citocinas podem diminuir a atividade geral do organismo, afetar o comportamento social e sexual, e diminuir a atividade exploratória do indivíduo. Como exemplo, a interleucina-1 (IL-1) é considerada um pirógeno endógeno, induzindo febre e alteração comportamental (PARMET *et al.*, 2002).

Com relação ao sistema imune, os distúrbios resultantes do estresse podem variar de acordo com o tipo, intensidade e duração do agente estressor. Da mesma forma, diferenças quanto à linhagem, idade e sexo dos animais de experimentação, bem como o momento de mensuração de determinados parâmetros, também podem afetar a avaliação da resposta imunológica (KIOUKIA-FOUGIA *et al.*, 2002).

O estresse crônico pode induzir imunossupressão (ELENKOV & CHROUSOS, 1999). Glicocorticóides são considerados imunossupressores, sendo clinicamente utilizados como agente anti-inflamatório. Como linfócitos, monócitos, macrófagos e granulócitos possuem receptores para vários mediadores do SNA e do eixo HPA, como catecolaminas e corticóides (PADGETT & GLASER, 2003), observa-se, no estresse

crônico, inibição na síntese de prostaglandinas, na produção de citocinas e proliferação de linfócitos (DHABHAR, 2002). A produção de IL-3, uma das citocinas fundamentais para hematopoiese, também encontra-se inibida em resposta ao estresse e em indivíduos com depressão, sugerindo a interligação entre sistema nervoso e sistema imune (BESSLER *et al.*, 2000). DOMÍNGUEZ-GERPE & REY-MÉNDEZ (2001) relataram alterações em subpopulações de linfócitos após estresse crônico por imobilização, demonstrando que células T foram mais afetadas que células B.

A resposta imune celular é mais afetada durante o estresse, em relação à imunidade humoral. O estresse pode afetar o padrão Th1/Th2 de resposta, direcionando-o mais para Th2, podendo suprimir a imunidade celular e reforçar a imunidade humoral, por ação de glicocorticóides e catecolaminas (IWAKABE *et al.*, 1998; ELENKOV & CHROUSOS, 1999; ELENKOV, 2004).

A imunossupressão causada pelo estresse pode afetar a metástase de tumores e promover a progressão do câncer (BEN-ELIYAHU, 2003).

O câncer de pele é a terceira neoplasia mais comum a acometer o homem, e sua incidência global tem sido progressiva. As formas mais comuns de neoplasias de pele são os carcinomas de células basais, os carcinomas de células escamosas e os melanomas – a forma mais perigosa e letal. Quando diagnosticado na fase inicial, 80% dos casos de melanomas são curados através de remoção cirúrgica, mas sua forma metastática é altamente refratária às terapias existentes (GRAY-SCHOPFER *et al.*, 2007).

Considerando que o estresse pode induzir imunossupressão e, conseqüentemente, o desenvolvimento de tumores, introduzimos, em nosso modelo experimental, o tratamento dos animais com própolis, visto que este produto apícola apresenta propriedades imunomoduladora e antitumoral. Ademais, poucos são os

trabalhos encontrados na literatura sobre os efeitos da própolis no sistema imunológico, principalmente quanto à determinação de citocinas.

Assim, nosso trabalho visou contribuir quanto ao efeito da própolis sobre a produção de citocinas pró-inflamatórias e de padrão Th1/Th2, bem como sobre o desenvolvimento tumoral *in vivo*, pois há inúmeros trabalhos sobre a ação antitumoral da própolis, sendo a maioria deles, porém, realizada com ensaios *in vitro*.

2. Objetivos

- Avaliar o efeito da própolis sobre a produção de citocinas pró-inflamatórias (IL-1 β e IL-6), de padrão Th1 (IL-2 e IFN- γ) e Th2 (IL-4 e IL-10) através da técnica de ELISA;
- Observar o possível papel imunorestaurador da própolis quanto à produção destas citocinas em animais estressados;
- Quantificar a expressão gênica das citocinas de padrão Th1 e Th2 produzidas por esplenócitos, pela técnica de PCR em tempo real;
- Observar a ação imunomoduladora da própolis em camundongos estressados, portadores ou não de melanoma;
- Avaliar os níveis séricos de corticosterona dos animais.

3. Referências

- ALVES DE LIMA, R.O., BAZO, A.P., SAID, R.A., SFORCIN, J.M., BANKOVA, V., DARROS, B.R., SALVADORI, D.M.F. Modifying effect of propolis on dimethylhydrazine-induced DNA damage but not colonic aberrant crypt foci in rats. **Environmental and Molecular Mutagenesis**, v. 45, p. 8-16, 2005.
- BANKOVA, V., BOUDOUROVA-KRASTEVA, G., POPOV, S., SFORCIN, J.M., FUNARI, S.R.C. Seasonal variations of the chemical composition of Brazilian propolis. **Apidologie**, v. 29, p. 361-67, 1998.
- BANKOVA, V., BOUDOUROVA-KRASTEVA, G., SFORCIN, J.M., FRETE, X., KUJUMGIEV, A., MAIMONI-RODELLA, R., POPOV, S. Phytochemical evidence for the plant origin of Brazilian propolis from São Paulo State. **Zeitschrift für Naturforschung**, v. 54c, p. 401-5, 1999.
- BANKOVA, V. Recent trends and important developments in propolis research. **Evidence-based Complementary and Alternative Medicine**, v. 2, p. 29-32, 2005.
- BANSKOTA, A.H., TEZUKA, V., KADOTA, S. Recent progress in pharmacological research of propolis. **Phytotherapy Research**, v. 15, p. 561-71, 2001.
- BASSANI-SILVA, S., SFORCIN, J.M., AMARAL, A.S., GASPAR, L.F.J., ROCHA, N.S. Propolis effect *in vitro* on venereal transmissible canine tumor. **Revista Portuguesa de Ciências Veterinárias**, no prelo.

*UNIVERSIDADE ESTADUAL PAULISTA. Coordenadoria Geral de Bibliotecas, Editora UNESP. *Normas para publicações da UNESP*. São Paulo: Editora UNESP, 1994. v. 2.: Referências bibliográficas.

BIOSIS. Serial Sources for the BIOSIS' previews data base. Philadelphia, 1991. 451p.

- BAUER, M.E., VEDHARA, K., PERKS, P., WILCOCK, G.K., LIGHTMAN, S.L., SHANKS, N. Chronic stress in caregivers of dementia is associated with reduced lymphocyte sensitivity to glucocorticoids. **Journal of Neuroimmunology**, v. 103, p. 84-92, 2000.
- BAUER, M.E., PERKS, P., LIGHTMAN, S.L., SHANKS, N. Restraint stress is associated with changes in glucocorticoid immunoregulation. **Physiology & Behavior**, v. 73, p. 525-32, 2001.
- BAZO, A.P., RODRIGUES, M.A.M., SFORCIN, J.M., CAMARGO, J.L.V., RIBEIRO, L.R., SALVADORI, D.M.F. Protective action of propolis on the rat colon carcinogenesis. **Teratogenesis, Carcinogenesis and Mutagenesis**, v. 22, p. 183-94, 2002.
- BEN-ELIYAHU, S. The promotion of tumor metastasis by surgery and stress: immunological basis and implications for psychoneuroimmunology. **Brain, Behavior, and Immunity**, v. 17, p. S27-S36, 2003.
- BESEDOVSKY, H.O., DEL REY, A. Immune-endocrine interactions: Facts and hypotheses. **Endocrine Reviews**, v. 17, p. 64-102, 1996.
- BESSLER, H., BERGMAN, M., SALMAN, H. Interleukin-3 and stress. **Biomedicine & Pharmacotherapy**, v. 54, p. 299-304, 2000.
- BOUDOUROVA-KRASTEVA, G., BANKOVA, V., SFORCIN, J.M., NIKOLOVA, N., POPOV, S. Phenolics from Brazilian propolis. **Zeitschrift für Naturforschung**, v. 52c, p. 676-79, 1997.
- BÚFALO, M.C. **Efeito do extrato alcoólico de própolis sobre a replicação do vírus sincicial respiratório humano (RSVH) em células HEp-2**. Monografia apresentada ao Departamento de Microbiologia e Imunologia, Instituto de Biociências, UNESP, Campus de Botucatu, 2006.

- BÚFALO, M.C., CANDEIAS, J.M.G., SFORCIN, J.M. *In vitro* cytotoxic effect of Brazilian green propolis on human laryngeal epidermoid carcinoma (HEp-2) cells. **Evidence-based Complementary and Alternative Medicine**, no prelo.
- CHIAO, C., CAROTHERS, A.M., GRUNBERGER, D., SOLOMON, G., PRESTON, G.A., BARRETT, J.C. Apoptosis and altered redox state induced by caffeic acid phenethyl ester (CAPE) in transformed rat fibroblast cells. **Cancer Research**, v. 55, p. 3576-83, 1995.
- CUNHA, I.B.S., SAWAYA, A.C.H.F., CAETANO, F.M., SHIMIZU, M.T., MARCUCCI, M.C., DREZZA, F.T., POVIA, G.S., CARVALHO, P.O. Factors that influence the yield and composition of Brazilian propolis extracts. **Journal of the Brazilian Chemical Society**, v. 15, p. 964-70, 2004.
- DHABHAR, F.S. Stress-induced augmentation of immune function – the role of stress hormones, leukocyte trafficking, and cytokines. **Brain, Behavior, and Immunity**, v. 16, p. 785-98, 2002.
- DOMÍNGUEZ-GERPE, L., REY-MÉNDEZ, M. Alterations induced by chronic stress in lymphocyte subsets of blood and primary and secondary immune organs of mice. **Immunology**, v. 2, p. 7, 2001.
- ELENKOV, I.J., CHROUSOS, G.P., Stress hormones, Th1/Th2 patterns, pro/anti-inflammatory cytokines and susceptibility to disease. **Trends in Endocrinology and Metabolism**, v. 10, p. 359-68, 1999.
- ELENKOV, I.J. Glucocorticoids and the Th1/Th2 balance. **Annals of the New York Academy of Sciences**, v. 1024, p. 138-46, 2004.
- FREITAS, S.F., SHINOHARA, L., SFORCIN, J.M., GUIMARÃES, S. *In vitro* effects of propolis on *Giardia duodenalis* trophozoites. **Phytomedicine**, v. 13, p. 170-5, 2006.

- GREENAWAY, W., MAJ., J., SAYSBROOK, T., WHATLEY, F.R. Identification by gas chromatography-mass spectrometry of 150 compounds in propolis. **Zeitschrift für Naturforschung**, v. 46, p. 111-21, 1991.
- GRAY-SCHOPFER, V., WELLBROCK, C., MARAIS, R. Melanoma biology and new targeted therapy. **Nature**, v. 445, p. 851-7, 2007.
- IWAKABE, K., SHIMADA, M., YAHATA, T., OHMI, Y., HABU, S., NISHIMURA, T. The restraint stress drives a shift in Th1/Th2 balance toward Th2-dominant immunity in mice. **Immunology Letters**, v. 62, p. 39-43, 1998.
- KIOUKIA-FOUGIA, N., ANTONIOU, K., BEKRIS, S., LIAPI, C., CHRISTOFIDIS, I., PAPADOPOULOU-DAIFOTI, Z. The effects of stress exposure on the hypothalamic-pituitary-adrenal axis, thymus, thyroid hormones and glucose levels. **Progress in Neuro-Psychopharmacology & Biological Psychiatry**, v. 26, p. 823-30, 2002.
- LOPES, F.C., BANKOVA, V., SFORCIN, J.M. Effect of three vegetal sources of propolis on macrophages activation. **Phytomedicine**, v. 10, p. 343, 2003.
- LUNDBERG, U. Stress hormones in health and illness: the roles of work and gender. **Psychoneuroendocrinology**, v. 30, p. 1017-21, 2005.
- MANI, F., DAMASCENO, H.C.R., NOVELLI, E.L.B., MARTINS, E.A.M., SFORCIN, J.M. Propolis: effect of different concentrations, extracts and intake period on seric biochemical variables. **Journal of Ethnopharmacology**, v. 105, p. 95-8, 2006.
- MISSIMA, F., SILVA FILHO, A.A., NUNES, G.A., BUENO, P.C.P., DE SOUSA, J.P.B., BASTOS, J.K., SFORCIN, J.M. Effects of *Baccharis dracunculifolia* D.C. (Asteraceae) extracts and its isolated compounds on macrophage activation. **Journal of Pharmacy and Pharmacology**, v. 59, p. 463-8, 2007.

- MURAD, J.M., CALVI, S.A., SOARES, A.M.V.C., BANKOVA, V., SFORCIN, J.M.
Effects of propolis from Brazil and Bulgaria on fungicidal activity of macrophages against *Paracoccidioides brasiliensis*. **Journal of Ethnopharmacology**, v. 79, p. 331-34, 2002.
- OLFF, M. Stress, depression and immunity: the role of defense and coping styles. **Psychiatry Research**, v. 85, p. 7-15, 1999.
- ORSI, R.O., BANKOVA, V., SFORCIN, J.M. Effects of Brazilian and Bulgarian propolis on bactericidal activity of macrophages against *Salmonella* Typhimurium. **International Immunopharmacology**, v.5, p. 359-68, 2005.
- ORSI, R.O., FUNARI, S.R.C., SOARES, A.M.V.C., CALVI, S.A., OLIVEIRA, S.L., SFORCIN, J.M. BANKOVA, V. Immunomodulatory action of propolis on macrophage activation. **The Journal of Venomous Animals and Toxins**, v. 6, n. 2, p. 205-19, 2000.
- ORSI, R.O., SFORCIN, J.M., FUNARI, S.R.C., FERNANDES JR., A., BANKOVA, V. Synergistic effect of propolis and antibiotics on the *Salmonella* Typhi. **Brazilian Journal of Microbiology**, v. 37, p. 108-12, 2006.
- ORSOLIC, N., KNEZEVIC, A.H., SVER, L., TERZIC, S., BASIC, I. Immunomodulatory and antimetastatic action of propolis and related polyphenolic compounds. **Journal of Ethnopharmacology**, v. 94, p. 307-15, 2004.
- OTTAVIANI, E. FRANCESCHI, C. The neuroimmunology of stress from invertebrates to man. **Progress in Neurobiology**, v. 48, p. 421-40, 1996.
- PADGET, D.A., GLASER, R. How stress influences the immune response. **Trends in Immunology**, v. 24, p. 444-8, 2003.

- PARMET, P., KELLEY, K.W., BLUTHÉ, R.M., DANTZER, R. Expression and regulation of interleukin-1 receptors in the brain. Role of cytokine-induced sickness behavior. **Journal of Neuroimmunology**, v. 125, p. 15-14, 2002.
- PEREIRA, A.D., ANDRADE, S.F., SWERTS, M.S.O., MAISTRO, E.L. First in vivo evaluation of the mutagenic effect of Brazilian green propolis by comet assay and micronucleus test. **Food and Chemical Toxicology**, 2008, doi:10.1016/j.fct.2008.04.001.
- RAO, C.V., DESAI, D., RIVENSON, A., SIMI, B., AMIN, S., REDDY, B.S. Chemoprevention of colon carcinogenesis by phenylethyl-3-methylcaffeate. **Cancer Research**, v. 55, p. 2310-5, 1995.
- SCHELLER, S., GADZA, G., PIETSZ, G., GABRYS, J., SZUMLAS, J., ECKERT, L., SHANI, J. The ability of ethanol extract of propolis to stimulate plaque formation in immunized mouse spleen cells. **Pharmacological Research Communications**, v. 20, p. 323-8, 1988.
- SELYE, H. **The stress of life**. McGraw-Hill Book Co.: New York, 1978.
- SFORCIN, J. M., FERNANDES JÚNIOR, A., LOPES, C. A. M., BANKOVA, V., FUNARI, S. R. C. Seasonal effect on Brazilian propolis antibacterial activity. **Journal of Ethnopharmacology**, v. 73, p. 243-49, 2000.
- SFORCIN, J. M., FERNANDES JÚNIOR, A., LOPES, C. A. M., FUNARI, S. R. C., BANKOVA, V. Seasonal effect of Brazilian propolis on *Candida albicans* and *Candida tropicalis*. **The Journal of Venomous Animals and Toxins**, v. 7, p. 139-44, 2001.
- SFORCIN, J.M., KANENO, R., FUNARI, S.R.C. Absence of seasonal effect on the immunomodulatory action of Brazilian propolis on natural killer activity. **The Journal of Venomous Animals and Toxins**, v. 8, p. 19-29, 2002a.

- SFORCIN, J.M., NOVELLI, E.L.B., FUNARI, S.R.C. Seasonal effect of Brazilian propolis on seric biochemical variables. **The Journal of Venomous Animals and Toxins**, v. 8, p. 244-54, 2002b.
- SFORCIN, J.M., ORSI, R.O., BANKOVA, V. Effect of propolis, some isolated compounds and its source plants on antibody production. **Journal of Ethnopharmacology**, v. 98, p. 301-5, 2005.
- SFORCIN, J.M. Propolis and the immune system: a review. **Journal of Ethnopharmacology**, v. 113, p. 1-14, 2007.
- TATEFUJI, T., IZUMI, N., OHTA, T., ARAI, S., IKEDA, M., KURIMOTO, M. Isolation and identification of compounds from Brazilian propolis which enhance macrophage spreading and mobility. **Biological & Pharmaceutical Bulletin**, v. 19, p. 966-70, 1996.
- TAVARES, D.C., BARCELOS, G.R.M., SILVA, L.F., TONIN, C.C.C., BASTOS, J.K. Propolis-induced genotoxicity and antigenotoxicity in Chinese hamster ovary cells. **Toxicology In Vitro**, v. 20, 1154-1158, 2006.

*Manuscrito 1**

* De acordo com as normas da revista Brain, Behavior, and Immunity. As normas constam no anexo 1 deste trabalho (página 63).

Propolis effect on Th1/Th2 cytokines expression and production by melanoma-bearing mice submitted to stress

Fabiane Missima, Ana C. Pagliarone, Cláudio L. Orsatti, João P. Araújo Júnior, José
Maurício Sforcin*

Department of Microbiology and Immunology, Biosciences Institute, UNESP, Campus
of Botucatu, 18618-000, Brazil

* Corresponding author. Address: Department of Microbiology and Immunology,
Biosciences Institute, UNESP, Campus of Botucatu, 18618-000, Brazil. Fax: +55 14
3811 6058 236. E-mail address: sforcin@ibb.unesp.br (J. M. Sforcin)

Abstract

Several studies have shown that stress affects the immune system, promoting the progression of cancer. Since propolis is a honeybee product showing immunomodulatory and antitumoral activities, the goal of this work was to evaluate propolis effect on Th1 (IL-2 and IFN- γ) and Th2 (IL-4 and IL-10) cytokines' expression and production by melanoma-bearing mice submitted to immobilization stress. In groups without melanoma, stress did not affect IL-2, IL-4 and IL-10 gene expression. Propolis treatment of stressed animals did not influence these cytokines expression as well. On the other hand, IL-2 and IL-10 expression was inhibited in melanoma-bearing mice, stressed or not. Th1 cytokines production was also inhibited in melanoma-bearing mice. Propolis administration to melanoma-bearing mice submitted to stress stimulated IL-2 expression, as well as Th1 (IL-2 and IFN- γ) production, indicating the activation of antitumor cell-mediated immunity. Propolis also stimulated IL-10 expression and production, what may be related to immunoregulatory effects. Our data indicate that propolis showed an immunomodulatory activity in this assay, what may be related to its antitumoral action in vivo. Further investigation will help to understand propolis usefulness during stress.

Keywords: Stress; Melanoma; Cytokines; Propolis; Immunomodulation

1. Introduction

Stress comprises several events, activating physiologic systems in the body to cope with perceived stimuli, and altering the homeostatic state of the organism. Evidences have shown a communication between the immune, endocrine and central nervous systems during stress, resulting in behavioral, endocrine and immunological changes. Stress responses induce activation of both autonomic nervous system and hypothalamic-pituitary-adrenal (HPA) axis, with consequent release of catecholamines and glucocorticoids, respectively (Bauer et al., 2001; Dhabhar, 2002).

Viveros-Paredes et al. (2006) reported that glucocorticoids inhibit the production of IL-2 and IFN- γ by Th1 cells, up-regulate the production of IL-4, IL-10 and IL-13 by Th2 cells, and increase the production of TGF- β by T cells, suppressing cellular immunity and stimulating humoral immunity. Because of glucocorticoids' immunosuppressive effects, stress plays a role in the etiology of many diseases, such as cancer, being detrimental to health.

Melanoma is among the most immunogenic of all solid cancers, and the presence of tumor antigen-specific antibodies and tumor-specific cytotoxic T cells in the peripheral blood of melanoma patients has been well established (Fang et al., 2008). B16F10 is a selective variant cell line obtained from pulmonary metastasis of a melanoma, syngeneic to black C57BL/6 mice (Sá-Rocha et al., 2006). While stress does not seem to affect humoral immunity, a Th2 cytokine profile is associated with down-regulation of the immune response and progression of melanoma (McCarter et al., 2005).

The interaction between stress and immunity has been well documented, but the use of natural products during stress deserves investigation, mainly in melanoma-bearing mice. Propolis is a resinous hive product showing several therapeutic activities,

including immunomodulatory and antitumoral. In vitro and in vivo assays demonstrated the modulatory action of propolis on murine peritoneal macrophages, increasing their microbicidal activity. Its stimulant action on the lytic activity of natural killer cells against tumor cells, and on antibody production was demonstrated (Sforcin, 2007). The antitumoral property of propolis has been investigated both in vivo and in vitro (Sforcin et al., 2002; Búfalo et al., in press).

Based on these observations, the aim of this work was to analyze propolis effects on Th1/Th2 cytokines' expression and production by melanoma-bearing mice submitted to chronic stress, in order to investigate a possible immunorestorative action of propolis in these animals.

2. Methods

2.1. Animals

C57BL/6 male mice aged between 8 and 12 weeks were kept in rooms at 21-25 °C, with a 12 h /12 h light/dark cycle. Food and water were provided *ad libitum*. This work agrees with Ethical Principles in Animal Research adopted by Brazilian College of Animal Experimentation (n° 464).

2.2. Melanoma cells

B16F10 cells were cultured in DMEM (Cultilab, Campinas, SP, Brazil), supplemented with 25 mM HEPES (Sigma – Aldrich, St. Louis, MO, USA) and 10% fetal calf serum. Cell suspensions were detached from the culture flasks using 0.2% trypsin (Nutricell, Campinas, SP, Brazil), and viable cells were counted using a

hemocytometer. Mice were inoculated with 5×10^4 cells in 0.1 mL of phosphate-buffered saline subcutaneously (s.c.) into the right flank region. Tumor development was monitored weekly.

2.3. Propolis sample

Propolis was collected in the Beekeeping Section, UNESP. Propolis was ground and 30% ethanolic extracts of propolis were prepared (30 g of propolis, completing the volume to 100 mL with 70% ethanol), in the absence of bright light, at room temperature, with moderate shaking. After a week, extracts were filtered and the dry weight of the extracts was calculated (120 mg/mL) (Búfalo et al., in press). Propolis chemical composition was investigated using thin-layer chromatography (TLC), gas-chromatography (GC), and gas chromatography-mass spectrometry (GC-MS) analysis (Bankova et al., 1998).

2.4. Experimental groups and stress procedure

Mice were divided into 8 groups (G1, G2, G3, G4, G5, G6, G7, G8), of 8 animals each, according to the experimental design:

	G1	G2	G3	G4	G5	G6	G7	G8
STRESS	-	+	-	+	-	+	-	+
PROPOLIS	-	-	+	+	-	-	+	+
MELANOMA	-	-	-	-	+	+	+	+

G1 was considered as control, and received physiologic solution (NaCl 0.9%). G2 was submitted to restraint stress in a well-ventilated immobilization tube (restrainer) of about 50 mL capacity for 15, 30, 45, 60, 75, 90 and 120 minutes during 7 consecutive days, and from 7th to 14th day, mice were submitted to restraint stress for 2 hours a day, at a fixed time between 8:00 and 11:00 a.m. This procedure is easily performed and causes no physical pain to the animals (Domínguez-Gerpe and Rey-Méndez, 2001; Sforcin et al., 2007).

G3 was treated daily with propolis (200 mg/kg in 0.1 mL, orally), and G4 was treated daily with propolis and submitted to the same stress protocol (Missima and Sforcin, 2008). After 24 h of the respective treatments, animals were sacrificed using a CO₂ inhalation chamber.

G5 was inoculated with B16F10 (5×10^4 cells) s.c. into the right flank. G6 was inoculated with B16F10 cells and submitted to stress. G7 was inoculated with B16F10 cells and treated daily with propolis. G8 was inoculated with B16F10 cells, treated daily with propolis and submitted to stress. All groups had no water and food during stress. After 14 days of melanoma inoculation, mice were sacrificed and metastases were investigated.

70% ethanol (propolis solvent) effects were also investigated after its administration for 14 days to mice.

2.5. Corticosterone determination

Before sacrifice, blood was collected by cardiac puncture and serum was stored at -20° C. Corticosterone concentrations were determined by

radioimmunoassay, using a commercial kit (Coat-A-count, DPC, Los Angeles, CA, USA).

2.6. Spleen cells cultures and cytokines determination

After sacrifice, spleens were aseptically removed and cells were suspended at a concentration of $5 \times 10^6/\text{mL}$ in RPMI 1640 (Cultilab, Campinas, SP, Brazil) supplemented with 10% fetal calf serum and 1% L-glutamine and cultured in flat-bottomed 24-well plates. Cells were cultured in triplicates (1 mL/well) and stimulated with concanavalin A (Con A – 5 $\mu\text{g}/\text{mL}$) for 48h and 5% CO_2 .

Supernatants of spleen cell cultures were collected and assayed for Th1 (IL-2 and IFN- γ) and Th2 (IL-4 and IL-10) cytokines determination by enzyme-linked immunosorbent assay (ELISA), according to manufacturer's instructions (BD Biosciences, San Diego, USA). Briefly, a 96-well flat bottom Maxisorp (Nunc, USA) was coated with capture antibody specific to each cytokine. The plate was washed and blocked before 100 μL of the supernatants and serially diluted specific standards were added to the respective wells. Following a series of washing, the captured cytokine was detected using the specific conjugated detection antibody. The chromogen/substrate reagent was added into each well and, after color development, the plate was read at 450 nm, using an ELISA plate reader (Tan et al., 2006).

2.7. Cytokines gene expression

2.7.1. mRNA extraction and cDNA synthesis

A portion of 30 mg of the spleens was kept in RNA safer (Omega Bio-tek, Inc. USA). Total RNA was extracted with RNAspin extraction kit (GE Healthcare UK

Limited) and further treated with RQ1 RNase-free DNase (Promega, Madison, WI, USA) for 30 min at 37 °C to avoid false-positive results due to amplification of genomic DNA. cDNA was synthesized from 4 µg of total RNA in a final volume of 80 µl reaction using the Improm RT II (Promega, Madison, WI, USA), with random primers at 12.5 ng/µl following manufacturer's protocol.

2.7.2. Primers

Primers were designed based on sequences retrieved from Genbank using the Primer Express (Applied Biosystems, USA) and IDTSciTools (<http://www.idtdna.com>) softwares. For IL-2: primers forward 5'- CCC AAG CAG GCC ACA GAA TTG AAA-3' and reverse 5'- AGT CAA ATC CAG AAC ATG CCG CAG-3' (accession no. **X01772**); for IFN-γ: primers forward 5'- AGA GGA TGG TTT GCA TCT GGG TCA-3' and reverse 5'- ACA ACG CTA TGC AGC TTG TTC GTG-3' (accession no. **M28381**); for IL-4: primers forward 5'- AGA TGG ATG TGC CAA ACG TCC TCA-3' and reverse 5'- AAT ATG CGA AGC ACC TTG GAA GCC-3' (accession no. **NM_021283**); for IL-10: primers forward 5'- GGA CAA CAT ACT GCT AAC CGA C-3' and reverse 5'- TGG ATC ATT TCC GAT AAG GCT TG-3' (accession no. **M37897**). As an endogenous control, primers for β-actin found in literature were used (forward 5'-AAG TGT GAC GTT GAC ATC CGT AA- 3' and reverse 5'-TGC CTG GGT ACA TGG TGG TA-3') (Yang et al., 2006).

2.7.3. Real-time PCR

cDNA (4 μ L) was amplified by real-time PCR with 2X Power Sybr® Green PCR Master Mix and 300 nM target primers. β -actin was used as the housekeeping gene. Each sample was analyzed in duplicates using SDS 7300 (Applied Biosystems, USA). The reaction conditions were as follows: 95 °C/10 min, 40 cycles of 95 °C/15 s, 60 °C/1 min, and the melting curve: 95 °C/15 s, 60 °C/15 s and 95 °C/15 s. PCR amplification was related to a standard-curve. To the smallest dilution of cDNA standard it was given the relative value 100 and, following the same reason of dilution, the other 3 points were 25, 6.25 and 1.56. The results were expressed as percentage of differences relative to normal controls (relative expression) using 7300 systems SDS software (Applied Biosystems, USA).

2.8. Data analysis

Analyses of variance (ANOVA) and Tukey-Kramer multiple comparison test were used to determine differences between the groups. A probability (p) of .05 was chosen as the significant level (Zar, 1999).

3. Results

All melanoma-bearing groups developed a tumor area, reflecting the successful experimental model. Tumors were weighed and no significant differences were seen among the groups due to a high variability. However, stress induced a higher tumor

area, while propolis-treated mice, stressed or not, showed a melanoma development similar to control (data not shown). As to mice behavior, stressed mice (G2) treated with propolis (G4) and bearing melanoma (G6 and G8) were dirty, sweating and with piloerection and tachycardia when leaving the restrainer, in comparison to non-stressed groups. However, a significant increase in corticosterone serum levels (stress indicator) was found only in melanoma-bearing mice submitted to stress (G6) ($F(3.74) = 12.428$, $p < .001$) (Fig. 1).

3.1. *Th1 cytokines' expression and production*

IL-2 gene expression was significantly inhibited in melanoma-bearing groups, stressed or not (G5 and G6) ($F(2.20) = 26.833$, $p < .05$ and $F(3.02) = 26.833$, $p < .01$, respectively). Melanoma-bearing mice treated with propolis, stressed or not, (G7 and G8) showed a significant increase in IL-2 expression ($F(3.38) = 26.833$, $p < .001$) (Fig. 2), what is associated to the basal production of this cytokine in these groups (Fig. 3).

No significant differences were seen in IL-2 basal production in G1-G4. However, increased levels of IL-2 basal production were found in melanoma-bearing mice (G5-G8) ($F(3.38) = 26.775$, $p < .001$), without differences between stressed mice, treated or not with propolis (Fig. 3).

In Con A-stimulated cultures, stress (G2) did not influence IL-2 production ($p > .05$). On the other hand, propolis-treated mice, stressed or not (G3 and G4), showed a significant inhibition in this cytokine production ($F(3.51) = 21.951$, $p < .001$) (Fig. 3). IL-2 was also inhibited in melanoma-bearing mice (G5) in comparison to control ($F(3.12) = 21.951$, $p < .01$). Stressed animals (G6) showed increased concentrations of

IL-2 in comparison to G5 ($F(3.51) = 21.951, p < .001$). Propolis also stimulated IL-2 production by G7 and G8, in comparison to G5, but not significantly (Fig. 3).

mRNA IFN- γ was not detected in the experimental groups. Maybe this cytokine expression occurred under the threshold detection of real-time PCR, or its expression in vivo may have occurred before spleens removal. No differences were found in IFN- γ basal production in G1-G4. Propolis stimulated this cytokine production in melanoma-bearing mice, submitted to stress (G8) ($F(3.38) = 11.838, p < .001$). In Con A-stimulated cultures, propolis inhibited IFN- γ production (G3) ($F(3.38) = 75.585, p < .001$), as well as in stressed mice (G4) ($F(3.38) = 75.585, p < .001$). Melanoma also inhibited IFN- γ production ($F(3.38) = 75.585, p < .001$); nevertheless, propolis treatment of melanoma-bearing mice submitted to stress (G8) led to a higher IFN- γ production in comparison to G5 ($F(3.38) = 75.585, p < .001$) (Fig. 4).

3.2. Th2 cytokines' expression and production

No alterations were found in IL-4 expression in all groups (Fig. 5). IL-4 basal production was not detected, and no significant differences were seen in this cytokine production in Con A-stimulated cultures (Fig. 6).

Propolis treatment (G3) enhanced significantly IL-10 gene expression (Fig. 7) ($F(3.12) = 16.479, p < .01$). Among the melanoma-bearing groups, G5, G6 and G7 showed a significant inhibition in IL-10 expression ($F(2.25) = 16.479, p < .05$); however, propolis treatment of melanoma-bearing mice submitted to stress (G8) induced a similar IL-10 expression in comparison to control (G1).

No alterations were seen in IL-10 basal production, and increased basal IL-10 was observed only in G8 ($F(3.38) = 9.972, p < .001$) (Fig. 8). In Con A-stimulated

cultures, propolis-treated mice, stressed or not, showed a significant inhibition in IL-10 production ($F(3.38) = 48.577, p < .001$). In melanoma-bearing mice stressed or not (G7 and G8), propolis stimulated significantly IL-10 production, ($F(3.38) = 48.577, p < .001$) (Fig. 8). Propolis treatment induced higher IL-10 concentrations stimulated or not by Con A in comparison to G5 ($F(3.38) = 48.577, p < .001$).

Ethanol (propolis solvent) did not influence cytokines production, and propolis effects were exclusively due to its chemical constituents.

4. Discussion

Propolis has been widely investigated lately, and its effect on stressed mice has been the subject of our group (Sforcin et al., 2008). In this work, Th1 and Th2 cytokines' expression and production were analyzed after B16F10 cells inoculation in mice. Although the importance of stress in life appears to be significant, the cellular and molecular mechanisms involved in the pathophysiology of stress still remain largely unclear, and it is important to investigate how stress can influence Th1/Th2 homeostasis, resulting in resistance or facilitation of tumor progression.

A successful B16F10 cells inoculation was observed in our model, and no metastasis was seen after s.c. tumor inoculation. Sá-Rocha et al. (2006) found metastasis in the lung of mice submitted to social stress after 14 days, but B16F10 cells were inoculated into the tail vein. Stressed mice showed a higher tumor area, whereas propolis treatment did not promote tumor growth.

All groups submitted to stress protocols left the restrainer with tachycardia, piloerection, hyperlocomotion, sweating and dirty. HPA axis has an important role in behavioral and immunological responses during stress, and glucocorticoids play a major regulatory role in long-term stress in mammals. Higher levels of corticosterone were

found only in melanoma-bearing mice submitted to stress. Controversial data on corticosterone concentrations are found in literature, and biological effects depend on the type of stress, intensity, time of measurement of a particular parameter, and mice strains (Kioukia-Fougia et al., 2002). Bowers et al. (2007) related that different stressors commonly used in research may not activate the physiological response to the same extent, and variations in the measured immune responses may reflect differential glucocorticoid activation, differential metabolic pathways, or both processes in response to specific stressors.

Previous observations of our group revealed that stress (immobilization for 7 consecutive days) induced a higher hydrogen peroxide (H_2O_2) generation by peritoneal macrophages. Propolis treatment potentiated H_2O_2 generation and counteracted the alterations found in the spleen of stressed mice (Missima and Sforcin, 2008). However, propolis effect on cytokines' expression and production during chronic stress was little investigated.

Cytokines play an important role in controlling tumor growth and metastasis. A vigorous Th1 response is required for the destruction of tumor cells, while a Th2 response would create a tolerogenic environment in which melanoma could grow (McCarter et al., 2005).

In groups without melanoma, stress did not affect Th1/Th2 cytokines' expression or production, while propolis inhibited Th1 (IFN- γ and IL-2) and Th2 (IL-10) production, suggesting the anti-inflammatory action of this apitherapeutic agent after 14 of administration. Melanoma-bearing mice showed an inhibition in Th1 cytokines production. Thejass and Kuttan (2007) observed decreased IL-2 plasma levels in C57BL/6 mice of B16F10 inoculation. On the other hand, Posevitz et al. (2003) found no consistent changes in the mRNAs of TNF- α , TNF- β , IL-1 α , IL-1 β , IL-6, IL-4,

IL-5, IFN- γ , IL-2, and IL-3 of C57BL/6 or DBA/2 melanoma-bearing mice submitted to immobilization. Propolis administration to melanoma-bearing mice submitted to stress enhanced IL-2 expression, an increased the production of IFN- γ and IL-2, indicating the activation of antitumor cell-mediated immunity.

IL-2 is known for its potent ability to activate CD8⁺ T cells and natural killer cells, and several studies have been conducted to investigate the combinations of IL-2 and chemotherapy in advanced melanoma (Fang et al., 2008). IFN- γ might be involved in antitumor and antiangiogenic activities (Corthay et al., 2005). Gollob et al. (2005), through DNA microarray analysis, found that the antimelanoma effect of IFN- γ in human melanoma DM6 cells was associated with the down-regulation of multiple genes involved in G-protein signaling and phospholipase C activation (including Rap2B and calpain 3) as well as the down-regulation of genes involved in melanocyte/melanoma survival (MITF and SLUG), apoptosis inhibition (Bcl2A1 and galectin-3), and cell cycling (CDK2). However, although IFN- γ has been generally thought to enhance antitumor immune responses, it has also been reported that IFN- γ could promote tumor immune evasion (He et al., 2005).

Patients with advancing or progressing melanoma have frequently circulating cytokine levels of Th2 pattern (McCarter et al., 2005). These authors suggested that melanoma does have a direct effect on T cells, but the mechanisms involved in the genesis of a sustained Th2 response in cancer patients is still unclear. IL-10 exerts immunosuppressive effects and may act as a cancer-promoting agent. No changes in IL-4 expression and production were seen. Propolis stimulated IL-10 expression and production, what may be associated to immunoregulatory effects. However, it has been reported that IL-10 also possesses immunostimulatory and anti-angiogenic effects and thus may inhibit tumor growth (Vuoristo, 2007).

In our work, systemic levels of cytokines were not determined, nor melanoma-derived cytokines, but data from our laboratory showed that propolis stimulated pro-inflammatory cytokines production (IL-1 β and IL-6) by melanoma-bearing mice submitted to stress (Missima et al., not submitted). TNF- α , IL-1 β and IL-6 may act as autocrine growth factors for tumor angiogenesis (Thejass and Kuttan, 2007). However, both tumor-promoting and tumor-suppressing effects of TNF- α have been reported. Corthay et al. (2005) reported that TNF- α may act in synergy with IFN- γ for the activation of tumoricidal macrophages. Since propolis induced higher levels of cytokines in G8, one may speculate that a synergistic effect of IFN- γ and pro-inflammatory cytokines could inhibit tumor growth in vivo by inducing the production of antiangiogenic factors by tumor cells.

Herbal drugs are recognized as cancer chemopreventive agents. Several plant extracts and isolated compounds are well known to suppress tumor cell growth due to apoptosis in different tumor cell lines (Guruvayoorappan and Kuttan, 2007). It has been demonstrated in vitro that propolis possesses antitumoral active compounds, such as caffeic acid phenethyl ester and artemisinic acid; however, the antiproliferative action of propolis on tumor cells may be the result of the synergistic effect of propolis constituents (Búfalo et al., in press). In vivo, propolis-treated rats showed an increased cytotoxic activity of natural killer cells (Sforcin et al., 2002). Orsolic et al. (2006) suggested that the antitumor activity of propolis and some of its constituents is associated with the augmentation of non-specific antitumor immunity, via macrophages activation, which in turn could produce soluble factors and interfere directly in the tumor cells or in the functions of other immune cells. Propolis administration to rats after treatment with the carcinogen 1,2 dimethylhydrazine reduced significantly the number of aberrant crypt foci, reflecting a suppression of the clonal expansion of the

initiated cells that characterizes the promotion step of carcinogenesis (Bazo et al., 2002).

The polarization of Th1 cells is thought to contribute to cellular immune response and increase CD8⁺ cytotoxic T lymphocytes, playing an important role in anti-tumor immune surveillance (Hu et al., 2007). Our data indicate that propolis showed an immunomodulatory activity in this assay, what may be related to its antitumoral action in vivo, but this aspect still deserves further investigation to understand propolis efficacy in tumor-bearing mice, mainly during stress.

Acknowledgements

We thank Dr. Alexandre Corthay, Institute of Immunology, University of Oslo, Norway, for the critical reading of the manuscript, and Dr. Lídia Raquel de Carvalho, Biosciences Institute, UNESP, Campus of Botucatu, for the statistical analysis. This work was supported by FAPESP (2005/52566-1) and CAPES (199-01/2007).

References

- Bankova, V., Boudourova-Krasteva, G., Popov, S., Sforcin, J.M., Funari, S.R.C., 1998. Seasonal variations of the chemical composition of Brazilian propolis. *Apidologie* 29, 361-367.
- Bauer, M.E., Perks, P., Lightman, S.L., Shanks, N., 2001. Restraint stress is associated with changes in glucocorticoid immunoregulation. *Physiol. Behav.* 73, 525-532.
- Bazo, A.P., Rodrigues, M.A.M., Sforcin, J.M., Camargo, J.L.V., Ribeiro, L.R., Salvadori, D.M.F., 2002. Protective action of propolis on the rat colon carcinogenesis. *Teratog. Carcinog. Mutagen.* 22, 183-194.

- Bowers, S.L., Bilbo, S.D., Dhabhar, F.S., Nelson, R.J., 2008. Stressor-specific alterations in corticosterone and immune responses in mice. *Brain Behav. Immun.* 22, 105-113.
- Búfalo, M.C., Candeias, J.M.G., Sforcin, J.M. *In vitro* cytotoxic effect of Brazilian green propolis on human laryngeal epidermoid carcinoma (HEp-2) cells. *Evid. Based Complement. Alternat. Med.* In press.
- Corthay, A., Skovseth, D.K., Lundin, K.U., Rosjo, E., Omholt, H., Hofgaard, P.O., Haraldsen, G., Bogen, B., 2005. Primary antitumor immune response mediated by CD4⁺ T cells. *Immunity* 22, 371-383.
- Dhabhar, F.S., 2002. Stress-induced augmentation of immune function – the role of stress hormones, leukocyte trafficking, and cytokines. *Brain Behav. Immun.* 16, 785-798.
- Domínguez-Gerpe, L., Rey-Méndez, M., 2001. Alterations induced by chronic stress in lymphocyte subsets of blood and primary and secondary immune organs of mice. *BMC Immunol.* 2, 7.
- Fang, L., Lonsdorf, A.S., Hwang, S.T., 2008. Immunotherapy for advanced melanoma. *J. Invest. Dermatol.* 128, 2596-2605.
- Gollob, J.A., Sciambi, C.J., Huang, Z., Dressman, H.K., 2005. Gene expression changes and signaling events associated with the direct antimelanoma effect of IFN- γ . *Cancer Res.* 65, 8869-8877.
- Guruvayoorappan, C., Kuttan, G., 2007. Apoptotic effect of *Biophytum sensitivum* on B16F-10 cells and its regulatory effects on nitric oxide and cytokine production on tumor-associated macrophages. *Integr. Cancer Ther.* 6, 373-380.
- He, Y., Wang, X., Zhang, G., Chen, H., Zhang, H., Feng, Z., 2005. Sustained low-level expression of interferon- γ promotes tumor development: potential insights in

- tumor prevention and tumor immunotherapy. *Cancer Immunol. Immunother.* 54, 891-897.
- Hu, K., Xiong, J., Ji, K., Sun, H., Wang, J., Liu, H., 2007. Recombined CC chemokine ligand 2 into B16 cells induces production of Th2-dominant cytokines and inhibits melanoma metastasis. *Immunol. Lett.* 113, 19-28.
- Kioukia-Fougia, N., Antoniou, K., Bekris, S., Liapi, C., Christofidis, I., Papadopoulou-Daifoti, Z., 2002. The effects of stress exposure on the hypothalamic-pituitary-adrenal axis, thymus, thyroid hormones and glucose levels. *Progr. NeuroPsychopharmacol. Biol Psychiatry* 26, 823-830.
- Mccarter, M., Clarke, J., Richter, D., Wilson, C., 2005. Melanoma skews dendritic cells to facilitate a T helper 2 profile. *Surgery* 138, 321-328.
- Missima, F., Pagliarone, A.C., Orsatti, C.L., Sforcin, J.M. Propolis effect on pro-inflammatory cytokines produced by melanoma-bearing mice submitted to stress. *J. Apiprodukt and ApiMedical Sci.*, not submitted.
- Missima, F., Sforcin, J.M., 2008. Green Brazilian propolis action on macrophages and lymphoid organs of chronically stressed mice. *Evid. Based Complement. Alternat. Med.* 5, 71-75.
- Orsolic, N., Saranovic, A.B., Basic, I., 2006. Direct and indirect mechanism(s) of antitumor activity of propolis and its polyphenolic compounds. *Planta Med.* 72, 20-27.
- Posevitz, V., Vizler, C., Benyhe, S., Duda, E., Borsodi, A., 2003. Restraint stress and anti-tumor immune response in mice. *Acta Biol. Hung.* 54, 167-176.
- Sá-Rocha, V.M., Sá-Rocha, L.C., Palermo-Neto, J., 2006. Variations in behavior, innate immunity and host resistance to B16F10 melanoma growth in mice that present social stable hierarchical ranks. *Physiol. Behav.* 88, 108-115.

- Sforcin, J.M., Nunes, G.A., Missima, F., Sá-Nunes, A., Faccioli, L.H., 2007. Effect of a leukotriene inhibitor (MK886) on nitric oxide and hydrogen peroxide production by macrophages of acutely and chronically stressed mice. *J. Pharm. Pharmacol.* 59, 1249-1254.
- Sforcin, J.M., Kaneno, R., Funari, S.R.C., 2002. Absence of seasonal effect on the immunomodulatory action of Brazilian propolis on natural killer activity. *J. Venomous Anim. Toxins* 8, 19-29.
- Sforcin, J.M., Missima, F., Orsatti, C., Pagliarone, A., Kaneno, R., 2008. Propolis effect on Th1/Th2 cytokine profile in melanoma-bearing mice submitted to stress. *Scand. J. Immunol.* 68, 216-217.
- Sforcin, J.M., 2007. Propolis and the immune system: a review. *J. Ethnopharmacol.* 113, 1-14.
- Tan, E.L., Selvaratnam, G., Kananathan, R., Sam, C.K., 2006. Quantification of Epstein-Barr virus DNA load, interleukin-6, interleukin-10, transforming growth factor- β 1 and stem cell factor in plasma of patients with nasopharyngeal carcinoma. *BMC Cancer* 6, 227.
- Thejass, P., Kuttan, G., 2007. Inhibition of endothelial cell differentiation and proinflammatory cytokine production during angiogenesis by allyl isothiocyanate and phenyl isothiocyanate. *Integr. Cancer Ther.* 6, 389-399.
- Viveros-Paredes, J.M., Puebla-Pérez, A.M., Gutiérrez-Coronado, O., Sandoval-Ramírez, L., Villaseñor-García, M.M., 2006. Dysregulation of the Th1/Th2 cytokine profile is associated with immunosuppression induced by hypothalamic-pituitary-adrenal axis activation in mice. *Int. Immunopharmacol.* 6, 774-781.

- Vuoristo, M.S., 2007. The polymorphisms of interleukin-10 gene influence the prognosis of patients with advanced melanoma. *Cancer Genet. Cytogenet.* 176, 54-57.
- Yang, H., Xu, Z., Iuvone, P.M., Grossniklaus, H.E., 2006. Angiostatin decreases cell migration and vascular endothelium growth factor (VEGF) to pigment epithelium derived factor (PEDF) RNA ratio in vitro and in a murine ocular melanoma model. *Mol. Vis.* 12, 511-517.
- Zar, J.H., 1999. *Biostatistical Analysis*. New Jersey, Prentice Hall.

Graphs

Fig. 1. Corticosterone concentration (ng/mL). G1: control; G2: stress; G3: propolis; G4: propolis + stress; G5: melanoma; G6: melanoma + stress; G7: melanoma + propolis; G8: melanoma + propolis + stress. Data represent means and standard-deviation of 8 animals. * significantly different from G1 ($p < .001$).

Fig. 2. Relative IL-2 mRNA quantification. Control group received the relative value of 100. G1: control; G2: stress; G3: propolis; G4: propolis + stress; G5: melanoma; G6: melanoma + stress; G7: melanoma + propolis; G8: melanoma + propolis + stress. Data represent means and standard-deviation of 8 animals. * significantly different from G1 ($p < .05$). ** significantly different from G1 ($p < .01$). *** significantly different from G1 ($p < .001$).

Fig. 3. IL-2 production (pg/mL) by spleen cells stimulated or not with concanavalin A (5 μ g/mL) for 48 h. G1: control; G2: stress; G3: propolis; G4: propolis + stress; G5: melanoma; G6: melanoma + stress; G7: melanoma + propolis; G8: melanoma + propolis + stress. Data represent means and standard-deviation of 8 animals. * significantly different from G1 ($p < .001$). # significantly different from G1 + Con A

($p<.01$). ^{##} significantly different from G1 + Con A ($p<.001$). [♦] significantly different from G5 + Con A ($p<.001$).

Fig. 4. IFN- γ production (pg/mL) by spleen cells stimulated or not with concanavalin A (5 μ g/mL) for 48 h. G1: control; G2: stress; G3: propolis; G4: propolis + stress; G5: melanoma; G6: melanoma + stress; G7: melanoma + propolis; G8: melanoma + propolis + stress. Data represent means and standard-deviation of 8 animals. * significantly different from G1 ($p<.001$). [#] significantly different from G1 + Con A ($p<.001$). [♦] significantly different from G5 + Con A ($p<.001$).

Fig. 5. Relative IL-4 mRNA quantification. Control group received the relative value of 100. G1: control; G2: stress; G3: propolis; G4: propolis + stress; G5: melanoma; G6: melanoma + stress; G7: melanoma + propolis; G8: melanoma + propolis + stress. Data represent means and standard-deviation of 8 animals.

Fig. 6. IL-4 production (pg/mL) by spleen cells stimulated with concanavalin A (5 μ g/mL) for 48 h. G1: control; G2: stress; G3: propolis; G4: propolis + stress; G5: melanoma; G6: melanoma + stress; G7: melanoma + propolis; G8: melanoma + propolis + stress. Data represent means and standard-deviation of 8 animals.

Fig. 7. Relative IL-10 mRNA quantification. Control group received the relative value of 100. G1: control; G2: stress; G3: propolis; G4: propolis + stress; G5: melanoma; G6: melanoma + stress; G7: melanoma + propolis; G8: melanoma + propolis + stress. Data represent means and standard-deviation of 8 animals. * significantly different from G1 ($p<.01$).

Fig. 8. IL-10 production (pg/mL) by spleen cells stimulated or not with concanavalin A (5 μ g/mL) for 48 h. G1: control; G2: stress; G3: propolis; G4: propolis + stress; G5: melanoma; G6: melanoma + stress; G7: melanoma + propolis; G8: melanoma + propolis + stress. Data represent means and standard-deviation of 8 animals. * significantly different from G1 ($p<.001$). [#] significantly different from G1 + Con A ($p<.001$). [♦] significantly different from G5 ($p<.001$). [•] significantly different from G5 + Con A ($p<.001$).




Fig. 1.




Fig. 2.




Fig. 3.




Fig. 4.




Fig. 5.




Fig.6.




Fig. 7.




Fig. 8.

*Manuscrito 2**

De acordo com as normas da revista Journal of ApiProduct and ApiMedical Science. As normas constam no anexo 2 deste trabalho (página 72).

Propolis effect on pro-inflammatory cytokines produced by melanoma-bearing mice submitted to chronic stress

Running title: Propolis effect on melanoma-bearing mice

Fabiane Missima, Ana C. Pagliarone, Cláudio L. Orsatti, José M. Sforcin

*Department of Microbiology and Immunology, Biosciences Institute, UNESP,
Campus of Botucatu, 18618-000, Brazil*

* Corresponding author:

Prof. José M. Sforcin

Tel.: +55 14 3811 6058, Fax: +55 14 3811 6058 236.

E-mail sforcin@ibb.unesp.br

Summary

Altered levels of pro-inflammatory cytokines could be prometastatic and proangiogenic factors. Stress leads to somatic disorders such as cancer, and animal models have been used to investigate stress effects on tumorigenesis and immune response modulation. Propolis is a honeybee product and its immunomodulatory effects on stressed mice have been the goal of our recent researches. In this work, propolis effect on pro-inflammatory (IL-1 β and IL-6) cytokines production by melanoma-bearing mice submitted to immobilization stress were analyzed. Propolis administration to stressed mice inhibited pro-inflammatory cytokines production. On the other hand, propolis induced higher levels of IL-1 β and IL-6 in melanoma-bearing mice submitted or not to chronic stress. Since propolis also stimulated Th1 cytokines production in these mice, one may speculate that a synergistic effect of IFN- γ and pro-inflammatory cytokines could inhibit tumor growth in vivo by inducing the production of antiangiogenic factors. Further investigation will help to understand propolis usefulness during stress.

Keywords: Interleukin-1 β ; Interleukin-6; Stress; Melanoma; Propolis; Immunomodulation

Introduction

Propolis is a resinous hive product, used in folk medicine since ancient times, due to its many biological properties, such as antimicrobial (Sforcin *et al.*, 2000; Freitas *et al.*, 2006; Búfalo *et al.*, submitted), antitumor (Banskota *et al.*, 2001; Bazo *et al.*, 2002), immunomodulatory (Sforcin, 2007), among others. Analyses of its chemical composition have identified at least 300 compounds in propolis (De Castro, 2001), and its effects on stressed mice have been investigated by our research group (Missima and Sforcin, 2008).

Stress comprises several nonspecific events, altering the homeostatic state of the organism and leading to “sickness behavior”. Stress responses induce co-activation of both sympathoadrenomedullary system and hypothalamic-pituitary-adrenal (HPA) axis, with consequent release of catecholamines and glucocorticoids, respectively (Dhabhar, 2002). Psychological and behavioral changes are associated to physiological changes, evidencing a communication between the immune, endocrine and central nervous systems during stress.

Catecholamines, glucocorticoids and pro-inflammatory cytokines are considered among the principal messengers responsible for the bi-directional communication between the central nervous system and the immune system (Maes *et al.*, 1998). IL-1 β , also called endogenous pyrogen, is synthesized primarily by monocytes and macrophages, and contributes to the pathogenesis of chronic inflammatory diseases (Pope and Tschopp, 2007). IL-6 is a typical pleiotropic cytokine, playing an important role in the homeostasis of the immune system. IL-6 is one of the major cytokines that stimulates the HPA axis during inflammatory stress (Sjogren *et al.*, 2006).

Because of glucocorticoids' immunosuppressive effects, stress plays a role in the etiology of many diseases, such as cancer, being detrimental to health. Melanoma is among the most immunogenic of all solid cancers, and the presence of tumor antigen-specific antibodies and tumor-specific cytotoxic T cells in the peripheral blood of melanoma patients has been well established (Fang *et al.*, 2008). B16F10 is a selective variant cell line obtained from pulmonary metastasis of a melanoma, syngeneic to black C57BL/6 mice (Sá-Rocha *et al.*, 2006). Moreover, altered levels of IL-1 β , IL-6 and TNF- α could be prometastatic and proangiogenic factors and their deregulated expression directly correlates with the metastatic potential of several forms of cancer.

Whereas the relationship between stress and immunity has been extensively studied in the last decades, the use of natural products during stress deserves investigation, mainly in melanoma-bearing mice. We report herein propolis' immunomodulatory action on pro-inflammatory (IL-1 β and IL-6) cytokines production by melanoma-bearing mice submitted to chronic stress.

Materials and Methods

Propolis sample

Propolis was collected in the Beekeeping Section, UNESP. Propolis was ground and 30% ethanolic extracts of propolis were prepared (30 g of propolis, completing the volume to 100 mL with 70% ethanol), in the absence of bright light, at room temperature, with moderate shaking. After a week, extracts were filtered and the dry weight of the extracts was calculated (120 mg/mL) (Sforcin *et al.*, 2005). Propolis chemical composition was investigated using thin-layer chromatography (TLC), gas-

chromatography (GC), and gas chromatography-mass spectrometry (GC-MS) analysis (Bankova *et al.*, 1998).

Melanoma cells

B16F10 cells were cultured in DMEM (Cultilab, Campinas, SP, Brazil), supplemented with 25 mM HEPES (Sigma – Aldrich, St. Louis, MO, USA) and 10% fetal calf serum. Cell suspensions were detached from the culture flasks using 0.2% trypsin, and viable cells were counted using a hemocytometer. Mice were inoculated with 5×10^4 cells in 0.1 mL of phosphate-buffered saline subcutaneously (s.c.) into the right flank region. Tumor development was monitored weekly.

Animals, experimental groups and stress procedure

C57BL/6 male mice aged between 8 and 12 weeks were kept in rooms at 21-25°C, with a 12 h /12 h light/dark cycle. Food and water were provided *ad libitum*. This work agrees with Ethical Principles in Animal Research adopted by Brazilian College of Animal Experimentation (n° 464).

Mice were divided into 8 groups (G1, G2, G3, G4, G5, G6, G7, G8), of 8 animals each. G1 was considered as control, and received physiologic solution (NaCl 0.9%, 0.1 mL). G2 was submitted to restraint stress in a well-ventilated immobilization tube (restrainer) of about 50 mL capacity for 15, 30, 45, 60, 75, 90 and 120 minutes during 7 consecutive days, and from 7th to 14th day, mice were submitted to restraint stress for 2 hours a day, at a fixed time between 8.00 and 11.00 a.m. This procedure is easily performed and causes no physical pain to the animals (Dominguez-Gerpe and Rey-Méndez, 2001; Sforcin *et al.*, 2007).

G3 was treated daily with propolis (200 mg/kg in 0.1 mL, orally), and G4 was treated daily with propolis and submitted to the same stress protocol (Missima and Sforcin, 2008). After 24 h of the respective treatments, animals were sacrificed using a CO₂ inhalation chamber.

G5 was inoculated with B16F10 (5×10^4 cells) s.c. into the right flank. G6 was inoculated with B16F10 cells and submitted to stress. G7 was inoculated with B16F10 cells and treated daily with propolis. G8 was inoculated with B16F10 cells, treated daily with propolis and submitted to stress. All groups had no water and food during stress. After 14 days of melanoma inoculation, mice were sacrificed and metastases were investigated.

70% ethanol (propolis solvent) effects were also investigated after its administration for 14 days to mice.

Corticosterone determination

Before sacrifice, blood was collected by cardiac puncture and serum was stored at -20°C . Corticosterone concentrations were determined by radioimmunoassay, using a commercial kit (Coat-A-count, DPC, Los Angeles, CA, USA).

Spleen cells cultures and cytokines determination

After sacrifice, spleens were aseptically removed and cells were suspended at a concentration of $5 \times 10^6/\text{mL}$ in RPMI 1640 (Cultilab, Campinas, SP, Brazil) supplemented with 10% fetal calf serum and 1% L-glutamine and cultured in flat-bottomed 24-well plates. Cells were cultured in triplicates (1 mL/well) and stimulated with lipopolysaccharide (LPS – $5 \mu\text{g}/\text{mL}$) for 48h at 37°C and 5% CO₂.

Supernatants of spleen cell cultures were collected and assayed for IL-1 β and IL-6 cytokines determination by enzyme-linked immunosorbent assay (ELISA), according to manufacturer's instructions (BD Biosciences, San Diego, USA). Briefly, a 96-well flat bottom Maxisorp (Nunc, USA) was coated with capture antibody specific to each cytokine. The plate was washed and blocked before 100 μ L of the supernatants and serially diluted specific standards were added to the respective wells. Following a series of washing, the captured cytokine was detected using the specific conjugated detection antibody. The substrate reagent was added into each well and, after color development, the plate was read at 450 nm, using an ELISA plate reader (Tan *et al.*, 2006).

Statistical analysis

Analyses of variance (ANOVA) and Tukey-Kramer multiple comparison test were used to determine differences between the groups. A probability (p) of 0.05 was chosen as the significant level (Zar, 1999).

Results

All melanoma-bearing groups developed a tumor area, reflecting a successful experimental model. Tumors were weighed and no significant differences were seen among the groups due to a high variability. However, stress induced a higher tumor area, while propolis-treated mice, stressed or not, showed a melanoma development similar to control (data not shown). As to mice behavior, stressed mice (G2) treated with propolis (G4) and bearing melanoma (G6 and G8) were dirty, sweating and with piloerection and tachycardia when leaving the restrainer, in comparison to non-stressed

groups. However, a significant increase in corticosterone serum levels (stress indicator) was found only in melanoma-bearing mice submitted to stress ($p<0.001$) (Fig. 1).

IL-1 β concentration was detected only in LPS-stimulated cultures. Propolis-treated mice and stressed or not (G3 and G4) showed a significant inhibition this cytokine production ($p<0.01$). In melanoma-bearing groups, increasing concentrations of IL-1 β were seen in propolis-treated mice submitted or not to stress (G7 and G8) (Fig. 2).

IL-6 basal production was elevated only melanoma-bearing groups ($p<0.05$). In LPS-stimulated cultures, propolis treatment of stressed mice (G4) inhibited IL-6 production ($p<0.05$). On the other hand, increased concentrations of IL-6 ($p<0.05$) were found after propolis administration to melanoma-bearing mice submitted or not to stress (G7 and G8) (Fig. 3).

Ethanol (propolis solvent) did not influence cytokines nor corticosterone production, and propolis effects were exclusively due to its chemical constituents.

Discussion

Propolis action on stressed mice has attracted our group's interest (Missima and Sforcin, 2008; Sforcin *et al.*, 2008). Since propolis effect on cytokines production during stress was little investigated, pro-inflammatory cytokines production was analyzed after B16F10 cells inoculation in stressed mice.

A successful B16F10 cells inoculation was observed in our model, and no metastasis was seen after subcutaneous tumor inoculation. Although no significant differences were seen in tumor weight among the groups, stress induced a higher tumor

area, whereas propolis treatment did not affect tumor growth. Metastasis was found in the lung of mice submitted to social stress after 14 days when B16F10 cells were inoculated into the tail vein (Sá-Rocha *et al.*, 2006).

Pro-inflammatory cytokines are known to activate the HPA axis and consequently increase glucocorticoids levels. However, stressed mice (G2) showed no alterations in IL-1 β , IL-6 and corticosterone concentrations. With respect to mice behavior, stressed groups (G2, G4, G6 and G8) were dirty, sweating and with piloerection when leaving the restrainer, but higher levels of corticosterone were found only in melanoma-bearing mice submitted to stress (G6), what was associated to increased IL-6 production. HPA axis has an important role in behavioral and immunological responses during stress, but controversial data on corticosterone concentrations are found in literature. Moreover, biological effects depend on the intensity and type of stress, time of measurement of a particular parameter, and mice strains (Kioukia-Fougia *et al.*, 2002). Different stressors may not activate the physiological response to the same extent, and variations in immunological parameters reflect differential glucocorticoid activation and metabolic pathways in response to specific stressors (Bowers *et al.*, 2008).

Pro-inflammatory cytokines have overlapping activities, and their production is also increased in acute inflammatory responses associated with infection, injury, trauma and stress (Kamimura *et al.*, 2003; Avitsur *et al.*, 2006). Pro-inflammatory cytokines could be also prometastatic and proangiogenic factors (Thejass and Kuttan, 2007). Cytokines play an important role in controlling tumor growth and metastasis. A vigorous Th1 response is required for the destruction of tumor cells, while a Th2 response would create a tolerogenic environment in which melanoma could grow (McCarter *et al.*, 2005). Works of our laboratory revealed that propolis stimulated IFN- γ

and IL-2 production by melanoma-bearing mice submitted to stress (G8), indicating the activation of antitumor cell-mediated immunity (Missima *et al.*, not submitted). In this work, propolis administration to G8 induced higher levels of IL-1 β and IL-6. Both tumor-promoting and tumor-suppressing effects of TNF- α have been reported. TNF- α may act in synergy with IFN- γ for the activation of tumoricidal macrophages (Corthay *et al.*, 2005). Since propolis stimulated pro-inflammatory and Th1 cytokines production in G8, one may speculate that a synergistic effect of IFN- γ and pro-inflammatory cytokines could inhibit tumor growth *in vivo* by inducing the production of antiangiogenic factors. Our results are in agreement with other authors, who related that changes in the production of the pro-inflammatory cytokines and IFN- γ take part in the homeostatic responses to psychosocial stress in humans, and that stress-induced anxiety is related to a Th1 response (Maes *et al.*, 1998).

Several researchers have reported the antitumoral property of propolis *in vivo* and *in vitro*. Propolis antiproliferative activity on tumor cells has been demonstrated and some responsible compounds were isolated (Banskota *et al.*, 2001; Sforcin *et al.*, 2002; Orsolic *et al.*, 2006; Sforcin, 2007). On the basis of these findings, our data suggest that propolis exerted a possible antitumoral action *in vivo* enhancing pro-inflammatory and IFN- γ production, but further investigation is still needed to evaluate propolis usefulness in tumor-bearing mice, during stress conditions.

Acknowledgements

This work was supported by FAPESP (2005/52566-1) and CAPES (199-01/2007). Authors wish to thank Dr. Lídia Raquel de Carvalho, IB, UNESP, Campus of Botucatu, for the statistical analysis.

References

- AVITSUR R; HUNZEKER; J, SHERIDAN, J F (2006) Role of early stress in the individual differences in host response to viral infection. *Brain, Behavior, and Immunity* 20: 339-348.
- BANKOVA, V; BOUDOUROVA-KRASTEVA, G; POPOV, S; SFORCIN, J M; FUNARI, S R C (1998) Seasonal variations of the chemical composition of Brazilian propolis. *Apidologie* 29: 361–367.
- BANSKOTA, A H; TEZUKA, Y; KADOTA, S (2001) Recent progress in pharmacological research of propolis. *Phytotherapy Research* 15: 561–571.
- BAZO, A P; RODRIGUES, M A M; SFORCIN, J M; CAMARGO, J L V; RIBEIRO, L R; SALVADORI, D M F (2002) Protective action of propolis on the rat colon carcinogenesis. *Teratogenesis, Carcinogenesis and Mutagenesis* 22: 183-194.
- BOWERS, S L; BILBO, S D; DHABHAR, F S; NELSON, R J (2008) Stressor-specific alterations in corticosterone and immune responses in mice. *Brain, Behavior, and Immunity* 22: 105-113.
- BÚFALO, M C; FIGUEIREDO, A S; DE SOUSA, J P B; CANDEIAS, J M G; BASTOS, J K; SFORCIN, J M. Antiviral activity of *Baccharis dracunculifolia* and propolis and poliovirus quantification by real-time PCR. *Antiviral Research*, submitted.

- CORTHAY, A; SKOVSETH, D K; LUNDIN, K U; ROSJO, E; OMHOLT, H; HOFGAARD, P O; HARALDSEN, G; BOGEN, B (2005) Primary antitumor immune response mediated by CD4⁺ T cells. *Immunity* 22: 371-383.
- DE CASTRO, S L (2001) Propolis: biological and pharmacological activities. Therapeutic uses of this bee-product. *Annual Review of Biomedical Sciences* 3: 49-83.
- DHABHAR, F S (2002) Stress-induced augmentation of immune function – the role of stress hormones, leukocyte trafficking, and cytokines. *Brain, Behavior, and Immunity* 16: 785-798.
- DOMINGUEZ-GERPE, L; REY-MÉNDEZ, M (2001) Alterations induced by chronic stress in lymphocyte subsets of blood and primary and secondary immune organs of mice. *BMC Immunology* 2: 7.
- FANG, L; LONSDORF, A S; HWANG, S T (2008) Immunotherapy for advanced melanoma. *Journal of Investigative Dermatology* 128: 2596-2605.
- FREITAS, S F; SHINOHARA, L; SFORCIN, J M; GUIMARÃES, S (2006) *In vitro* effects of propolis on *Giardia duodenalis* trophozoites. *Phytomedicine* 13: 170-175.
- KAMIMURA, D; ISHIHARA, K; HIRANO, T (2003) IL-6 signal transduction and its physiological roles: the signal orchestration model. *Reviews of Physiology, Biochemistry & Pharmacology* 149: 1-38.
- KIOUKIA-FOUGIA, N; ANTONIOU, K; BEKRIS, S; LIAPI, C; CHRISTOFIDIS, I; PAPADOUPOLOU-DAIFOTI, Z (2002) The effects of stress exposure on the hypothalamic-pituitary-adrenal axis, thymus, thyroid hormones and glucose levels. *Progress in Neuropsychopharmacology and Biological Psychiatry* 26: 823-830.
- MAES, M; SONG, C; LIN, A; JONGH, R D; GASTEL, A V; KENIS, G; BOSMANS, E; MEESTER, I D; BENOY, I; NEELS, H; DEMEDTS, P; JANCA, A; SCHARPÉ, S; SMITH, R S (1998) The effects of psychosocial stress on humans: increased

- production of pro-inflammatory cytokines and a Th1-like response in stress-induced anxiety. *Cytokine* 10: 313-318.
- McCARTER, M; CLARKE, J; RICHTER, D; WILSON, C (2005) Melanoma skews dendritic cells to facilitate a T helper 2 profile. *Surgery* 138: 321-328.
- MISSIMA, F; PAGLIARONE, A C; ORSATTI, C L; SFORCIN, J M. Propolis effect on Th1/Th2 cytokines' expression and production by melanoma-bearing mice submitted to stress. *Brain, Behavior, and Immunity*, not submitted.
- MISSIMA, F; SFORCIN, J M (2008) Green Brazilian propolis action on macrophages and lymphoid organs of chronically stressed mice. *Evidence-based Complementary and Alternative Medicine* 5: 71-75.
- ORSOLIC, N; SARANOVIC, A B; BASIC, I (2006) Direct and indirect mechanism(s) of antitumour activity of propolis and its polyphenolic compounds. *Planta Medica* 72: 20-27.
- POPE, R M; TSCHOPP, J (2007) The role of interleukin-1 and the inflammasome in gout: implications for therapy. *Arthritis & Rheumatism* 56: 3183-3188.
- SÁ-ROCHA, V M; SÁ-ROCHA, L C; PALERMO-NETO, J (2006) Variations in behavior, innate immunity and host resistance to B16F10 melanoma growth in mice that present social stable hierarchical ranks. *Physiology & Behavior* 88: 108-115.
- SFORCIN, J M (2007) Propolis and the immune system: a review. *Journal of Ethnopharmacology* 113: 1-14.
- SFORCIN, J M; FERNANDES JR., A; LOPES, C A M; BANKOVA, V; FUNARI, S R C (2000) Seasonal effect on Brazilian propolis antibacterial activity. *Journal of Ethnopharmacology* 73: 243-249.

- SFORCIN, J M; KANENO, R; FUNARI, S R C (2002) Absence of seasonal effect on the immunomodulatory action of Brazilian propolis on natural killer activity. *Journal of Venomous Animals and Toxins* 8: 19-29.
- SFORCIN, J M; ORSI, R O; BANKOVA, V (2005) Effects of propolis, some isolated compounds and its source plant on antibody production. *Journal of Ethnopharmacology* 98: 301-305.
- SFORCIN, J M; NUNES, G A; MISSIMA, F; SÁ-NUNES, A; FACCIOLI, L H (2007) Effect of a leukotriene inhibitor (MK886) on nitric oxide and hydrogen peroxide production by macrophages of acutely and chronically stressed mice. *Journal of Pharmacy and Pharmacology* 59:1249-1254.
- SFORCIN, J M; MISSIMA, F; ORSATTI, C; PAGLIARONE, A; KANENO, R (2008) Propolis effect on Th1/Th2 cytokine profile in melanoma-bearing mice submitted to stress. *Scandinavian Journal of Immunology* 68: 216-217.
- SJOGREN, E; LEANDERSON, P; KRISTENSON, M; ERNERUDH, J (2006) Interleukin-6 levels in relation to psychosocial factors: studies on serum, saliva, and in vitro production by blood mononuclear cells. *Brain, Behavior, and Immunity* 20: 270-278.
- TAN, E L; SELVARATNAM, G; KANANATHAN, R; SAM, C K (2006) Quantification of Epstein-Barr virus DNA load, interleukin-6, interleukin-10, transforming growth factor- β 1 and stem cell factor in plasma of patients with nasopharyngeal carcinoma. *BMC Cancer* 6: 227.
- THEJASS, P; KUTTAN, G (2007) Inhibition of endothelial cell differentiation and proinflammatory cytokine production during angiogenesis by allyl isothiocyanate and phenyl isothiocyanate. *Integrative Cancer Therapies* 6: 389-399.
- ZAR, J H (1999) *Biostatistical Analysis*. Prentice Hall; New Jersey, USA. 663 pp.

Graphs

Fig. 1. Corticosterone concentrations (ng/mL). G1: control; G2: stress; G3: propolis; G4: propolis + stress; G5: melanoma; G6: melanoma + stress; G7: melanoma + propolis; G8: melanoma + propolis + stress. Data represent means and standard-deviation of 8 animals. * significantly different from G1 ($p<0.001$).

Fig. 2. IL-1 β production (pg/mL) by spleen cells stimulated with LPS (5 μ g/mL) for 48 h. G1: control; G2: stress; G3: propolis; G4: propolis + stress; G5: melanoma; G6: melanoma + stress; G7: melanoma + propolis; G8: melanoma + propolis + stress. Data represent means and standard-deviation of 8 animals. * significantly different from G1 ($p<0.01$).

Fig. 3. IL-6 production (pg/mL) by spleen cells stimulated or not with LPS (5 μ g/mL) for 48 h. G1: control; G2: stress; G3: propolis; G4: propolis + stress; G5: melanoma; G6: melanoma + stress; G7: melanoma + propolis; G8: melanoma + propolis + stress. Data represent means and standard-deviation of 8 animals. * significantly different from G1 ($p<0.05$); # significantly different from G1 + LPS ($p<0.05$).




Fig. 1.




Fig. 2.




Fig. 3.

Anexo 1

BRAIN, BEHAVIOR, AND IMMUNITY

Guide for Authors

Brain, Behavior, and Immunity, founded in 1987, is the official journal of the Psychoneuroimmunology Research Society (PNIRS). This innovative journal publishes peer-reviewed basic, experimental, and clinical studies dealing with behavioral, neural, endocrine, and immune system interactions in humans and animals. It is an international, interdisciplinary journal devoted to investigation of the physiological systems that integrate behavioral and immunological responses. The journal welcomes original research in neuroscience, immunology, integrative physiology, behavioral biology, psychiatry, psychology, and clinical medicine and is inclusive of research at the molecular, cellular, social, and organismic levels. The journal features online submission and review. Manuscripts are usually peer-reviewed and returned to authors within 30 days of submission, leading to timely publication of experimental results. There are no submission fees or page charges for *Brain, Behavior, and Immunity*, which is published eight times a year. Detailed instructions for authors can be found at <http://ees.elsevier.com/bbi/>.

Research areas include:

- Neurochemical and hormonal mechanisms that convey messages to and from the immune system and brain
- Stress and immunity, including the effects of stress-related hormones and neurotransmitters on the immune system and brain
- Actions of cytokines and growth factors on neuronal and glial cells to regulate behavior, cognition, clinical pain and neuroendocrine function
- Clinical translational research, such as depression, other mental disorders, fatigue, chronic inflammatory diseases
- Roles of hormones, growth factors, and cytokines in the immune and central nervous systems
- Inflammation, neuroscience, and behavior
- Neuroimmunopharmacology and the immunomodulating effects of psychotropic drugs and drugs of abuse
- Sleep, exercise, immunity, and health
- Roles of cytokines, hormones, and neurotransmitters in the aging immune system and brain
- Cancer, brain, and immunity
- Regulation of nerve injury and repair by the immune system
- Psychosocial, behavioral, and neuroendocrine influences on immunity and on the development and progression of immunologically-mediated disease processes
- Genomics of behavior and immunity

Electronic Submission of Manuscripts

Manuscripts must be written in English and submitted electronically at <http://ees.elsevier.com/bbi/>. New contributors should first register at this site and then log into the Elsevier Editorial System (EES) with their user name and password.

There are eight steps that must be completed to submit a manuscript: Enter Article Title; Select Article Type; Add/Edit Remove Author (corresponding author does not need to be the person who submits the paper); Submit Abstract; Enter Key Words; Select Document Classification; Enter Comments (recommend expert reviewers); Attach Files. All sections except the last one can be “copied and pasted” into text boxes from existing files.

The files that must be attached separately are: cover letter to the Editor-in-Chief, manuscript, figures, and tables. An introductory cover letter must outline the most important research findings and their significance. Complete legends (captions) for both figures and tables should be placed at the end of the manuscript. Figures must be attached as separate files or as a single file. Tables must also be attached as either individual tables or a single file with all the tables. All files containing figures or tables must clearly identify each figure or part of figure by adding, at the top of each figure or table, the name of the first author and abbreviated title of the manuscript. Authors can also upload supplementary material such as video, audio, movie and other files (which will be available as a link in the PDF file that the system generates). After the files are attached, the EES system will create a PDF file, which may require a few minutes. You will then be asked to approve the PDF file, a step that must be completed before the new submission is sent to the Editor-in-Chief who will initiate the review process.

The editorial office of *Brain, Behavior, and Immunity* is:

Dr. Keith W. Kelley

Editor-in-Chief, Brain, Behavior, and Immunity

University of Illinois at Urbana-Champaign

Laboratory of Immunophysiology

Integrative Immunology and Behavior Program

Department of Animal Sciences

212 Edward R. Madigan Laboratory

1201 West Gregory Drive

Urbana, IL 61801, USA

bbieditor@illinois.edu

Telephone: (217) 333-9641

Fax: (217) 244-5617

<http://ees.elsevier.com/bbi/>

The authors should provide names and e-mail addresses of **at least three** qualified, expert reviewers.

Ethics

Manuscripts are accepted for review with the understanding that the same work has not been and will not be published elsewhere (except in abstract form) and that it is not being submitted for review elsewhere. It is also understood that all persons listed as authors have given their approval for the submission of the paper and that any person cited as a source of personal communications has approved such citation. Written authorization may be required at the Editor's discretion. Articles and any other material published in *Brain, Behavior, and Immunity* represent the opinions of the author(s) and should not be construed to reflect the opinions of the Editor(s) or the Publisher.

Copyright

Upon acceptance of an article, authors will be asked to sign a “Journal Publishing Agreement” (for more information on this and copyright, see <http://www.elsevier.com>). Acceptance of this agreement will ensure the widest possible dissemination of information. An e-mail (or letter) will be sent to the corresponding author confirming receipt of the manuscript, together with a “Journal Publishing Agreement” form or a link to the online version of this agreement.

If material from other copyrighted works is included, the author(s) must obtain written permission from the copyright owners and credit the source(s) in the article. Elsevier has preprinted forms for use by authors in these cases: contact Elsevier Global Rights Department, P.O. Box 800, Oxford OX5 1DX, UK; phone: (+44) 1865 843830, fax: (+44) 1865 853333, e-mail: permissions@elsevier.com. Requests may also be completed online via the Elsevier home page (<http://www.elsevier.com/locate/permissions>). Authors are responsible for retaining the signed permission forms following approval by the copyright owners.

Manuscripts Using Human Subjects

Brain, Behavior, and Immunity requires the use of guidelines known as “Consolidated Standards of Reporting Trials” (CONSORT) for publishing results of clinical trials. Information about these guidelines can be found at <http://www.consort-statement.org/>. Authors should note that CONSORT requires a checklist that should be submitted along with the cover letter. Inclusion of a figure depicting subject flow and allocation to the intervention and control conditions must also be contained within the manuscript.

Types of Articles

The chief criteria for the acceptance of submitted papers are the quality, originality, and clarity of the work reported. Original full-length research reports, full-length review articles, brief commentaries, and letters to the editor will be considered for publication. The journal also publishes invited minireviews and short communications. The Editor-in-Chief invites timely articles to be published as minireviews. These minireviews are aimed at being short and focused on a contemporary topic. Invited minireviews consist of approximately 6,000 words of text and 25-30 scientific references. Minireviews must contain one figure highlighting the theme of the article, complete with an explanatory figure legend. If appropriate, a color version of the figure can be published in the online publication, with a black-and-white figure in the print version. If the author chooses this option, the figure legend must be self-explanatory in the absence of color-coding. Manuscripts published as short communications are, primarily, reports of noteworthy findings that do not require extensive introduction or discussion. They may include findings of special interest, some kinds of preliminary results, or descriptions of methods, techniques, or reanalysis of data. Short communications are subject to peer review. Papers will be considered short communications if the *text, references, and a maximum of two tables or figures (or one of each) are limited to 3,500 words*. Authors may elect to include additional illustrations, but the limitation to 3,500 words will remain. Letters to the editor should be of high scientific quality, contain less than 500 words, and cite no more than 5 scientific references. If the letter is directed to a paper

published in *Brain, Behavior, and Immunity*, the author of that paper will be provided an opportunity to respond. Both the letter to the editor and the author's response will be published simultaneously.

Announcements

Brain, Behavior, and Immunity will consider for publication announcements of interest to the readership such as notices of scientific meetings.

Language Services

Authors who require information about language editing and copyediting services pre- and post-submission please visit <http://www.elsevier.com/locate/languagepolishing> or contact

authorsupport@elsevier.com for more information. Please note Elsevier neither endorses nor takes responsibility for any products, goods or services offered by outside vendors through our services or in any advertising. For more information please refer to our terms and Conditions <http://www.elsevier.com/termsandconditions>

Form

Manuscripts should be prepared using a 12-point font, double-spaced throughout (including tables, footnotes, references, and figure captions) with 1-in. margins on all sides. Unusual typeface is acceptable only if it is clear and legible. For initial submission, all manuscripts must be prepared and submitted in one of the following formats: Microsoft Word (.doc), WordPerfect (.wps), or Rich Text Format (.rtf). All figures and tables should be clearly labeled at the top.

Revised manuscripts should not be marked using underlined or bolded words to indicate changes from the original submission. Instead, changes in the revised manuscript must be explained in a rebuttal letter. Submission of all revised manuscripts requires both figures and tables to be submitted separately from the manuscript text: do not insert figures and tables at the end of the text for revised manuscripts. Instead, the electronic submission system requires identification and submission of figures and tables separate from the text of revised manuscripts (see information below for graphs, scans, and illustrations). For more information, please also see the Author Gateway Web page for *Brain, Behavior, and Immunity* available through the journal home page at <http://www.elsevier.com/locate/ybrbj>.

Page 1 should contain the article title, the names and affiliations of all authors, and, at the bottom of the page, the name, mailing address, e-mail address, telephone number, and Fax number of the person to whom correspondence, proofs, and reprint requests should be sent.

Page 2 should contain an abstract of the article that is not to exceed 250 words. A list of up to 10 **keywords** or phrases suitable for indexing should be provided in the electronic submission process.

Headings. The organization of the paper should be clearly indicated by appropriate headings and subheadings. After an introductory statement of the background and problem to be investigated, most manuscripts would be arranged under the following

main headings: Methods, Results, Discussion, and References.

Abbreviations. Do not use periods after abbreviations of measure (cm, s, kg, mA, etc.) in text or tables, except for "in." (inch). The American Chemical Society *Style Guide* should be used as a reference for proper abbreviations.

Statistical analyses. Analysis of variance tables should not be submitted, but important *F* ratios should be presented in the text. Letters used as statistical symbols should be typed in italics. Common statistical tests need not be referenced. The appropriate form for presentation of statistical information should include degrees of freedom and *p* value (e.g., $F(3,17) = 4.16, p < .05$; $t(22) = 2.97, p < .01$; $r(2,15) = 0.58, p < .05$).

Footnotes. Footnotes should be used only when absolutely necessary. They should be identified in the text by consecutive superscript Arabic numerals and typed double-spaced on a separate page.

References. The name and date system should be used for citing all references. All citations in the text should refer to:

1. *Single author:* the author's name (without initials, unless there is ambiguity) and the year of publication.
2. *Two authors:* both authors' names and the year of publication.
3. *Three or more authors:* first author's name followed by et al. and the year of publication.

Examples: "as demonstrated (Allan, 1996a, 1996b, 1999; Allan and Jones, 1995).
Kramer et al. (2000) have recently shown"

References should be arranged first alphabetically and then further sorted chronologically if necessary. More than one reference from the same author(s) in the same year must be identified by the letters "a", "b", "c", etc., placed after the year of publication.

For journal names follow the abbreviations in the most recent edition of the *Chemical Abstracts Service Source Index*.

Note the following examples:

Antoni, M., 2002. Stress Management Intervention for Women with Breast Cancer. American Psychological Association, Washington, DC.

French, R.A., Zachary, J.F., Dantzer, R., Frawley, L.S., Chizzonite, R., Parnet, P., Kelley, K.W., 1996. Dual expression of p80 type I and p68 type II interleukin-1 receptors on anterior pituitary cells synthesizing growth hormone. *Endocrinology* 137, 4027-4036.

Kiecolt-Glaser, J.K., 1999. Stress, personal relationships, and immune function: health implications. *Brain Behav. Immun.* 13, 61-72.

SAS Institute, 1988. SAS/STAT User's Guide. SAS Institute, Cary, NC.

Sluzki, C.E., Beavin, J., 1977. Symmetry and complementarity. In: Watzlawick, P., Weakland, J.H. (Eds.), *The Interactional View*. Norton, New York, pp. 71-87. Reprinted from: *Acta Psiquiatrica y Psicologica America Latin* 11, 321-330 (1965) (reprint).

Turner-Cobb, J., Sephton, S., Spiegel, D., 2001. Psychosocial effects on immune function and disease progression in cancer: human studies. In: Ader, R., Felten, D., Cohen, N. (Eds.), *Psychoneuroimmunology*. Academic Press, San Diego, pp. 565-582.

World Health Organization, 1998. Obesity: Preventing and managing the global epidemic: Report of a WHO Consultation on Obesity, Geneva, June 3-5, 1997. World Health Organization, Geneva.

Tables. Number tables consecutively with Arabic numerals in order of appearance in the text. Tables should be double-spaced on separate pages that follow the References. The title and footnotes (typed beneath the table) should contain all the information necessary to understand and interpret the table without reference to the text. Footnotes to tables should be identified by superscript lowercase letters and assigned alphabetically in order of appearance as the table is read *horizontally*.

Graphs. Graphs and charts should be in a finished form suitable for high-quality journal reproduction. Units of measurement should be given in the ordinate and abscissa. Graphs and charts can be submitted as Powerpoint (.ppt), Tagged-Image File Format (.tiff or .tif), or Encapsulated PostScript (.eps) files. If graphs are submitted in color, they must be saved in the CMYK (Cyan, Magenta, Yellow, Black) mode for color. If a .tiff (.tif) format is being used and the file is greater than 10 MB, file size should be reduced by using LZW compression. Following preparation of graphs, scans, and illustrations (see below), all figures should be numbered consecutively with Arabic numerals in order of appearance in the text, including figures with more than one part (i.e., Fig. 1A, Fig. 1B, Fig. 1C). All images should then be inserted at the end of the manuscript (one illustration per page). Each graph must be supplied as a separate file in a .ppt, .tif, or .eps format. For further information on the preparation of electronic artwork, please see ➡ <http://www.elsevier.com/artworkinstructions>. The cost of print color reproduction will be charged to the author(s).

Scans and illustrations. Scans should be prepared using high-quality image editing software such as Adobe Photoshop or Corel Photo-Paint. Illustrations should be constructed with high-resolution software such as Adobe Illustrator or CorelDRAW. All images should be in a finished form suitable for publication. Scans and drawings may be either 87 mm (one column or one-half page) or 178 mm (two column or full page) wide. Both widths require resolutions yielding 600 dots per inch/pixels per inch (dpi/ppi) or greater. If scans or illustrations are to be printed in color, it is important that the images be saved in the CMYK mode for color. Scans and illustrations can be submitted either as .tiff (.tif) or .eps files. If a .tiff (.tif) format is being used and the file is greater than 10 MB, file size should be reduced by using LZW compression. All images should be numbered consecutively with Arabic numerals in order of appearance in the text, including figures with more than one part (i.e., Fig. 1A, Fig. 1B, Fig. 1C). Each high-resolution scan or illustration must be supplied as an individual source file in either a .tiff or .eps format.

Color on the Web. The cost of print **color** reproduction will be charged to the author(s). However, if together with your accepted article, you submit usable color figures, then Elsevier will ensure, *at no additional charge*, that these figures will appear in color on the Web (e.g., ScienceDirect and other sites) regardless of whether these illustrations are reproduced in color in the printed version. For color reproduction in print, you will receive information regarding the costs from Elsevier after receipt of your accepted article. For further information on the preparation of electronic artwork, please see <http://www.elsevier.com/artworkinstructions>.

Please note: Because of technical complications that can arise in converting color figures to "gray scale" (for the printed version should you not opt for color in print), please submit in addition usable black-and-white electronic figures corresponding to all the color illustrations, using the criteria set forth above.

GenBank/DNA sequence linking. Authors wishing to enable other scientists to use the accession numbers cited in their papers via links to these sources should type this information in the following manner:

For each and every accession number cited in an article, authors should type the accession number in **bold, underlined text**. Letters in the accession number should always be capitalized (*see example below*). This combination of letters and format will enable Elsevier's typesetters to recognize the relevant texts as accession numbers and add the required link to GenBank's sequences.

Example: GenBank accession nos. **AI631510**, **AI631511**, **AI632198**, and **BF223228**), a B-cell tumor from a chronic lymphatic leukemia (GenBank accession no. **BE675048**), and a T-cell lymphoma (GenBank accession no. **AA361117**).

Authors are encouraged to check accession numbers used very carefully. **An error in a letter or number can result in a dead link.**

In the final version of the **printed article**, the accession number text will not appear bold or underlined. In the final version of the *electronic* copy, the accession number text will be linked to the appropriate source in the NCBI databases, enabling readers to go directly to that source from the article.

Preparation of supplementary material

Elsevier now accepts electronic supplementary material to support and enhance your scientific research. Supplementary files offer additional possibilities for publishing supporting applications, movies, animation sequences, high-resolution images, background datasets, sound clips, and more. Supplementary files supplied will be published online alongside the electronic version of your article in Elsevier Web products, including ScienceDirect (<http://www.sciencedirect.com>). To ensure that your submitted material is directly usable, please provide the data in one of our recommended file formats. Authors should submit the material in electronic format together with the article and supply a concise and descriptive caption for each file. Please note, however, that supplementary material will not appear in the printed journal. For more detailed instructions, please visit <http://www.elsevier.com/artworkinstructions>.

Special Subject Repositories

As a service to our authors, Elsevier will deposit to PubMed Central (PMC) author manuscripts on behalf of Elsevier authors reporting NIH funded research. This service is a continuation of Elsevier's 2005 agreement with the NIH when the NIH introduced their voluntary 'Public Access Policy.'

The service will help authors comply with the National Institutes of Health (NIH) revised "Public Access Policy," effective April 7, 2008. The NIH's revised policy requires that NIH-funded authors submit to PubMed Central (PMC), or have submitted on their behalf, their peer-reviewed author manuscripts, to appear on PMC no later than 12 months after final publication.

Elsevier will send to PMC the final peer-reviewed manuscript, which was accepted for publication and sent to Elsevier's production department, and that reflects any author-agreed changes made in response to peer-review comments. Elsevier will authorize the author manuscript's public access posting 12 months after final publication. Following the deposit by Elsevier, authors will receive further communications from Elsevier and NIH with respect to the submission.

Authors are also welcome to post their accepted author manuscript on their personal or institutional web site. Please note that consistent with Elsevier's author agreement, authors should not post manuscripts directly to PMC or other third party sites. Individual modifications to this general policy may apply to some Elsevier journals and society publishing partners.

As a leading publisher of scientific, technical and medical (STM) journals, Elsevier has led the industry in developing tools, programs and partnerships that provide greater access to, and understanding of, the vast global body of STM information. This service is an example of Elsevier willingness to work cooperatively to meet the needs of all participants in the STM publishing community.

Proofs

One set of page proofs in PDF format will be sent by e-mail to the corresponding Author (if we do not have an e-mail address then paper proofs will be sent by post). Elsevier now sends PDF proofs which can be annotated; for this you will need to download Adobe Reader version 7 available free from <http://www.adobe.com/products/acrobat/readstep2.html>. Instructions on how to annotate PDF files will accompany the proofs. The exact system requirements are given at the Adobe site: <http://www.adobe.com/products/acrobat/acrrsystemreqs.html#70win>

If you do not wish to use the PDF annotations function, you may list the corrections (including replies to the Query Form) and return to Elsevier in an e-mail. Please list your corrections quoting line number. If, for any reason, this is not possible, then mark the corrections and any other comments (including replies to the Query Form) on a printout of your proof and return by fax, or scan the pages and e-mail or send by post.

Please use this proof only for checking the typesetting, editing completeness and

correctness of the text, tables and figures. Significant changes to the article as accepted for publication will only be considered at this stage with permission from the Editor. We will do everything possible to get your article published quickly and accurately. Therefore, it is important to ensure that all of your corrections are sent back to us in one communication: please check carefully before replying, as inclusion of any subsequent corrections cannot be guaranteed. Proofreading is solely your responsibility. **Return corrections within 2 days of receipt of the proofs.** Note that Elsevier may proceed with the publication of your article if no response is received.

Offprints

The corresponding author, at no cost, will be provided with a PDF file of the article via e-mail or, alternatively, 25 free paper offprints. The PDF file is a watermarked version of the published article and includes a cover sheet with the journal cover image and a disclaimer outlining the terms and conditions of use.

Additional paper offprints can be ordered by the authors. An order form with prices will be sent to the corresponding author.

Author Enquiries

For enquiries relating to the submission of articles please visit this journal's homepage at ➡ <http://www.elsevier.com/locate/ybrbi>. You can track accepted articles at ➡ <http://www.elsevier.com/trackarticle> and set up e-mail alerts to inform you of when an article's status has changed, as well as copyright information, frequently asked questions and more.

Contact details for questions arising after acceptance of an article, especially those relating to proofs, are provided after registration of an article for publication.

Disclaimer

Whilst every effort is made by the publishers and editorial board to see that no inaccurate or misleading data, opinion or statement appears in this journal, they wish to make it clear that the data and opinions appearing in the articles and advertisements herein are the sole responsibility of the contributor or advertiser concerned. Accordingly, the publishers, the editorial board and editors and their respective employees, officers and agents accept no responsibility or liability whatsoever for the consequences of any inaccurate or misleading data, opinion or statement.

Anexo 2

Journal of ApiProduct and ApiMedical Science

JAAS

The international link between bee product science and practice

Guidelines for authors © 2008 IBRA

16, NORTH ROAD,
CARDIFF, CF10 3DY
UNITED KINGDOM
TEL: +44(0)2920 372409
FAX: +44(0)5601 135640
joneshr@ibra.org.uk
Richard Jones
Director

Those considering writing for the journal should study the following guidelines. Manuscripts that are correctly formatted will be processed much more quickly than those requiring considerable copy-editing, and submissions deviating significantly from these guidelines will not be considered by the editorial team.

Ethical considerations will be taken into account in considering the acceptability of papers, and the editorial board's decision on this, as on other aspects, will be final. When submitting a manuscript, the corresponding author should confirm that all the authors have read and approved of the material being submitted, and that the findings reported are their own work. Papers submitted or data contained therein must not have been published or accepted for publication in any other journal, or be under consideration for publication elsewhere.

Scope

The *Journal of ApiProduct and ApiMedical Science* publishes original research articles, original theoretical papers, notes, comments and authoritative reviews. The scope of the journal is two fold. On one hand it covers the relevant properties or substances of the six hive products honey, pollen, propolis, wax, royal jelly, and bee venom. A special attention is given to the biological and health relevant properties. On the other hand it will cover studies concerning biological and medical effects of the bee products in experiments with animal and humans.

Editing and peer review

The journal is committed to the rapid processing and quality reviewing of manuscripts. The editors handle the scientific assessment of papers, including their initial review, allocation of referees and the subsequent assessment of review and authors' revisions. IBRA undertakes further editing of papers once accepted, to eliminate any final problems and ensure they conform to the journal's style.

Access

Open Access means that on payment of an up-front fee by the authors, articles will be made available online free of charge to anybody. Authors will retain copyright of their article, and will then be able to grant anybody the right to disseminate and reproduce it, provided that it is cited correctly and that no errors are introduced. As a special introductory offer for articles submitted to the journal during 2009, the cost for this service will be £375 (€500 US\$725) per article, to be payable on acceptance. From 2010

onwards the cost will be £650 (€950 US\$1500). Increasingly, grant awarding bodies are supporting, or even requiring, publication of research that they have funded under Open Access arrangements. Open Access is of course merely an additional option, and normal publication in the journal will remain free of page charges.

Structure of papers

Original Research Articles

Original research papers should be organized in the following way: Title, Author(s), Author affiliation, Summary, Keywords (provide up to 10), Introduction (including the objectives of the work), Materials and methods, Results, Discussion, Acknowledgements, References, and a short title (running head). The Results and Discussion should not be combined. Theoretical papers should contain some of these elements although they may require modified organization.

Notes and Comments

This section is intended as a forum for two kinds of paper:

- Brief research notes, reporting techniques and findings that are too narrow in scope to justify a larger research report.
- Constructive criticisms of research papers that have recently appeared in the journal or elsewhere.

Papers intended for this section should not normally exceed 1000 words, and should contain no more than one figure or table, and no more than five references. They should contain keywords (up to 10), but no summary; and should not have the subheadings of Introduction, Materials and Methods, etc. All papers published in this section will be subject to the same review process as Original Research Articles. Where it seems appropriate, the editors may solicit comments on an accepted paper from scientists known to have a strong interest in the subject. Comments contributed as a result of invitations from the editors will be evaluated by the journal's standard review process. Upon the recommendation of the referees, authors of Original Research Articles may be invited to restructure their papers to fit the comment format.

Papers received this way may not require additional review.

Reviews

These are authoritative reviews of specific subjects written by experts who have helped advance the subjects they are reviewing. They should cover a clearly delimited topic in bee biology or the economics, technology, sociology or history of beekeeping. An important function of a review is to give an appraisal of the relative significance of different work, and to point out which is the most important. Reviews are accompanied by a comprehensive bibliography giving references to the most important papers that have been published within the period under review; this is a vital component of the review and extends its usefulness considerably.

Writing style

Papers should be in clear, concise English. Preferably write in the first person and avoid the use of impersonal phrases such as 'the present authors'. The full scientific (Latin) name for the species must be cited in full the first time any organism is mentioned. In subsequent use, the genus should be abbreviated to its initial (e.g. *B. hortorum*).

Authorities for specific names are not required except for taxonomic papers. Sufficient information on procedures must be supplied to make it possible for experiments to be repeated elsewhere. Statistical evaluation of results must be given where appropriate. The sources of equations and the names of statistical tests must be given in full at their first citation. An arithmetic mean should preferably be stated in the form: mean $\pm$ s.e. (n). The metric system (SI) must be used for all weights and measures, and temperatures must be given in $^{\circ}\text{C}$. Abbreviations for units must conform to usage in the journal and can be given in the following ways: '6.5–10.9 $^{\circ}\text{C}$ ' and 'from 6.5 to 10.9 $^{\circ}\text{C}$ ' or '14.2 $\pm$ s.e. 0.5 $^{\circ}\text{C}$ ($n = 50$)'. Durations of time should be given as '3 h 30 min' or '210 min'. Times of day should be given in the format '14.15 h'. Authors for whom English is not their primary language are encouraged to seek professional editorial assistance in the preparation of manuscripts. The editors reserve the right to reject, without further review, papers which do not meet acceptable standards of English.

References

References should conform to the Harvard (name and date) system. When quoting references in the text, give the name of the author(s) and the year of publication, and if there are more than two authors give the name of the first followed by '*et al.*' References cited together in the text should be arranged chronologically, for example: (Goodwin, 1985; Anderson, 1986, 1987).

Arrange the list of references at the end of the paper alphabetically by author's name in the following order: publications of the single author, chronologically ordered; publications of the same author with one or more than one co-authors, chronologically ordered. It is the author's responsibility to check that the references in the text correspond exactly with the list at the end of the paper. The style and punctuation of references listed should follow the format given here. Titles of periodicals are not to be abbreviated. Authors should consult recent issues for further examples.

IBRA is committed to implementing the DOI (Digital Object Identifier) system throughout its journal output. **It is the responsibility of authors to ensure that references to articles bearing DOI numbers include the correct number.**

Article: BENDA, N D; BOUCIAS, D; TORTO, B; TEAL, P (2008) Detection and Characterisation of *Kodamaea ohmeri* associated with small hive beetle *Aethina tumida* infesting honey bee hives. *Journal of Apicultural Research* 47(3): 194-201. DOI: 10.3827/IBRA.1.47.3.07

Book: ROUBIK, D W (1989) *Ecology and natural history of tropical bees*. Cambridge University Press; Cambridge, UK. 514 pp.

Conference proceedings: DE GUZMAN, L; FRAKE, A (2007) Russian honey bees and their response towards brood infested with small hive beetles. In *Proceedings of 40th International Apicultural Congress, Melbourne, Australia, 9-14 September 2007*. p185.

Article in a book: MAURIZIO, A (1979) Microscopy of honey. In Crane, E (ed). *Honey, a comprehensive survey (2nd edition)*. Heinemann; London, UK. pp. 240–257.

Illustrations and graphics

When you first e-mail your paper for consideration by the editors, you should send a single Microsoft Word file containing the text **followed** by any figures and tables. **Do not include tables or figures within the text.** Once the paper has been accepted, you will need to send the text and image files formatted as described below.

Please read these instructions carefully to avoid delays in processing your paper. Colour photos or illustrations can be included. Figures should be numbered serially, and a separate list provided with the titles of each typed in double-spacing. Titles should enable the figures to be understood without reference to the text. Micrographs must have calibration lines in the bottom right hand corner, even if magnification factors (such as '×1000') are given, as they may be resized.

Authors who use applications such as Microsoft Excel to create graphs and illustrations can submit the whole spreadsheet with the graph embedded within it. Authors who do this will be able to take advantage of the high-resolution image-setter used by the journal's designers. **Please do not use colours or patterned fills within charts or graphs, and instead use shades of grey to differentiate items.**

Photographs should be no less than 180 pixels per cm/225 pixels per inch. Files should be supplied as 8 bit greyscale (or 256 greys), and saved as a TIFF or maximum quality JPEG. If using a digital camera, files should be supplied at a minimum resolution of three megapixels at the highest quality JPEG setting possible. Colour images should conform to the same resolution as greyscale images, and can be sent as TIFF or JPEG files.

Tables

Tables should be self-explanatory, and headings should be self-contained. The reader should not have to search the text for relevant material. Footnotes are for incidental information (e.g. *observation delayed by bad weather; **two samples only). Make sure that the type for all headings can be set within the width of a page. A table can be printed sideways if absolutely necessary, though this should be avoided if possible. Authors are encouraged to use the tables feature in Microsoft Word or similar programs and ensure that every value is assigned a unique row and column. **Tables made manually with tabs or spaces frequently scramble values when viewed on a different computer.**

Submitting manuscripts

E-mail is the preferred method of submission. Printed copy sent by mail will only be considered with prior agreement of the editors.

When you first e-mail your paper for consideration by the editors, you should send a single Microsoft Word file containing the text **followed** by any figures and tables. **Do not include tables or figures within the text. Please keep formatting to a minimum and do not include electronic footnotes etc.** Once the paper has been accepted, you will need to send the text and image files separately as described in the 'Illustrations and Graphics' section of these guidelines (see above).

Please e-mail original research articles, notes and comments or reviews to jaas@ibra.org.uk

Editorial Board:

Name and contact information Main Field of Competence

Dr. Stefan Bogdanov

Bee Product Science

Moosstrasse 7

3127 Muehlethurnen, Switzerland

Info@bee-hexagon.net

Bee Product Science

Dr. Noori Al-Waili

Al Waili Science and Trading

New York, USA

noori786@yahoo.com

Api Medicine

Prof. Vassya Bankova

Institute of Organic Chemistry

Acad. Bonchev bl. 9

1113 Sofia, Bulgaria

bankova@orgchm.bas.bg

Propolis Science

Prof. Maria Campos

Laboratory of Pharmacognosy

Fac. Coimbra, Universidade de Coimbra

PT-3000-295 Coimbra Portugal

mgcampos@ff.uc.pt

Pollen Science

Prof. Rose Cooper,

Centre Biomedical Sciences, School of Applied Sciences,

University of Wales Institute, Cardiff, Wales, UK

RCooper@uwic.ac.uk

Honey Microbiology and

Biological Research

Dr. Mitsuo Matsuka

Tamagawa University,

Research Institute Machida-shi,

JP-194-8610

Tokyo, Japan

mmat@agr.tamagawa.ac.jp

Propolis and Royal Jelly

Science

Dr. Cristina Matteescu

Institute for Beekeeping Research

42 Ficusului Blv. Sector 1

RO-013975 Bucharest

cristina.mateescu@icdapicultura.com

Applied Apitherapy

Dr. Keith Morris

Centre Biomedical Sciences, School of Applied Sciences,

University of Wales Institute, Cardiff, Wales, UK

kmorris@uwic.ac.uk

Haematology and Statistics

Prof. Karsten Muenstedt

University clinic Giessen

Klinikstrasse 32

35392 Giessen – Germany

Karsten.Muenstedt@gyn.med.uni-giessen.de

Api Medical Research

Prof. Nada Orsolic

Univ Zagreb,

Fac Sci, Dept Anim Physiol,
Zagreb 10000, Croatia
norsolic@yahoo.com
Api Medical Research

Once you have received the editor's letter confirming that your paper has been accepted in its final form, e-mail a copy of the revised manuscript and accompanying illustrations and tables to the editor dealing with your paper as separate files:-

- File 1: the text of the article in Microsoft Word format
- File 2: any tables
- File 3: the captions to the figures
- Any images or graphics, sent each as a separate file (see above for details of file format)

Please ensure that the files have been scanned using the latest virus scanning software with an up-to-date virus definitions file.

Proofs and reprints

Proofs will be sent to the first author indicated on the title page unless otherwise specified. To speed up the proofing process, the proofs will be e-mailed as an Acrobat portable document format (.pdf) file, unless hard copy was specifically requested at the time of acceptance (you must inform IBRA of this as soon as the paper is accepted). To read the file, you will need the Adobe Acrobat Reader that is free to download from www.adobe.com/products/acrobat/readstep2.html

The proofs should be corrected carefully and must be returned to IBRA within 48 hours of receiving them. If this period is exceeded, the proofs will be checked only by the editorial staff of the journal and printed without the author's corrections. If substantial changes from the original manuscript are requested at the proof stage, they will be subject to editorial review and made at the author's expense.

A high resolution pdf file will be sent free of charge to the first author or corresponding author for each paper.

Copyright statement

The copyright to articles is transferred to IBRA (for US government employees, to the extent transferable) effective when the article is accepted for publication. The copyright transfer covers the exclusive right to reproduce and distribute the article, including reprints, translations, photographic reproductions, microform, electronic form (offline, online) or any other reproductions of similar nature.

As an additional service to authors, IBRA allows authors to post a pdf file of their article on their personal websites. An acknowledgement to IBRA should be included: 'Provided with permission of IBRA, the International Bee Research Association (www.ibra.org.uk).' If possible, please link the IBRA URL to the IBRA website.

All communications on proofs and reprints should be mailed to
Ms. S L Jones, Assistant Editor, JAAS, jonessl@ibra.org.uk
©IBRA: 06/11/2008