

First low frequency all-sky search for continuous gravitational wave signals

J. Aasi,¹ B. P. Abbott,¹ R. Abbott,¹ T. D. Abbott,² M. R. Abernathy,¹ F. Acernese,^{3,4} K. Ackley,⁵ C. Adams,⁶ T. Adams,^{7,8} P. Addesso,⁹ R. X. Adhikari,¹ V. B. Adya,¹⁰ C. Affeldt,¹⁰ M. Agathos,¹¹ K. Agatsuma,¹¹ N. Aggarwal,¹² O. D. Aguiar,¹³ A. Ain,¹⁴ P. Ajith,¹⁵ B. Allen,^{10,16,17} A. Allocca,^{18,19} D. V. Amariutei,⁵ M. Andersen,²⁰ S. B. Anderson,¹ W. G. Anderson,¹⁶ K. Arai,¹ M. C. Araya,¹ C. C. Arceneaux,²¹ J. S. Areeda,²² N. Arnaud,²³ G. Ashton,²⁴ S. M. Aston,⁶ P. Astone,²⁵ P. Aufmuth,¹⁷ C. Aulbert,¹⁰ S. Babak,²⁶ P. T. Baker,²⁷ F. Baldaccini,^{28,29} G. Ballardini,³⁰ S. W. Ballmer,³¹ J. C. Barayoga,¹ S. E. Barclay,³² B. C. Barish,¹ D. Barker,³³ F. Barone,^{3,4} B. Barr,³² L. Barsotti,¹² M. Barsuglia,³⁴ J. Bartlett,³³ M. A. Barton,³³ I. Bartos,³⁵ R. Bassiri,²⁰ A. Basti,^{36,19} J. C. Batch,³³ C. Baune,¹⁰ V. Bavigadda,³⁰ B. Behnke,²⁶ M. Bejger,³⁷ C. Belczynski,³⁸ A. S. Bell,³² B. K. Berger,¹ J. Bergman,³³ G. Bergmann,¹⁰ C. P. L. Berry,³⁹ D. Bersanetti,^{40,41} A. Bertolini,¹¹ J. Betzwieser,⁶ S. Bhagwat,³¹ R. Bhandare,⁴² I. A. Bilenko,⁴³ G. Billingsley,¹ J. Birch,⁶ R. Birney,⁴⁴ S. Biscans,¹² M. Bitossi,³⁰ C. Biwer,³¹ M. A. Bizouard,²³ J. K. Blackburn,¹ C. D. Blair,⁴⁵ D. Blair,⁴⁵ S. Bloemen,^{11,46} O. Bock,¹⁰ T. P. Bodiya,¹² M. Boer,⁴⁷ G. Bogaert,⁴⁷ P. Bojtos,⁴⁸ C. Bond,³⁹ F. Bondu,⁴⁹ R. Bonnand,⁸ R. Bork,¹ M. Born,¹⁰ V. Boschi,^{19,36} Sukanta Bose,^{14,50} C. Bradaschia,¹⁹ P. R. Brady,¹⁶ V. B. Braginsky,⁴³ M. Branchesi,^{51,52} V. Branco,⁵³ J. E. Brau,⁵⁴ T. Briant,⁵⁵ A. Brillet,⁴⁷ M. Brinkmann,¹⁰ V. Brisson,²³ P. Brockill,¹⁶ A. F. Brooks,¹ D. A. Brown,³¹ D. Brown,⁵ D. D. Brown,³⁹ N. M. Brown,¹² C. C. Buchanan,² A. Buikema,¹² T. Bulik,³⁸ H. J. Bulten,^{56,11} A. Buonanno,^{57,26} D. Buskulic,⁸ C. Buy,³⁴ R. L. Byer,²⁰ L. Cadonati,⁵⁸ G. Cagnoli,⁵⁹ J. Calderón Bustillo,⁶⁰ E. Calloni,^{61,4} J. B. Camp,⁶² K. C. Cannon,⁶³ J. Cao,⁶⁴ C. D. Capano,¹⁰ E. Capocasa,³⁴ F. Carbognani,³⁰ S. Caride,⁶⁵ J. Casanueva Diaz,²³ C. Casentini,^{66,67} S. Caudill,¹⁶ M. Cavaglià,²¹ F. Cavalier,²³ R. Cavalieri,³⁰ C. Celerier,²⁰ G. Cella,¹⁹ C. Cepeda,¹ L. Cerboni Baiardi,^{51,52} G. Cerretani,^{36,19} E. Cesarini,^{66,67} R. Chakraborty,¹ T. Chalermsoongsak,¹ S. J. Chamberlin,¹⁶ S. Chao,⁶⁸ P. Charlton,⁶⁹ E. Chassande-Mottin,³⁴ X. Chen,^{55,45} Y. Chen,⁷⁰ C. Cheng,⁶⁸ A. Chincarini,⁴¹ A. Chiummo,³⁰ H. S. Cho,⁷¹ M. Cho,⁵⁷ J. H. Chow,⁷² N. Christensen,⁷³ Q. Chu,⁴⁵ S. Chua,⁵⁵ S. Chung,⁴⁵ G. Ciani,⁵ F. Clara,³³ J. A. Clark,⁵⁸ F. Cleva,⁴⁷ E. Coccia,^{66,74} P.-F. Cohadon,⁵⁵ A. Colla,^{75,25} C. G. Collette,⁷⁶ M. Colombini,²⁹ M. Constancio Jr.,¹³ A. Conte,^{75,25} L. Conti,⁷⁷ D. Cook,³³ T. R. Corbitt,² N. Cornish,²⁷ A. Corsi,⁷⁸ C. A. Costa,¹³ M. W. Coughlin,⁷³ S. B. Coughlin,⁷ J.-P. Coulon,⁴⁷ S. T. Countryman,³⁵ P. Couvares,³¹ D. M. Coward,⁴⁵ M. J. Cowart,⁶ D. C. Coyne,¹ R. Coyne,⁷⁸ K. Craig,³² J. D. E. Creighton,¹⁶ J. Cripe,² S. G. Crowder,⁷⁹ A. Cumming,³² L. Cunningham,³² E. Cuoco,³⁰ T. Dal Canton,¹⁰ M. D. Damjanic,¹⁰ S. L. Danilishin,⁴⁵ S. D'Antonio,⁶⁷ K. Danzmann,^{17,10} N. S. Darman,⁸⁰ V. Dattilo,³⁰ I. Dave,⁴² H. P. Daveloza,⁸¹ M. Davies,²³ G. S. Davies,³² E. J. Daw,⁸² R. Day,³⁰ D. DeBra,²⁰ G. Debreczeni,⁸³ J. Degallaix,⁵⁹ M. De Laurentis,^{61,4} S. Deléglise,⁵⁵ W. Del Pozzo,³⁹ T. Denker,¹⁰ T. Dent,¹⁰ H. Dereli,⁴⁷ V. Dergachev,¹ R. De Rosa,^{61,4} R. T. DeRosa,² R. DeSalvo,⁹ S. Dhurandhar,¹⁴ M. C. Díaz,⁸¹ L. Di Fiore,⁴ M. Di Giovanni,^{75,25} A. Di Lieto,^{36,19} I. Di Palma,²⁶ A. Di Virgilio,¹⁹ G. Dojcinoski,⁸⁴ V. Dolique,⁵⁹ E. Dominguez,⁸⁵ F. Donovan,¹² K. L. Dooley,^{1,21} S. Doravari,⁶ R. Douglas,³² T. P. Downes,¹⁶ M. Drago,^{86,87} R. W. P. Drever,¹ J. C. Driggers,¹ Z. Du,⁶⁴ M. Ducrot,⁸ S. E. Dwyer,³³ T. B. Edo,⁸² M. C. Edwards,⁷³ M. Edwards,⁷ A. Effler,² H.-B. Eggenstein,¹⁰ P. Ehrens,¹ J. M. Eichholz,⁵ S. S. Eikenberry,⁵ R. C. Essick,¹² T. Etzel,¹ M. Evans,¹² T. M. Evans,⁶ R. Everett,⁸⁸ M. Factourovich,³⁵ V. Fafone,^{66,67,74} S. Fairhurst,⁷ Q. Fang,⁴⁵ S. Farinon,⁴¹ B. Farr,⁸⁹ W. M. Farr,³⁹ M. Favata,⁸⁴ M. Fays,⁷ H. Fehrmann,¹⁰ M. M. Fejer,²⁰ D. Feldbaum,^{5,6} I. Ferrante,^{36,19} E. C. Ferreira,¹³ F. Ferrini,³⁰ F. Fidecaro,^{36,19} I. Fiori,³⁰ R. P. Fisher,³¹ R. Flaminio,⁵⁹ J.-D. Fournier,⁴⁷ S. Franco,²³ S. Frasca,^{75,25} F. Frasconi,¹⁹ M. Frede,¹⁰ Z. Frei,⁴⁸ A. Freise,³⁹ R. Frey,⁵⁴ T. T. Fricke,¹⁰ P. Fritschel,¹² V. V. Frolov,⁶ P. Fulda,⁵ M. Fyffe,⁶ H. A. G. Gabbard,²¹ J. R. Gair,⁹⁰ L. Gammaitoni,^{28,29} S. G. Gaonkar,¹⁴ F. Garufi,^{61,4} A. Gatto,³⁴ N. Gehrels,⁶² G. Gemme,⁴¹ B. Gendre,⁴⁷ E. Genin,³⁰ A. Gennai,¹⁹ L. Á. Gergely,⁹¹ V. Germain,⁸ A. Ghosh,¹⁵ S. Ghosh,^{11,46} J. A. Giaime,^{2,6} K. D. Giardina,⁶ A. Giazotto,¹⁹ J. R. Gleason,⁵ E. Goetz,^{10,65} R. Goetz,⁵ L. Gondan,⁴⁸ G. González,² J. Gonzalez,^{36,19} A. Gopakumar,⁹² N. A. Gordon,³² M. L. Gorodetsky,⁴³ S. E. Gossan,⁷⁰ M. Gosselin,³⁰ S. Goßler,¹⁰ R. Gouaty,⁸ C. Graef,³² P. B. Graff,^{62,57} M. Granata,⁵⁹ A. Grant,³² S. Gras,¹² C. Gray,³³ G. Greco,^{51,52} P. Groot,⁴⁶ H. Grote,¹⁰ K. Grover,³⁹ S. Grunewald,²⁶ G. M. Guidi,^{51,52} C. J. Guido,⁶ X. Guo,⁶⁴ A. Gupta,¹⁴ M. K. Gupta,⁹³ K. E. Gushwa,¹ E. K. Gustafson,¹ R. Gustafson,⁶⁵ J. J. Hacker,²² B. R. Hall,⁵⁰ E. D. Hall,¹ D. Hammer,¹⁶ G. Hammond,³² M. Haney,⁹² M. M. Hanke,¹⁰ J. Hanks,³³ C. Hanna,⁸⁸ M. D. Hannam,⁷ J. Hanson,⁶ T. Hardwick,² J. Harms,^{51,52} G. M. Harry,⁹⁴ I. W. Harry,²⁶ M. J. Hart,³² M. T. Hartman,⁵ C.-J. Haster,³⁹ K. Haughian,³² A. Heidmann,⁵⁵ M. C. Heintze,^{5,6} H. Heitmann,⁴⁷ P. Hello,²³ G. Hemming,³⁰ M. Hendry,³² I. S. Heng,³² J. Hennig,³² A. W. Heptonstall,¹ M. Heurs,¹⁰ S. Hild,³² D. Hoak,⁹⁵ K. A. Hodge,¹ J. Hoelscher-Obermaier,¹⁷ D. Hofman,⁵⁹ S. E. Hollitt,⁹⁶ K. Holt,⁶ P. Hopkins,⁷ D. J. Hosken,⁹⁶ J. Hough,³² E. A. Houston,³² E. J. Howell,⁴⁵ Y. M. Hu,³² S. Huang,⁶⁸ E. A. Huerta,⁹⁷ D. Huet,²³ B. Hughey,⁵³ S. Husa,⁶⁰ S. H. Huttner,³² M. Huynh,¹⁶ T. Huynh-Dinh,⁶ A. Idrisy,⁸⁸ N. Indik,¹⁰ D. R. Ingram,³³ R. Inta,⁷⁸ G. Islas,²²

- J. C. Isler,³¹ T. Isogai,¹² B. R. Iyer,¹⁵ K. Izumi,³³ M. B. Jacobson,¹ H. Jang,⁹⁸ P. Jaranowski,⁹⁹ S. Jawahar,¹⁰⁰ Y. Ji,⁶⁴ F. Jiménez-Forteza,⁶⁰ W. W. Johnson,² D. I. Jones,²⁴ R. Jones,³² R.J.G. Jonker,¹¹ L. Ju,⁴⁵ Haris K,¹⁰¹ V. Kalogera,¹⁰² S. Kandhasamy,²¹ G. Kang,⁹⁸ J. B. Kanner,¹ S. Karki,⁵⁴ J. L. Karlen,⁹⁵ M. Kasprzak,^{23,30} E. Katsavounidis,¹² W. Katzman,⁶ S. Kaufer,¹⁷ T. Kaur,⁴⁵ K. Kawabe,³³ F. Kawazoe,¹⁰ F. Kéfélian,⁴⁷ M. S. Kehl,⁶³ D. Keitel,¹⁰ D. B. Kelley,³¹ W. Kells,¹ J. Kerrigan,⁹⁵ J. S. Key,⁸¹ F. Y. Khalili,⁴³ Z. Khan,⁹³ E. A. Khazanov,¹⁰³ N. Kijbunchoo,³³ C. Kim,⁹⁸ K. Kim,¹⁰⁴ N. G. Kim,⁹⁸ N. Kim,²⁰ Y.-M. Kim,⁷¹ E. J. King,⁹⁶ P. J. King,³³ D. L. Kinzel,⁶ J. S. Kissel,³³ S. Klimenko,⁵ J. T. Kline,¹⁶ S. M. Koehlenbeck,¹⁰ K. Kokeyama,² S. Koley,¹¹ V. Kondrashov,¹ M. Korobko,¹⁰ W. Z. Korth,¹ I. Kowalska,³⁸ D. B. Kozak,¹ V. Kringel,¹⁰ B. Krishnan,¹⁰ A. Królak,^{105,106} C. Krueger,¹⁷ G. Kuehn,¹⁰ A. Kumar,⁹³ P. Kumar,⁶³ L. Kuo,⁶⁸ A. Kutynia,¹⁰⁵ B. D. Lackey,³¹ M. Landry,³³ B. Lantz,²⁰ P. D. Lasky,^{80,107} A. Lazzarini,¹ C. Lazzaro,^{58,77} P. Leaci,^{26,75} S. Leavey,³² E. O. Lebigot,^{34,64} C. H. Lee,⁷¹ H. K. Lee,¹⁰⁴ H. M. Lee,¹⁰⁸ J. Lee,¹⁰⁴ J. P. Lee,¹² M. Leonardi,^{86,87} J. R. Leong,¹⁰ N. Leroy,²³ N. Letendre,⁸ Y. Levin,¹⁰⁷ B. M. Levine,³³ J. B. Lewis,¹ T. G. F. Li,¹ A. Libson,¹² A. C. Lin,²⁰ T. B. Littenberg,¹⁰² N. A. Lockerbie,¹⁰⁰ V. Lockett,²² D. Lodhia,³⁹ J. Logue,³² A. L. Lombardi,⁹⁵ M. Lorenzini,⁷⁴ V. Lorette,¹⁰⁹ M. Lormand,⁶ G. Losurdo,⁵² J. D. Lough,^{31,10} M. J. Lubinski(Ski),³³ H. Lück,^{17,10} A. P. Lundgren,¹⁰ J. Luo,⁷³ R. Lynch,¹² Y. Ma,⁴⁵ J. Macarthur,³² E. P. Macdonald,⁷ T. MacDonald,²⁰ B. Machenschalk,¹⁰ M. MacInnis,¹² D. M. Macleod,² D. X. Madden-Fong,²⁰ F. Magaña-Sandoval,³¹ R. M. Magee,⁵⁰ M. Mageswaran,¹ E. Majorana,²⁵ I. Maksimovic,¹⁰⁹ V. Malvezzi,^{66,67} N. Man,⁴⁷ I. Mandel,³⁹ V. Mandic,⁷⁹ V. Mangano,^{75,25,32} N. M. Mangini,⁹⁵ G. L. Mansell,⁷² M. Manske,¹⁶ M. Mantovani,³⁰ F. Marchesoni,^{110,29} F. Marion,⁸ S. Márka,³⁵ Z. Márka,³⁵ A. S. Markosyan,²⁰ E. Maros,¹ F. Martelli,^{51,52} L. Martellini,⁴⁷ I. W. Martin,³² R. M. Martin,⁵ D. V. Martynov,¹ J. N. Marx,¹ K. Mason,¹² A. Masserot,⁸ T. J. Massinger,³¹ S. Mastrogiovanni,^{75,25} F. Matichard,¹² L. Matone,³⁵ N. Mavalvala,¹² N. Mazumder,⁵⁰ G. Mazzolo,¹⁰ R. McCarthy,³³ D. E. McClelland,⁷² S. McCormick,⁶ S. C. McGuire,¹¹¹ G. McIntyre,¹ J. McIver,⁹⁵ S. T. McWilliams,⁹⁷ D. Meacher,⁴⁷ G. D. Meadors,¹⁰ M. Mehmet,¹⁰ J. Meidam,¹¹ M. Meinders,¹⁰ A. Melatos,⁸⁰ G. Mendell,³³ R. A. Mercer,¹⁶ M. Merzougui,⁴⁷ S. Meshkov,¹ C. Messenger,³² C. Messick,⁸⁸ P. M. Meyers,⁷⁹ F. Mezzani,^{25,75} H. Miao,³⁹ C. Michel,⁵⁹ H. Middleton,³⁹ E. E. Mikhailov,¹¹² L. Milano,^{61,4} J. Miller,¹² M. Millhouse,²⁷ Y. Minenkov,⁶⁷ J. Ming,²⁶ S. Mirshekari,¹¹³ C. Mishra,¹⁵ S. Mitra,¹⁴ V. P. Mitrofanov,⁴³ G. Mitselmakher,⁵ R. Mittleman,¹² B. Moe,¹⁶ A. Moggi,¹⁹ M. Mohan,³⁰ S. R. P. Mohapatra,¹² M. Montani,^{51,52} B. C. Moore,⁸⁴ D. Moraru,³³ G. Moreno,³³ S. R. Morris,⁸¹ K. Mossavi,¹⁰ B. Mours,⁸ C. M. Mow-Lowry,³⁹ C. L. Mueller,⁵ G. Mueller,⁵ A. Mukherjee,¹⁵ S. Mukherjee,⁸¹ A. Mullavey,⁶ J. Munch,⁹⁶ D. J. Murphy IV,³⁵ P. G. Murray,³² A. Mytidis,⁵ M. F. Nagy,⁸³ I. Nardecchia,^{66,67} L. Naticchioni,^{75,25} R. K. Nayak,¹¹⁴ V. Necula,⁵ K. Nedkova,⁹⁵ G. Nelemans,^{11,46} M. Neri,^{40,41} G. Newton,³² T. T. Nguyen,⁷² A. B. Nielsen,¹⁰ A. Nitz,³¹ F. Nocera,³⁰ D. Nolting,⁶ M. E. N. Normandin,⁸¹ L. K. Nuttall,¹⁶ E. Ochsner,¹⁶ J. O'Dell,¹¹⁵ E. Oelker,¹² G. H. Ogin,¹¹⁶ J. J. Oh,¹¹⁷ S. H. Oh,¹¹⁷ F. Ohme,⁷ M. Okounkova,⁷⁰ P. Oppermann,¹⁰ R. Oram,⁶ B. O'Reilly,⁶ W. E. Ortega,⁸⁵ R. O'Shaughnessy,¹¹⁸ C. D. Ott,⁷⁰ D. J. Ottaway,⁹⁶ R. S. Ottens,⁵ H. Overmier,⁶ B. J. Owen,⁷⁸ C. T. Padilla,²² A. Pai,¹⁰¹ S. A. Pai,⁴² J. R. Palamos,⁵⁴ O. Palashov,¹⁰³ C. Palomba,²⁵ A. Pal-Singh,¹⁰ H. Pan,⁶⁸ Y. Pan,⁵⁷ C. Pankow,¹⁶ F. Pannarale,⁷ B. C. Pant,⁴² F. Paoletti,^{30,19} M. A. Papa,^{26,16} H. R. Paris,²⁰ A. Pasqualetti,³⁰ R. Passaquieti,^{36,19} D. Passuello,¹⁹ Z. Patrick,²⁰ M. Pedraza,¹ L. Pekowsky,³¹ A. Pele,⁶ S. Penn,¹¹⁹ A. Perreca,³¹ M. Phelps,³² O. Piccinni,^{75,25} M. Pichot,⁴⁷ M. Pickenpack,¹⁰ F. Piergiovanni,^{51,52} V. Pierro,⁹ G. Pillant,³⁰ L. Pinard,⁵⁹ I. M. Pinto,⁹ M. Pitkin,³² J. H. Poeld,¹⁰ R. Poggiani,^{36,19} A. Post,¹⁰ J. Powell,³² J. Prasad,¹⁴ V. Predoi,⁷ S. S. Premachandra,¹⁰⁷ T. Prestegard,⁷⁹ L. R. Price,¹ M. Prijatelj,³⁰ M. Principe,⁹ S. Privitera,²⁶ R. Prix,¹⁰ G. A. Prodi,^{86,87} L. Prokhorov,⁴³ O. Puncken,^{81,10} M. Punturo,²⁹ P. Puppo,²⁵ M. Pürner,⁷ J. Qin,⁴⁵ V. Quetschke,⁸¹ E. A. Quintero,¹ R. Quitzow-James,⁵⁴ F. J. Raab,³³ D. S. Rabeling,⁷² I. Rácz,⁸³ H. Radkins,³³ P. Raffai,⁴⁸ S. Raja,⁴² M. Rakhmanov,⁸¹ P. Rapagnani,^{75,25} V. Raymond,²⁶ M. Razzano,^{36,19} V. Re,^{66,67} C. M. Reed,³³ T. Regimbau,⁴⁷ L. Rei,⁴¹ S. Reid,⁴⁴ D. H. Reitze,^{1,5} F. Ricci,^{75,25} K. Riles,⁶⁵ N. A. Robertson,^{1,32} R. Robie,³² F. Robinet,²³ A. Rocchi,⁶⁷ A. S. Rodger,³² L. Rolland,⁸ J. G. Rollins,¹ V. J. Roma,⁵⁴ J. D. Romano,⁸¹ R. Romano,^{3,4} G. Romanov,¹¹² J. H. Romie,⁶ D. Rosińska,^{120,37} S. Rowan,³² A. Rüdiger,¹⁰ P. Ruggi,³⁰ K. Ryan,³³ S. Sachdev,¹ T. Sadecki,³³ L. Sadeghian,¹⁶ M. Saleem,¹⁰¹ F. Salemi,¹⁰ L. Sammut,⁸⁰ E. Sanchez,¹ V. Sandberg,³³ J. R. Sanders,⁶⁵ I. Santiago-Prieto,³² B. Sassolas,⁵⁹ B. S. Sathyaprakash,⁷ P. R. Saulson,³¹ R. Savage,³³ A. Sawadsky,¹⁷ P. Schale,⁵⁴ R. Schilling,¹⁰ P. Schmidt,¹ R. Schnabel,¹⁰ R. M. S. Schofield,⁵⁴ A. Schönbeck,¹⁰ E. Schreiber,¹⁰ D. Schuette,¹⁰ B. F. Schutz,⁷ J. Scott,³² S. M. Scott,⁷² D. Sellers,⁶ D. Sentenac,³⁰ V. Sequino,^{66,67} A. Sergeev,¹⁰³ G. Serna,²² A. Seigny,³³ D. A. Shaddock,⁷² P. Shaffery,¹⁰⁸ S. Shah,^{11,46} M. S. Shahriar,¹⁰² M. Shaltev,¹⁰ Z. Shao,¹ B. Shapiro,²⁰ P. Shawhan,⁵⁷ D. H. Shoemaker,¹² T. L. Sidery,³⁹ K. Siellez,⁴⁷ X. Siemens,¹⁶ D. Sigg,³³ A. D. Silva,¹³ D. Simakov,¹⁰ A. Singer,¹ L. P. Singer,⁶² R. Singh,²

- A. M. Sintes,⁶⁰ B. J. J. Slagmolen,⁷² J. R. Smith,²² N. D. Smith,¹ R. J. E. Smith,¹ E. J. Son,¹¹⁷ B. Sorazu,³²
 T. Souradeep,¹⁴ A. K. Srivastava,⁹³ A. Staley,³⁵ M. Steinke,¹⁰ J. Steinlechner,³² S. Steinlechner,³²
 D. Steinmeyer,¹⁰ B. C. Stephens,¹⁶ S. Steplewski,⁵⁰ S. P. Stevenson,³⁹ R. Stone,⁸¹ K. A. Strain,³² N. Straniero,⁵⁹
 N. A. Strauss,⁷³ S. Strigin,⁴³ R. Sturani,¹¹³ A. L. Stuver,⁶ T. Z. Summerscales,¹²¹ L. Sun,⁸⁰ P. J. Sutton,⁷
 B. L. Swinkels,³⁰ M. J. Szczepanczyk,⁵³ M. Tacca,³⁴ D. Talukder,⁵⁴ D. B. Tanner,⁵ M. Tápai,⁹¹ S. P. Tarabrin,¹⁰
 A. Taracchini,²⁶ R. Taylor,¹ T. Theeg,¹⁰ M. P. Thirugnanasambandam,¹ M. Thomas,⁶ P. Thomas,³³
 K. A. Thorne,⁶ K. S. Thorne,⁷⁰ E. Thrane,¹⁰⁷ S. Tiwari,⁷⁴ V. Tiwari,⁵ K. V. Tokmakov,¹⁰⁰ C. Tomlinson,⁸²
 M. Tonelli,^{36,19} C. V. Torres,⁸¹ C. I. Torrie,¹ F. Travasso,^{28,29} G. Traylor,⁶ D. Trifirò,²¹ M. C. Tringali,^{86,87}
 M. Tse,¹² M. Turconi,⁴⁷ D. Ugolini,¹²² C. S. Unnikrishnan,⁹² A. L. Urban,¹⁶ S. A. Usman,³¹ H. Vahlbruch,¹⁰
 G. Vajente,¹ G. Valdes,⁸¹ M. Vallisneri,⁷⁰ N. van Bakel,¹¹ M. van Beuzekom,¹¹ J. F. J. van den Brand,^{56,11}
 C. van den Broeck,¹¹ L. van der Schaaf,¹¹ M. V. van der Sluys,^{11,46} J. van Heijningen,¹¹ A. A. van Veggel,³²
 M. Vardaro,^{123,77} S. Vass,¹ M. Vasúth,⁸³ R. Vaulin,¹² A. Vecchio,³⁹ G. Vedovato,⁷⁷ J. Veitch,³⁹ P. J. Veitch,⁹⁶
 K. Venkateswara,¹²⁴ D. Verkindt,⁸ F. Vetrano,^{51,52} A. Viceré,^{51,52} J.-Y. Vinet,⁴⁷ S. Vitale,¹² T. Vo,³¹ H. Vocca,^{28,29}
 C. Vorvick,³³ W. D. Voudsen,³⁹ S. P. Vyatchanin,⁴³ A. R. Wade,⁷² M. Wade,¹⁶ L. E. Wade IV,¹⁶ M. Walker,²
 L. Wallace,¹ S. Walsh,¹⁶ G. Wang,⁷⁴ H. Wang,³⁹ M. Wang,³⁹ X. Wang,⁶⁴ R. L. Ward,⁷² J. Warner,³³ M. Was,⁸
 B. Weaver,³³ L.-W. Wei,⁴⁷ M. Weinert,¹⁰ A. J. Weinstein,¹ R. Weiss,¹² T. Welborn,⁶ L. Wen,⁴⁵ P. Weßels,¹⁰
 T. Westphal,¹⁰ K. Wette,¹⁰ J. T. Whelan,^{118,10} S. E. Whitcomb,¹ D. J. White,⁸² B. F. Whiting,⁵ K. J. Williams,¹¹¹
 L. Williams,⁵ R. D. Williams,¹ A. R. Williamson,⁷ J. L. Willis,¹²⁵ B. Willke,^{17,10} M. H. Wimmer,¹⁰ W. Winkler,¹⁰
 C. C. Wipf,¹ H. Wittel,¹⁰ G. Woan,³² J. Worden,³³ J. Yablon,¹⁰² I. Yakushin,⁶ W. Yam,¹² H. Yamamoto,¹
 C. C. Yancey,⁵⁷ M. Yvert,⁸ A. Zadrożny,¹⁰⁵ L. Zangrando,⁷⁷ M. Zanolin,⁵³ J.-P. Zendri,⁷⁷ Fan Zhang,¹² L. Zhang,¹
 M. Zhang,¹¹² Y. Zhang,¹¹⁸ C. Zhao,⁴⁵ M. Zhou,¹⁰² X. J. Zhu,⁴⁵ M. E. Zucker,¹² S. E. Zuraw,⁹⁵ and J. Zweizig¹

¹LIGO—California Institute of Technology, Pasadena, CA 91125, USA

²Louisiana State University, Baton Rouge, LA 70803, USA

³Università di Salerno, Fisciano, I-84084 Salerno, Italy

⁴INFN, Sezione di Napoli, Complesso Universitario di Monte S. Angelo, I-80126 Napoli, Italy

⁵University of Florida, Gainesville, FL 32611, USA

⁶LIGO Livingston Observatory, Livingston, LA 70754, USA

⁷Cardiff University, Cardiff CF24 3AA, United Kingdom

⁸Laboratoire d'Annecy-le-Vieux de Physique des Particules (LAPP),
 Université Savoie Mont Blanc, CNRS/IN2P3, F-74941 Annecy-le-Vieux, France

⁹University of Sannio at Benevento, I-82100 Benevento,

Italy and INFN, Sezione di Napoli, I-80100 Napoli, Italy

¹⁰Albert-Einstein-Institut, Max-Planck-Institut für Gravitationsphysik, D-30167 Hannover, Germany

¹¹Nikhef, Science Park, 1098 XG Amsterdam, The Netherlands

¹²LIGO—Massachusetts Institute of Technology, Cambridge, MA 02139, USA

¹³Instituto Nacional de Pesquisas Espaciais, 12227-010 São José dos Campos, SP, Brazil

¹⁴Inter-University Centre for Astronomy and Astrophysics, Pune 411007, India

¹⁵International Centre for Theoretical Sciences, Tata Institute of Fundamental Research, Bangalore 560012, India

¹⁶University of Wisconsin-Milwaukee, Milwaukee, WI 53201, USA

¹⁷Leibniz Universität Hannover, D-30167 Hannover, Germany

¹⁸Università di Siena, I-53100 Siena, Italy

¹⁹INFN, Sezione di Pisa, I-56127 Pisa, Italy

²⁰Stanford University, Stanford, CA 94305, USA

²¹The University of Mississippi, University, MS 38677, USA

²²California State University Fullerton, Fullerton, CA 92831, USA

²³LAL, Université Paris-Sud, IN2P3/CNRS, F-91898 Orsay, France

²⁴University of Southampton, Southampton SO17 1BJ, United Kingdom

²⁵INFN, Sezione di Roma, I-00185 Roma, Italy

²⁶Albert-Einstein-Institut, Max-Planck-Institut für Gravitationsphysik, D-14476 Golm, Germany

²⁷Montana State University, Bozeman, MT 59717, USA

²⁸Università di Perugia, I-06123 Perugia, Italy

²⁹INFN, Sezione di Perugia, I-06123 Perugia, Italy

³⁰European Gravitational Observatory (EGO), I-56021 Cascina, Pisa, Italy

³¹Syracuse University, Syracuse, NY 13244, USA

³²SUPA, University of Glasgow, Glasgow G12 8QQ, United Kingdom

³³LIGO Hanford Observatory, Richland, WA 99352, USA

³⁴APC, AstroParticule et Cosmologie, Université Paris Diderot,

CNRS/IN2P3, CEA/Irfu, Observatoire de Paris,

Sorbonne Paris Cité, F-75205 Paris Cedex 13, France

³⁵Columbia University, New York, NY 10027, USA

- ³⁶ *Università di Pisa, I-56127 Pisa, Italy*
³⁷ *CAMK-PAN, 00-716 Warsaw, Poland*
³⁸ *Astronomical Observatory Warsaw University, 00-478 Warsaw, Poland*
³⁹ *University of Birmingham, Birmingham B15 2TT, United Kingdom*
⁴⁰ *Università degli Studi di Genova, I-16146 Genova, Italy*
⁴¹ *INFN, Sezione di Genova, I-16146 Genova, Italy*
⁴² *RRCAT, Indore MP 452013, India*
⁴³ *Faculty of Physics, Lomonosov Moscow State University, Moscow 119991, Russia*
⁴⁴ *SUPA, University of the West of Scotland, Paisley PA1 2BE, United Kingdom*
⁴⁵ *University of Western Australia, Crawley, Western Australia 6009, Australia*
⁴⁶ *Department of Astrophysics/IMAPP, Radboud University Nijmegen, P.O. Box 9010, 6500 GL Nijmegen, The Netherlands*
⁴⁷ *ARTEMIS, Université Nice-Sophia-Antipolis, CNRS and Observatoire de la Côte d'Azur, F-06304 Nice, France*
⁴⁸ *MTA Eötvös University, "Lendület" Astrophysics Research Group, Budapest 1117, Hungary*
⁴⁹ *Institut de Physique de Rennes, CNRS, Université de Rennes 1, F-35042 Rennes, France*
⁵⁰ *Washington State University, Pullman, WA 99164, USA*
⁵¹ *Università degli Studi di Urbino 'Carlo Bo', I-61029 Urbino, Italy*
⁵² *INFN, Sezione di Firenze, I-50019 Sesto Fiorentino, Firenze, Italy*
⁵³ *Embry-Riddle Aeronautical University, Prescott, AZ 86301, USA*
⁵⁴ *University of Oregon, Eugene, OR 97403, USA*
⁵⁵ *Laboratoire Kastler Brossel, UPMC-Sorbonne Universités, CNRS, ENS-PSL Research University, Collège de France, F-75005 Paris, France*
⁵⁶ *VU University Amsterdam, 1081 HV Amsterdam, The Netherlands*
⁵⁷ *University of Maryland, College Park, MD 20742, USA*
⁵⁸ *Center for Relativistic Astrophysics and School of Physics, Georgia Institute of Technology, Atlanta, GA 30332, USA*
⁵⁹ *Laboratoire des Matériaux Avancés (LMA), IN2P3/CNRS, Université de Lyon, F-69622 Villeurbanne, Lyon, France*
⁶⁰ *Universitat de les Illes Balears—IEEC, E-07122 Palma de Mallorca, Spain*
⁶¹ *Università di Napoli 'Federico II', Complesso Universitario di Monte S. Angelo, I-80126 Napoli, Italy*
⁶² *NASA/Goddard Space Flight Center, Greenbelt, MD 20771, USA*
⁶³ *Canadian Institute for Theoretical Astrophysics, University of Toronto, Toronto, Ontario M5S 3H8, Canada*
⁶⁴ *Tsinghua University, Beijing 100084, China*
⁶⁵ *University of Michigan, Ann Arbor, MI 48109, USA*
⁶⁶ *Università di Roma Tor Vergata, I-00133 Roma, Italy*
⁶⁷ *INFN, Sezione di Roma Tor Vergata, I-00133 Roma, Italy*
⁶⁸ *National Tsing Hua University, Hsinchu Taiwan 300*
⁶⁹ *Charles Sturt University, Wagga Wagga, New South Wales 2678, Australia*
⁷⁰ *Caltech—CaRT, Pasadena, CA 91125, USA*
⁷¹ *Pusan National University, Busan 609-735, Korea*
⁷² *Australian National University, Canberra, Australian Capital Territory 0200, Australia*
⁷³ *Carleton College, Northfield, MN 55057, USA*
⁷⁴ *INFN, Gran Sasso Science Institute, I-67100 L'Aquila, Italy*
⁷⁵ *Università di Roma 'La Sapienza', I-00185 Roma, Italy*
⁷⁶ *University of Brussels, Brussels 1050, Belgium*
⁷⁷ *INFN, Sezione di Padova, I-35131 Padova, Italy*
⁷⁸ *Texas Tech University, Lubbock, TX 79409, USA*
⁷⁹ *University of Minnesota, Minneapolis, MN 55455, USA*
⁸⁰ *The University of Melbourne, Parkville, Victoria 3010, Australia*
⁸¹ *The University of Texas at Brownsville, Brownsville, TX 78520, USA*
⁸² *The University of Sheffield, Sheffield S10 2TN, United Kingdom*
⁸³ *Wigner RCP, RMKI, H-1121 Budapest, Konkoly Thege Miklós út 29-33, Hungary*
⁸⁴ *Montclair State University, Montclair, NJ 07043, USA*
⁸⁵ *Argentinian Gravitational Wave Group, Cordoba Cordoba 5000, Argentina*
⁸⁶ *Università di Trento, Dipartimento di Fisica, I-38123 Povo, Trento, Italy*
⁸⁷ *INFN, Trento Institute for Fundamental Physics and Applications, I-38123 Povo, Trento, Italy*
⁸⁸ *The Pennsylvania State University, University Park, PA 16802, USA*
⁸⁹ *University of Chicago, Chicago, IL 60637, USA*
⁹⁰ *University of Cambridge, Cambridge CB2 1TN, United Kingdom*
⁹¹ *University of Szeged, Dóm tér 9, Szeged 6720, Hungary*
⁹² *Tata Institute for Fundamental Research, Mumbai 400005, India*
⁹³ *Institute for Plasma Research, Bhat, Gandhinagar 382428, India*
⁹⁴ *American University, Washington, D.C. 20016, USA*

- ⁹⁵ *University of Massachusetts-Amherst, Amherst, MA 01003, USA*
⁹⁶ *University of Adelaide, Adelaide, South Australia 5005, Australia*
⁹⁷ *West Virginia University, Morgantown, WV 26506, USA*
⁹⁸ *Korea Institute of Science and Technology Information, Daejeon 305-806, Korea*
⁹⁹ *University of BiaŁystok, 15-424 BiaŁystok, Poland*
¹⁰⁰ *SUPA, University of Strathclyde, Glasgow G1 1XQ, United Kingdom*
¹⁰¹ *IISER-TVM, CET Campus, Trivandrum Kerala 695016, India*
¹⁰² *Northwestern University, Evanston, IL 60208, USA*
¹⁰³ *Institute of Applied Physics, Nizhny Novgorod, 603950, Russia*
¹⁰⁴ *Hanyang University, Seoul 133-791, Korea*
¹⁰⁵ *NCBJ, 05-400 Swierk-Otwock, Poland*
¹⁰⁶ *IM-PAN, 00-956 Warsaw, Poland*
¹⁰⁷ *Monash University, Victoria 3800, Australia*
¹⁰⁸ *Seoul National University, Seoul 151-742, Korea*
¹⁰⁹ *ESPCI, CNRS, F-75005 Paris, France*
¹¹⁰ *Università di Camerino, Dipartimento di Fisica, I-62032 Camerino, Italy*
¹¹¹ *Southern University and A&M College, Baton Rouge, LA 70813, USA*
¹¹² *College of William and Mary, Williamsburg, VA 23187, USA*
¹¹³ *Instituto de Física Teórica, University Estadual Paulista/ICTP South American Institute for Fundamental Research, São Paulo SP 01140-070, Brazil*
¹¹⁴ *IISER-Kolkata, Mohanpur, West Bengal 741252, India*
¹¹⁵ *Rutherford Appleton Laboratory, HSIC, Chilton, Didcot, Oxon OX11 0QX, United Kingdom*
¹¹⁶ *Whitman College, 280 Boyer Ave, Walla Walla, WA 9936, USA*
¹¹⁷ *National Institute for Mathematical Sciences, Daejeon 305-390, Korea*
¹¹⁸ *Rochester Institute of Technology, Rochester, NY 14623, USA*
¹¹⁹ *Hobart and William Smith Colleges, Geneva, NY 14456, USA*
¹²⁰ *Institute of Astronomy, 65-265 Zielona Góra, Poland*
¹²¹ *Andrews University, Berrien Springs, MI 49104, USA*
¹²² *Trinity University, San Antonio, TX 78212, USA*
¹²³ *Università di Padova, Dipartimento di Fisica e Astronomia, I-35131 Padova, Italy*
¹²⁴ *University of Washington, Seattle, WA 98195, USA and*
¹²⁵ *Abilene Christian University, Abilene, TX 79699, USA*

In this paper we present the results of the first low frequency all-sky search of continuous gravitational wave signals conducted on Virgo VSR2 and VSR4 data. The search covered the full sky, a frequency range between 20 Hz and 128 Hz with a range of spin-down between -1.0×10^{-10} Hz/s and $+1.5 \times 10^{-11}$ Hz/s, and was based on a hierarchical approach. The starting point was a set of short Fast Fourier Transforms (FFT), of length 8192 seconds, built from the calibrated strain data. Aggressive data cleaning, both in the time and frequency domains, has been done in order to remove, as much as possible, the effect of disturbances of instrumental origin. On each dataset a number of candidates has been selected, using the FrequencyHough transform in an incoherent step. Only coincident candidates among VSR2 and VSR4 have been examined in order to strongly reduce the false alarm probability, and the most significant candidates have been selected. The criteria we have used for candidate selection and for the coincidence step greatly reduce the harmful effect of large instrumental artifacts. Selected candidates have been subject to a follow-up by constructing a new set of longer FFTs followed by a further incoherent analysis, still based on the FrequencyHough transform. No evidence for continuous gravitational wave signals was found, therefore we have set a population-based joint VSR2-VSR4 90% confidence level upper limit on the dimensionless gravitational wave strain in the frequency range between 20 Hz and 128 Hz. This is the first all-sky search for continuous gravitational waves conducted at frequencies below 50 Hz. We set upper limits in the range between about 10^{-24} and 2×10^{-23} at most frequencies. Our upper limits on signal strain show an improvement of up to a factor of ~ 2 with respect to the results of previous all-sky searches at frequencies below 80 Hz.

PACS numbers: 04.80.Nn, 95.55.Ym, 97.60.Gb, 07.05.Kf

I. INTRODUCTION

Continuous gravitational wave signals (CW) emitted by asymmetric spinning neutron stars are among the sources currently sought in the data of interferometric gravitational wave detectors. The search for signals emit-

ted by spinning neutron stars with no electromagnetic counterpart requires the exploration of a large portion of the source parameter space, consisting of the source position, signal frequency and signal frequency time-derivative (spin-down). This kind of search, called all-sky, cannot be based on fully coherent methods, as in

targeted searches for known pulsars, see e.g. [1, 2], because of the huge computational resources that would be required.

For this reason various hierarchical analysis pipelines, based on the alternation of coherent and incoherent steps, have been developed [3], [4], [5], [6], [7]. They allow us to dramatically reduce the computational burden of the analysis, at the cost of a small sensitivity loss. In this paper we present the results of the first all-sky search for CW signals using the data of Virgo science runs VSR2 and VSR4 (discussed in Sec. III). The analysis has been carried out on the frequency band 20-128 Hz, using an efficient hierarchical analysis pipeline, based on the FrequencyHough transform [7]. No detection was made, so we established upper limits on signal strain amplitude as a function of the frequency. The rationale for the chosen frequency band is twofold. First, frequencies below 50 Hz have never been considered in all-sky searches for CW signals, and the estimated joint sensitivity of Virgo VSR2 and VSR4 data is better than that of data from LIGO science runs S5 and S6 below about 60-70 Hz. Second, lower frequencies potentially offer promising sources. Higher frequency signals would be in principle easier to detect because of their high signal amplitudes at fixed distance and ellipticity (see Eq. 5). On the other hand, based on the electromagnetically observed pulsar population and the results of population synthesis modelling, e.g. [8], [9], we expect that a substantial fraction of the galactic neutron star population emits gravitational waves at frequencies below ~ 100 Hz. This is particularly true when considering young, unrecycled, neutron stars, which could be more distorted than older objects. A search at low frequency, as described below, could detect signals from a potentially significant population of nearby neutron stars.

The plan of the paper is as follows. In Sec. II we describe the kind of gravitational wave (GW) signal we are searching for. In Sec. III we discuss the Virgo detector performance during VSR2 and VSR4 runs. In Sec. IV we briefly recap the analysis procedure, referring the reader to [7] for more details. Section V is focused on the cleaning steps applied at different stages of the analysis. Section VI is dedicated to candidate selection, and Sec. VII to their clustering and coincidences. Section VIII deals with the follow-up of candidates surviving the coincidence step. Section IX is dedicated to validation tests of the analysis pipeline, by using hardware-injected signals in VSR2 and VSR4 data. In Sec. X a joint upper limit on signal strain amplitude is derived as a function of the search frequency. Conclusions and future prospects are presented in Sec. XI. Appendix A contains a list of the 108 candidates for which the follow-up has been done, along with their main parameters. Appendix B is devoted to a deeper analysis of the three outliers found. Appendix C contains the list of frequency intervals excluded from the computation of the upper limits.

II. THE SIGNAL

The expected quadrupolar GW signal from a non-axisymmetric neutron star steadily spinning around one of its principal axes has a frequency f_0 twice the rotation frequency f_{rot} , with a strain at the detector of [7, 10]

$$h(t) = H_0(H_+A^+ + H_\times A^\times)e^{j(\omega(t)t + \Phi_0)}, \quad (1)$$

where taking the real part is understood and where Φ_0 is an initial phase. The signal's time-dependent angular frequency $\omega(t)$ will be discussed below. The two complex amplitudes H_+ and H_\times are given respectively by

$$H_+ = \frac{\cos 2\psi - \eta \sin 2\psi}{\sqrt{1 + \eta^2}}, \quad (2)$$

$$H_\times = \frac{\sin 2\psi + \eta \cos 2\psi}{\sqrt{1 + \eta^2}}, \quad (3)$$

in which η is the ratio of the polarization ellipse semi-minor to semi-major axis, and the polarization angle ψ defines the direction of the major axis with respect to the celestial parallel of the source (counterclockwise). The parameter η varies in the range $[-1, 1]$, where $\eta = 0$ for a linearly polarized wave, while $\eta = \pm 1$ for a circularly polarized wave ($\eta = 1$ if the circular rotation is counterclockwise). The functions $A^{+, \times}$ describe the detector response as a function of time, with a periodicity of one and two sidereal periods, and depend on the source position, detector position and orientation on the Earth [10].

As discussed in [1], the strain described by Eq.(1) is equivalent to the standard expression (see e.g. [11])

$$h(t) = \frac{1}{2}F_+(t, \psi)h_0(1 + \cos^2 \iota) \cos \Phi(t) + F_\times(t, \psi)h_0 \cos \iota \sin \Phi(t) \quad (4)$$

Here F_+ , F_\times are the standard beam-pattern functions and ι is the angle between the star's rotation axis and the line of sight. The amplitude parameter

$$h_0 = \frac{4\pi^2 G}{c^4} \frac{I_{zz} \varepsilon f_0^2}{d} \quad (5)$$

depends on the signal frequency f_0 and on the source distance d ; on I_{zz} , the star's moment of inertia with respect to the principal axis aligned with the rotation axis; and on ε , which is the fiducial equatorial ellipticity expressed in terms of principal moments of inertia as

$$\varepsilon = \frac{I_{xx} - I_{yy}}{I_{zz}}. \quad (6)$$

It must be stressed that it is not the fiducial ellipticity but the quadrupole moment $Q_{22} \propto I_{zz} \varepsilon$ that, in case of detection, can be measured independently of any assumption about the star's equation of state and moment

of inertia (assuming the source distance can be also estimated). There exist estimates of the maximum ellipticity a neutron star can sustain from both elastic and magnetic deformations. In the elastic case, these maxima depend strongly on the breaking strain of the solid portion sustaining the deformation (see, e.g., [12], [13] for calculations for the crust) as well as on the star's structure and equation of state, and the possible presence of exotic phases in the stars interior (like in hybrid or strange quark stars, see, e.g., [14]). In the magnetic case, the deformation depends on the strength and configuration of the star's internal magnetic field (see, e.g., [15]). However, the actual ellipticity of a given neutron star is unknown—the best we have are observational upper limits. The relations between H_0 , η and h_0 , ι are given, e.g., in [10]. In Eq. 1 the signal angular frequency $\omega(t)$ is a function of time, therefore the signal phase

$$\Phi(t) = \int_{t_0}^t \omega(t') dt' \quad (7)$$

is not that of a simple monochromatic signal. It depends on the rotational frequency and frequency derivatives of the neutron star, as well as on Doppler and propagation effects. In particular, the received Doppler-shifted frequency $f(t)$ is related to the emitted frequency $f_0(t)$ by the well-known relation (valid in the non-relativistic approximation)

$$f(t) = \frac{1}{2\pi} \frac{d\Phi(t)}{dt} = f_0(t) \left(1 + \frac{\vec{v}(t) \cdot \hat{n}}{c} \right), \quad (8)$$

where \vec{v} is the detector velocity with respect to the Solar System barycenter (SSB), \hat{n} is the unit vector in the direction to the source from the SSB and c is the light speed. Smaller relativistic effects, namely the *Einstein delay* and the *Shapiro delay* are not relevant for the search described in this paper, due to the use of short length FFTs, and are therefore neglected.

The intrinsic signal frequency $f_0(t)$ slowly decreases in time due to the source's spin-down, associated with the rotational energy loss following emission of electromagnetic and/or gravitational radiation. The spin-down can be described through a series expansion

$$f_0(t) = f_0 + \dot{f}_0(t - t_0) + \frac{\ddot{f}_0}{2}(t - t_0)^2 + \dots \quad (9)$$

In general, the frequency evolution of a CW depends on $3+s$ parameters: position, frequency and s spin-down parameters. In the all-sky search described in this paper we need to take into account only the first spin-down ($s = 1$) parameter (see Sec. IV).

III. INSTRUMENTAL PERFORMANCE DURING VSR2 AND VSR4 RUNS

Interferometric GW detectors, such as LIGO [16], Virgo [17], and GEO [18], have collected years of data,

from 2002 to 2011. For the analysis described in this paper we have used calibrated data from the Virgo VSR2 and VSR4 science runs. The VSR3 run was characterized by a diminished sensitivity level and poor data quality (highly non-stationary data, large glitch rate), and so was not included in this analysis. The VSR2 run began on July 7th, 2009 (21:00 UTC) and ended on January 8th, 2010 (22:00 UTC). The duty cycle was 80.4%, resulting in a total of ~ 149 days of *science mode* data, divided among 361 segments. The data used in the analysis have been produced using the most up-to-date calibration parameters and reconstruction procedure. The associated systematic error amounts to 5.5% in amplitude and ~ 50 mrad in phase [19].

The VSR4 run extended from June 3rd, 2011 (10:27 UTC) to September 5th, 2011 (13:26 UTC), with a duty factor of about 81%, corresponding to an effective duration of 76 days. Calibration uncertainties amounted to 7.5% in amplitude and $(40 + 50 f_{\text{kHz}})$ mrad in phase up to 500 Hz, where f_{kHz} is the frequency in kilohertz [20]. The uncertainty on the amplitude contributes to the uncertainty on the upper limit on signal amplitude, together with that coming from the finite size of the Monte Carlo simulation used to compute it (see Sec. X). A calibration error on the phase of this size can be shown to have a negligible impact on the analysis [1]. The low-frequency sensitivity of VSR4 was significantly better, up to a factor of 2, than that of previous Virgo runs, primarily due to the use of monolithic mirror suspensions, and nearly in agreement with the design sensitivity of the initial Virgo interferometer [17]. We show in Fig. 1 the average experimental strain amplitude spectral density for VSR2 and VSR4, in the frequency range 20-128 Hz, obtained by making an average of the periodograms (squared modulus of the FFTs) stored in the short FFT database (see next section).

IV. THE ANALYSIS PROCEDURE

All-sky searches are intractable using completely coherent methods because of the huge size of the parameter space, which poses challenging computational problems [21], [22]. Moreover, a completely coherent search would not be robust against unpredictable phase variations of the signal during the observation time.

For these reasons hierarchical schemes have been developed. The hierarchical scheme we have used for this analysis has been described in detail in [7]. In this section we briefly recall the main steps. The analysis starts from the detector calibrated data, sampled at 4096 Hz. The first step consists of constructing a database of *short Discrete Fourier Transforms* (SFDB) [23], computed through the FFT algorithm (the FFT is just an efficient algorithm to compute Discrete Fourier Transforms (DFT), but for historical reasons and for consistency with previous papers we will use the term FFT instead of DFT). Each FFT covers the frequency range from 20 to 128Hz and




FIG. 1. VSR2 (darker, black in the color version) and VSR4 (lighter, red in the color version) average strain amplitude spectral density in the frequency range from 20 Hz up to 128 Hz.

is built from a data chunk of duration (*coherence time*) short enough such that if a signal is present, its frequency (modified by Doppler and spin-down) does not shift more than a frequency bin. The FFT duration for this search is 8192 seconds. This corresponds to a natural frequency resolution $\delta f = 1.22 \times 10^{-4}$ Hz. The FFTs are interlaced by half and windowed with a Tukey window with a width parameter $\alpha = 0.5$. Before constructing the SFDB, short strong time domain disturbances are removed from the data. This is the first of several cleaning steps applied to the data (see Sec. V). The total number of FFTs for VSR2 run is 3896 and for VSR4 run is 1978.

From the SFDB we create a time-frequency map, called the *peakmap* [24]. This is obtained by selecting the most significant local maxima (which we call *peaks*) of the square root of equalized periodograms, obtained by dividing the periodogram by an auto-regressive average spectrum estimation. The threshold for peak selection has been chosen equal to $\sqrt{2.5} = 1.58$ [7] which, in the ideal case of Gaussian noise, would correspond to a probability of selecting a peak of 0.0755. The peakmap is cleaned by removing peaks clearly due to disturbances, as explained in Sec. V. The peakmap is then corrected for the Doppler shift for the different sky directions, by shifting the frequency of the peaks by an amount corresponding to the variation the frequency of a signal coming from a given direction would be subject to at a given time. A *coarse* grid in the sky is used in this stage of the analysis. The grid is built using ecliptic coordinates, as described in [7]. Figure 2 shows the number of sky points (“patches”) as a function of the frequency (in steps of 1 Hz), for both VSR2 and VSR4 analyses. The number of patches increases with the square of the frequency, and ranges from 2492 at 20 Hz to 81244 at 128 Hz. The total number of sky patches is $N_{\text{sky}} \approx 3.5 \times 10^6$.

Each corrected peakmap is the input of the inco-

herent step, based on the FrequencyHough transform [7, 25]. This is a very efficient implementation of the Hough transform (see [25] for efficiency tests and comparison with a different implementation) which, for every sky position, maps the points of the peakmap into the signal frequency/spin-down plane. In the FrequencyHough transform we take into account slowly varying non-stationarity in the noise and the varying detector sensitivity caused by the time-dependent radiation pattern [26], [27]. Furthermore, the frequency/spin-down plane is discretized by building a suitable grid [7]. As the transformation from the peakmap to the Hough plane is not computationally bounded by the size of the frequency bin (which only affects the size of the Hough map) we have increased the frequency resolution by a factor of 10, with respect to the natural step size δf , in order to reduce the digitalization loss, so that the actual resolution is $\delta f_H = \delta f/10 = 1.22 \times 10^{-5}$ Hz.

We have searched approximately over the spin-down range $[-1.0 \times 10^{-10}, +1.5 \times 10^{-11}]$ Hz/s. This choice has been dictated by the need to not increase too much the computational load of the analysis while, at the same time, covering a range of spin-down values including the values measured for most known pulsars. Given the spin-down bin width scales as T_{obs}^{-1} ($\delta \dot{f} = \delta f/T_{\text{obs}}$), this implies a different number of spin-down values for the two data sets: $N_{\text{sd}}=16$ for VSR2 with a resolution of $\delta \dot{f} = 7.63 \times 10^{-12}$ Hz/s, and $N_{\text{sd}}=9$ for VSR4 with a resolution of $\delta \dot{f} = 1.5 \times 10^{-11}$ Hz/s. The corresponding minimum gravitational-wave spin-down age, defined as $\tau_{\text{min}}(f) = f/4N_{\text{sd}}\delta \dot{f}$ (where $4N_{\text{sd}}\delta \dot{f}$ is the absolute value of the maximum spin-down we have searched over), is a function of the frequency, going from ~ 1600 yr to ~ 10200 yr for VSR2 and from ~ 1500 yr to ~ 9700 yr for VSR4. These values are large enough that only the first order spin-down is needed in the analysis [7]. In Tab. I some




FIG. 2. Number of sky patches in every 1 Hz band, from 20 Hz up to 128 Hz. The frequency on abscissa axis indicates the beginning frequency of each band. The number of patches increases with the square of the frequency, and is fixed by the highest frequency of each 1 Hz band.

quantities referring to the FFTs and peakmaps of VSR2 and VSR4 data sets are given. Tab. II contains a summary of the main parameters of the coarse step, among which the exact spin-down range considered for VSR2 and VSR4 analyses.

Run	T_{obs} [days]	T_{start} [mjd]	T_{FFT} [s]	N_{peaks} (after vetoes)
VSR2	185	55112	8192	191771835
VSR4	95	55762	8192	93896752

TABLE I. Some quantities referring to FFTs and peakmaps. For each run T_{obs} is the run duration, T_{start} is the run start epoch, T_{FFT} is the FFT time length, and N_{peaks} is the number of peaks in the peakmap, after applying all the vetoing procedures.

For a given sky position, the result of the FrequencyHough transform is a histogram in the signal frequency/spin-down plane, called the Hough map. The most significant candidates, i.e. the bins of the Hough map with the highest amplitude, are then selected using an effective way to avoid being blinded by particularly disturbed frequency bands (see Sec. VI and [7]). For each coarse (or *raw*) candidate a refined search, still based on the FrequencyHough, is run again on the neighborhood of the candidate parameters and the final *first-level* refined candidates are selected. The refinement results in a reduction of the digitalization effects, that is the sensitivity loss due to the use of a discrete grid in the parameter space. With regard to the frequency, which was already refined at the coarse step, no further over-resolution occurs. For the spin-down we have used an over-resolution factor $K_f = 6$, i.e., the coarse interval between the spin-down of each candidate and the next value (on both sides) is divided in 6 pieces. The refined search range includes

$2 K_f = 12$ bins on the left of the coarse original value, and $(2 K_f - 1) = 11$ bins on the right, so that two coarse bins are covered on both sides. This choice is dictated by the fact that the refinement is also done in parallel for the position of the source and, because of parameter correlation, a coarse candidate could be found with a refined spin-down value outside the original coarse bin. Using an over-resolution factor $K_f = 6$ is a compromise between the reduction of digitization effects and the increase of computational load.

The refinement in the sky position for every candidate is performed by using a rectangular region centered at the candidate coordinates. The distance between the estimated latitude (longitude) and the next latitude (longitude) point in the coarse grid is divided into $\hat{K}_{\text{sky}} = 5$ points, so that 25 sky points are considered in total, see discussion in Sec. IX-C of [7].

From a practical point of view, the incoherent step of the analysis has been done by splitting the full parameter space to be explored in several independent jobs. Each job covered a frequency band of 5 Hz, the full spin-down range, and a portion of the sky (the extent of which depended on the frequency band and was chosen to maintain balanced job durations). The full set of jobs was run on the European Grid Infrastructure (<http://www.egi.eu/>).

Candidates found in the analysis of VSR2 and VSR4 data are then *clustered*, grouping together those occupying nearby points in the parameter space. This is done to improve the computational efficiency of the next steps of the analysis. In order to significantly reduce the false alarm probability, *coincidences* are required among clusters of candidates obtained from the two data sets. The most significant coincident candidates are subject to a *follow-up* with greater coherence time, in order to con-

Run	δf_H [Hz]	N_f	$\delta \dot{f}$ [Hz/s]	N_{sd}	$\Delta \dot{f}$ [Hz/s]	τ_{min} [years]	N_{sky}
VSR2	$1.22 \cdot 10^{-5}$	8,847,360	$7.63 \cdot 10^{-12}$	15	$[-9.91, 1.52] \cdot 10^{-11}$	1600-10200	3,528,767
VSR4	$1.22 \cdot 10^{-5}$	8,847,360	$1.50 \cdot 10^{-11}$	8	$[-10.5, 1.50] \cdot 10^{-11}$	1500-9700	3,528,767

TABLE II. Summary of the main parameters for the coarse step of the analysis. δf_H is the frequency bin, N_f is the number of frequency bins in the analyzed band, $\delta \dot{f}$ is the spin-down bin, N_{sd} is the number of spin-down steps, $\Delta \dot{f}$ is the range of spin-down covered in the analysis, τ_{min} is the corresponding minimum spin-down age, N_{sky} is the total number of sky patches.

firm or discard them. Candidate selection and analysis are described in some detail in Sec. VI-VIII.

V. DATA CLEANING

Time and frequency domain disturbances in detector data affect the search and, if not properly removed, can significantly degrade the search sensitivity, in the worst case blinding the search at certain times or in certain frequency bands. The effects can vary depending on the nature and amplitude of the disturbance. As described in [7], we apply cleaning procedures to safely remove such disturbances or reduce their effect, without contaminating a possible CW signal. The disturbances can be catalogued as “time domain glitches”, which enhance the noise level of the detector in a wide frequency band; “spectral lines of constant frequency”, in most cases of known origin, like calibration lines or lines whose origin has been discovered by studying the behaviour of the detector and the surrounding environment, or “spectral wandering lines”, where the frequency of the disturbance changes in time (often of unknown origin and present only for a few days or even hours). Time-domain glitches are removed during the construction of the SFDB.

Spectral wandering lines and spectral lines of constant frequency are removed from the peakmaps [7]. Removal of spectral wandering lines (composed by peaks occurring at varying frequencies) is based on the construction of a histogram of low resolution (both in time and in frequency) peakmap entries, which we call the “gross histogram”. Based on a study on VSR2 and VSR4 data, we have chosen a time resolution $\Delta t_H = 12$ hours and a frequency resolution $\Delta f_H = 0.01$ Hz. In this way any true CW signal of plausible strength would be completely confined within one bin, but would not significantly contribute to the histogram, avoiding veto. As an example, Fig. 3 shows the time-frequency plot of the peaks removed by the “gross histogram” cleaning procedure on VSR2 and VSR4 data and Fig. 4 shows the histogram of the removed peaks, using a 10 mHz bin width, again both for VSR2 and VSR4 data.

A second veto, aimed at removing lines of constant frequency, is based on the “persistency” analysis of the peakmaps, defined as the ratio between the number of FFTs in which a given line was present and the total number of analyzed FFTs. The vetoing procedure consisted of histogramming the frequency bins and setting a

reasonable threshold to select lines to be removed. To evaluate the veto threshold, we have used a “robust” statistic, described in Appendix D of [7], which is based on the median rather than the mean, which is much less affected by tails in the distribution. The resulting threshold on the persistency is of the order of 10%. Figure 5 shows the lines of constant frequency vetoed on the basis of the persistency, given on the ordinate axis. In this way we have vetoed 710 lines for VSR2 and 1947 lines for VSR4, which we know to be more disturbed than VSR2. The “gross” histogram veto reduced the total number of peaks by 11.3% and 13.1% for VSR2 and VSR4 respectively. The persistency veto brings the fraction of removed peaks to 11.5 % and 13.5 % for VSR2 and VSR4 respectively. We end up with a total number of peaks which is 191,771,835 for VSR2 and 93,896,752 for VSR4. The use of the adaptivity in the FrequencyHough transform acts to reduce the effect of some of the remaining peaks, those corresponding to an average level of the noise higher than usual or to times when the detector-source relative position is unfavourable.

For testing purposes, ten simulated CW signals, called *hardware injections* (HIs), have been injected during VSR2 and VSR4 runs, by properly acting on the mirrors control system (see Sec. IX for more details). While we have developed a method to effectively remove HIs from the peakmap, it has not been applied for the present analysis as their presence allowed testing the whole analysis chain. We will show results from HIs analysis in Sec. IX.

VI. CANDIDATE SELECTION

As outlined in Sec. IV, after the FrequencyHough transform has been computed for a given dataset, the first level of refined candidates is selected. Their number is chosen as a compromise between a manageable size and an acceptable sensitivity loss, as explained in Sect. IX of [7]. Moreover, candidates are selected in a way to reduce as much as possible the effect of disturbances. Let us describe the selection procedure. The full frequency range is split into 1 Hz sub-bands, each of which is analyzed separately and independently. Following the reasoning in Sec. VIII of [7], we fixed the total number of candidates to be selected in each run to $N_{cand} = 10^8$. We distribute them in frequency by fixing their number in each 1 Hz band, $N_{cand,i}$, where $i = 20, \dots, 127$, in such a way to have the same expected number of coincidences in




FIG. 3. Time-frequency plot of the peaks removed by the “gross histogram” cleaning procedure for VSR2 (left) and VSR4 (right).




FIG. 4. Histogram of the peaks removed by the “gross histogram” veto for VSR2 (left) and VSR4 (right). The size of the bins in the histogram is 10 mHz.




FIG. 5. Lines of constant frequency vetoed on the basis of the persistency, shown in the ordinate axis, for VSR2 (left) and VSR4 (right). We have removed 710 lines for VSR2 and 1947 lines for VSR4.)

all the bands. Moreover, for each given frequency band we have decided to have the same expected number of coincidences for each sky cell. Then, the number of candidates to be selected for each 1 Hz band and for each sky cell is given by $N_{\text{cand},i}^{\text{sky}} = N_{\text{cand},i}/N_{\text{sky},i}$, where $N_{\text{sky},i}$ is the number of sky patches in the i th frequency band. Since $N_{\text{cand},i}$ linearly increases with the frequency (see Eq. 48 in [7]) and, for a given frequency band, the number of points in the sky increases with the square of the band maximum frequency, we have that the number of candidates selected per sky patch decreases with frequency.

We now focus on some practical aspects of the procedure. As previously explained, we have fixed the size of the frequency bands to 1 Hz, the total number of candidates to be selected in each of them ($N_{\text{cand},i}$) and the number of candidates in each cell of the sky ($N_{\text{cand},i}^{\text{sky}}$). We have now the problem of being affected by the presence of disturbances of unknown origin. They could still pollute sub-bands in the i th band, even after performing all the cleaning steps described in Sec.V. We have designed a procedure for candidate selection for this purpose. For each sky cell, we divide the i th band into $n_{\text{sb}} = N_{\text{cand},i}^{\text{sky}}$ sub-bands, and select the most significant candidate in each of them, i.e. the bin in the Hough map with the highest amplitude. In this way a uniform selection of candidates is done in every band and the contribution of possible large disturbances is strongly reduced. A further step consists of selecting the “second order” candidates. Once the most significant candidate in each sub-band has been selected, an empirically established exclusion region of ± 4 frequency bins around it is imposed. We now look for the second loudest candidate in that sub-band and select it only if it is well separated in frequency from the first one, by at least ± 8 frequency bins. In this way, we expect to select two candidates per sub-band in most cases and to have one candidate only when the loudest candidate is due to a big disturbance, or a particularly strong HI. Results are shown in Fig. 6 (left), which gives the number of candidates selected in every 1 Hz band and in Fig. 6 (right), which gives the number of candidates selected in every 1 Hz band and for each sky patch. In the end we have 194,457,048 candidates for VSR2 and 193,855,645 for VSR4, which means that we have selected also the “second order” candidates in $\sim 96\%$ of the cases.

VII. CANDIDATE CLUSTERING AND COINCIDENCE

We expect that nearly all the candidates selected in the analysis of a dataset will be false, arising from noise. In order to reduce the false alarm probability, coincidences among the two sets of candidates, found in VSR2 and VSR4 analysis, have been required. Indeed, given the persistent nature of CWs, a candidate in a dataset due to a real signal will (approximately) have the same pa-

rameters in another dataset, even if this covers a different time span. If a candidate is found to be coincident among the two datasets, it will be further analyzed as described in Sec. VIII.

In fact, due to computational reasons, candidates from each dataset are *clustered* before making coincidences. A cluster is a collection of candidates such that the distance d in the parameter space among any two is smaller than a threshold d_{clust} . We have used an empirically chosen value $d_{\text{clust}} = 2$. Each candidate is defined by a set of 4 parameters: position in ecliptic coordinates (λ, β) , frequency f and spin-down \dot{f} . Therefore, for two given candidates with $\vec{c}_1 = (\lambda_1, \beta_1, f_1, \dot{f}_1)$ and $\vec{c}_2 = (\lambda_2, \beta_2, f_2, \dot{f}_2)$ respectively, we define their distance as

$$d = \|\vec{c}_1 - \vec{c}_2\| = \sqrt{k_\lambda^2 + k_\beta^2 + k_f^2 + k_{\dot{f}}^2}. \quad (10)$$

Here $k_\lambda = |\lambda_2 - \lambda_1|/\delta\lambda$ is the difference in number of bins between the ecliptic longitudes of the two candidates, and $\delta\lambda = (\delta\lambda_1 + \delta\lambda_2)/2$ is the mean value of the width of the coarse bins in the ecliptic longitude for the two candidates (which can vary, as the resolution in longitude depends on the longitude itself), and similarly for the other terms. We find 94,153,784 clusters for VSR2 and 38,953,404 for VSR4.

Once candidates of both datasets have been clustered, candidate frequencies are referred to the same epoch, which is the beginning of VSR4 run. This means that the frequency of VSR2 candidate are shifted according to the corresponding spin-down value. Then coincidence requirements among clusters are imposed as follows. For each cluster of the first set, a check is performed on clusters in the second set. First, a cluster of the second set is discarded if its parameters are not compatible with the trial cluster in the first set. As an example, if the maximum frequency found in a cluster is smaller than the minimum frequency found in the other one, they cannot be coincident.

For each candidate of the first cluster of a potentially coincident pair, the distance from the candidates of the second cluster is computed. If a distance smaller than $d_{\text{coin}} = 2$ is found, then the two clusters are considered coincident: the distance between all the pairs of candidates of the two clusters is computed and the pair with the smallest distance constitutes a pair of coincident candidates. This means that each pair of coincident clusters produces just one pair of coincident candidates. The choice $d_{\text{coin}} = 2$, based on a study of software-injected signals, allows us to reduce the false alarm probability and is robust enough against the fact that true signals can be found with slightly different parameters in the two datasets. The total number of coincidences we found is 3,608,192.

At this point, coincident candidates are divided into bands of 0.1 Hz and subject to a *ranking* procedure in order to select the most significant ones. Let N be the number of coincident candidates in a given 0.1 Hz band.




FIG. 6. Number of candidates selected in every 1 Hz band (left) and in every 1 Hz band and in every sky patch (right). Asterisks (red in the online version) corresponds to VSR2, black circles to VSR4.

Let us order them in descending order of Hough map amplitude (i.e. from the highest to the smallest), separately for VSR2 and VSR4. We assign a rank to each of them, which is $1/N$ to the highest and 1 to the smallest. We have then two sets of ranks, one for VSR2 candidates and one for VSR4 candidates. Now we make the product of the ranks of coincident candidates. The smallest possible value is $1/N^2$, if a pair of coincident candidates has the highest Hough amplitude in both VSR2 and VSR4, while the largest possible value is 1, if a pair of coincident candidates has the smallest amplitude in both VSR2 and VSR4. For each 0.1 Hz band we select the candidates having the smallest rank product. We chose 0.1 Hz wide bands as a good compromise between having a statistic large enough in every band and reducing the effect of (strong) noise outliers in wider bands, as will be clear in the following. With this choice we have 1080 coincident candidates over the band 20-128 Hz.

At this point, among the ten candidates in ten consecutive 0.1 Hz bands we chose the most significant one, ending up with 108 candidates, one per 1 Hz band, which will be subject to a *follow-up* procedure, as described in the next section. The number of candidates selected at this stage depends on the amount of available computing resources and is constrained by the amount of time the analysis will take. Figure 7 shows the rank product for the final 108 candidates as a function of their frequency. The two smallest values corresponds to the hardware-injected signals at 52.8 Hz and 108.3 Hz. In Tab. VI the main parameters of the 108 selected candidates are given. They have been ordered as a function of the value of the ranking parameter, starting from the most significant one.

VIII. CANDIDATE FOLLOW-UP

The follow-up procedure consists of several steps applied to each of the 108 selected candidates. First, a fully

coherent step, using the candidate parameters, is done separately for the two runs. This is based on the same analysis pipeline used for targeted searches [1], [2], and the output is a down-sampled (at 1 Hz) time series where the Doppler effect, the spin-down and the Einstein delay for a source, having the same parameters as the candidate, have been corrected. A final cleaning stage is also applied, by removing the non-Gaussian tail of the data distribution.

From these data, a new set of longer FFTs is computed, from chunks of data of duration 81,920 seconds, which is ten times longer than in the initial step of the analysis. This procedure preserves true signals with parameters slightly different from those of the candidate, but within the uncertainty window because of the partial Doppler and spin-down correction just applied. If we assume that the true signal has a frequency different from that of the candidate by a coarse bin $1/T_{\text{FFT}}$, and a sky position also shifted by a coarse sky bin, the maximum allowed FFT duration can be numerically evaluated using Eq. 36 in [7], and is of about 290,000 seconds, significantly larger than our choice.

From the set of FFTs a new peakmap is also computed, selecting peaks above a threshold of $\sqrt{5.5} \simeq 2.34$ on the square root of the equalized spectra, significantly higher than that (1.58) initially used (see Sec. IV). This is justified by the fact that a signal present in the data, and strong enough to produce a candidate, would contribute similarly when lengthening the FFT (by a factor of 10) while the noise in a frequency bin decreases by a factor of 10 (in energy). In fact, a threshold of 2.34 is a very conservative choice which, nevertheless, reduces the expected number of noise peaks by a factor of 20 with respect to the initial threshold.

At this point the peakmaps computed for the two runs are combined, so that their peaks cover a total observation time of 788.67 days, from the beginning of VSR2 to the end of VSR4 run. Before applying the incoherent step, a grid is built around the current candidate.




FIG. 7. Rank product for the final 108 candidates, which will be subject to the follow-up procedure. The two candidates with smallest rank product, indicated by filled circles, correspond to two hardware-injected signals.

The frequency grid is built with an over-resolution factor of 10, as initially done, so that $\delta f = 1/81920 \times 10 = 1.22 \times 10^{-6}$ Hz, and covers 0.1 Hz around the candidate frequency. The grid on the spin-down is computed with a step 10 times smaller than the coarse step of the shorter run (VSR4), and covering ± 1 coarse step, so that 21 spin-down bins are considered. Finally, a grid of 41 by 41 sky points (1681 in total) is built around the candidate position, covering ± 0.75 times the dimension of the coarse patch (whose extension actually depends on the position in the sky and on the candidate frequency). For every sky point the peakmap is Doppler corrected by shifting each peak by an amount corresponding to the Doppler shift, and a FrequencyHough transform is computed. This means that for each candidate $41 \times 41 = 1681$ FrequencyHough transforms are computed. The loudest candidate, among the full set of FrequencyHough maps, is then selected.

The starting peakmap is now corrected using the parameters of the loudest candidate and projected on the frequency axis. On the resulting histogram we search for significant peaks. This is done by taking the maximum of the histogram over a search band covering a range of ± 2 coarse bins around the candidate frequency. The rest of the 0.1 Hz band is used to estimate the background noise: it is similarly divided in sub-intervals covering 4 coarse bins and the loudest candidate is taken in each of them. In total we have 408 sub-intervals. In the frequentist approach the significance of a candidate is measured by the p-value, that is the probability that a candidate as significant as that, or more, is due to noise. The smaller is the computed p-value and less consistent is the candidate with noise. We estimate a p-value associated to

the candidate by computing the fraction of sub-bands in which the loudest peak is larger than that in the search band. The smaller is the p-value, the higher is the candidate significance. If the p-value is smaller than 1% the candidate can be considered as “interesting”, namely it is not fully compatible with noise, and will be subject to a deeper scrutiny, looking at the consistency of the candidate between the two runs and, possibly, extending the analysis to other data sets or applying a further step of follow-up with a longer coherence time.

As an example, Fig. 8 shows the peakmap histogram for the candidate at frequency 121.81 Hz, in which no significant peak is found around the candidate frequency (identified by the vertical dashed line). The corresponding p-value is ~ 0.44 .

In contrast, Fig. 10 shows the peakmap histograms for the candidates at 52.80 Hz (left), corresponding to HI *pulsar_5*, and the candidate at 108.85 Hz (right), corresponding to HI *pulsar_3*. In each case a highly significant peak is visible at the frequency of the candidate (in fact it is the largest peak in the whole 0.1 Hz band around it), as detailed in Sec. IX.

The p-value for the 108 candidates is given in the last column of Tab. VI. In Fig. 9 the p-value for the 108 candidates is shown as a function of candidate frequency. Both the hardware injections have the lowest possible p-value (equal to $1/408 = 2.45 \times 10^{-3}$), that is the highest possible significance, providing a good check that the full analysis pipeline works correctly.

Moreover, there are three outliers, which are candidates not associated to injected signals, having a p-value very close or below the 1% threshold we have established. These three candidates have undergone further analysis




FIG. 8. Peakmap histogram around the candidate at frequency 121.81 Hz. The vertical line identifies the candidate frequency.




FIG. 9. Final p-values for the 108 candidates as a function of candidate frequency after performing the follow-up step. The minimum value is $1/408=0.00245$. The two filled circles identify hardware-injected signals.

in order to increase their significance or to discard them as possible GW signals. In fact, as described in Appendix , all of them appear to be incompatible with GW signals.

IX. HARDWARE INJECTION RECOVERY

Ten simulated CW signals have been continuously injected in the Virgo detector during VSR2 and VSR4 runs, by actuating on the mirror control system with the aim of testing analysis pipelines. Two of these *hardware* injections have frequency below 128 Hz and have been considered in this work: *pulsar_5*, with a frequency of about 52.8 Hz, and *pulsar_3*, at about 108.85 Hz. In Tab. III

parameters relevant for the current analysis are given. These signals are strong enough to appear as the most

HI	f_{gw} [Hz]	\dot{f} [Hz/s]	λ [deg]	β [deg]	H_0 $\times 10^{-24}$
<i>pulsar_5</i>	52.80832	-4.03×10^{-18}	276.8964	-61.1909	3.703
<i>pulsar_3</i>	108.85716	-1.46×10^{-17}	193.3162	-30.9956	8.296

TABLE III. Main parameters of the HIs in the band of the present analysis. The reference epoch for the frequency is MJD 52944.

significant candidates in the final list (Tab. VI). As such they have been also subject to the follow-up procedure.

In Fig. 10 we show the final peakmap histograms for the candidate corresponding to *pulsar_5* (left) and *pulsar_3* (right), where in both cases a very high peak is clearly visible in correspondence of the signal frequency. This is reflected in the high significance level shown in Tab. VI. Tab. IV contains the estimated parameters for the HIs, together with the associated error, which is the difference with respect to the injected values. For both signals the parameters are well recovered with an improved accuracy thanks to the followup.

X. RESULTS

As all the 106 non-injection candidates appear to be consistent with what we would expect from noise only, we proceed to compute an upper limit on the signal amplitude in each 1-Hz band between 20 Hz and 128 Hz, with the exception of the band 52-53 Hz and 108-109 Hz, which corresponds to the two HIs (in fact, see point 2. in this section, there are also several noisy sub-bands which have been excluded *a posteriori* from the computation of the upper limit). We follow a standard approach adopted for all-sky CW searches [4], [6] and detailed in the following. Many (of the order of 200,000 in our case) simulated signals are injected into the data, and the signal amplitude is determined such that 90 % of the signals (or the desired confidence level) are detected, and are more significant than the original candidate found in the actual analysis of the same frequency band.

In other words, for each 1-Hz band we generate 100 simulated signals in the time domain, with unitary amplitude and having frequency and the other parameters drawn from a uniform distribution: the spin-down within the search range, position over the whole sky, polarization angle ψ between $-\frac{\pi}{2}$ and $\frac{\pi}{2}$, and inclination of the rotation axis with respect to the line of observation, $\cos \iota$, between -1 and 1 . The time series containing the signals are then converted in a set of FFTs, with duration 8192 seconds, the same used for the analysis, and covering the band of 1 Hz. These “fake signal FFTs” are multiplied by an amplitude factor and summed to the original data FFTs. This is repeated using of the order of 10-15 different amplitudes, chosen in order to be around the expected 90% confidence level upper limit.

For each set of 100 injections an analysis is done using the all-sky search code over a reduced parameter space around each injection, consisting of a frequency band of ± 0.2 Hz around the injection frequency, the same spin-down range used in the production analysis, and 9 sky points around the coarse grid point nearest to the injection position. We have verified, by injecting software simulated signals into Virgo VSR2 and VSR4 data, that for signal amplitudes around the approximately expected upper limit values this volume is sufficiently large to contain all the candidates produced by a given signal, and at the same time is small enough to make the procedure reasonably fast. The output of this stage is, for each in-

jected signal, a set of candidates from VSR2 data and another set of candidates from VSR4. In the production analysis, candidates of each run are clustered before making coincidences. For the computation of the upper limit the clustering cannot be applied because of the density of signals which would strongly affect the cluster composition introducing a mixing of candidates belonging to different injections. We recall that the candidate clustering has been introduced in the production analysis with the main purpose to reduce the computational cost of the coincidence step. However, for the upper limit computation, a direct coincidence step among candidates is still affordable. Hence, for every candidate found in VSR2 we determine, if it exists, the nearest coincident candidate found in VSR4 with a distance smaller than 2. The coincident candidates are then ranked using the same procedure described in Sec. VII, and the most significant one is selected. Then, for each injected signal amplitude we have at most 100 coincident candidates. They are compared to the candidate found in the actual analysis in the same 1-Hz band, in order to count the fraction “louder” than that. Two issues arise here and must be properly addressed.

1. In principle, the comparison between the candidates found after injections and the actual analysis candidate should be done on the basis of their ranking. In fact this cannot be done because the numerical values of the ranking depend on the number of candidates present in the 1-Hz band, and this number is very different in the two cases, as the injection analysis is performed over a smaller portion of the parameter space. We have then decided to make the comparison using the Hough amplitudes of the candidates: a candidate found in the injection analysis is considered “louder” if the amplitudes of its parent candidates, found in VSR2 and VSR4 data, are both larger than the corresponding amplitudes of the parents of the actual analysis candidate. This will tend to slightly overestimate the upper limit.
2. We have verified that in some bands the upper limit can be heavily affected by the presence of gaps in the starting peakmap, namely frequency bands in which the number of peaks is significantly smaller than in the rest of the band. As an example, in Fig. 11 the peakmap peak distribution is shown for a “bad” frequency band between 35 and 36 Hz in VSR2 data, containing gaps, and for a good one, between 126 and 127 Hz. The gaps are the result of the various cleaning steps applied to the data, and the injections that, for a given amplitude, have frequency overlapping with a gap will not be able, in most cases, to produce detectable candidates because the amplitude in the Hough map near the signal frequency is reduced by the low noise contribution. As a consequence, the upper limit tends to be significantly worse with respect to the case in




FIG. 10. Peakmap histogram around the hardware injections *pulsar_5* (left) and *pulsar_3* (right). In both cases the highest peak corresponds to the signal frequency, with high accuracy.

HI	\hat{f} [Hz]	Δf	$\hat{\dot{f}}$ [Hz/s]	$\Delta \dot{f}$	$\hat{\lambda}$ [deg]	$\Delta \lambda$	$\hat{\beta}$ [deg]	$\Delta \beta$
<i>pulsar_5</i>	52.80830	-1.8	0.0	4×10^{-6}	276.93	0.12	-61.19	0.007
<i>pulsar_3</i>	108.85714	-1.2	1.18×10^{-12}	1.05	193.36	-0.05	-31.00	0.66

TABLE IV. Estimated parameters, and corresponding errors, for candidates corresponding to hardware injections present in the band between 20 Hz and 128 Hz. The parameters $\hat{f}, \hat{\dot{f}}, \hat{\lambda}, \hat{\beta}$ are the estimated frequency, spin-down, ecliptic coordinates; $\Delta f, \Delta \dot{f}, \Delta \lambda, \Delta \beta$ are the differences with respect to the injected parameter values in units of the corresponding bins.

which the gaps are not relevant. To cope with this issue we opted to exclude from the computation of the upper limit the parts of a 1-Hz band that correspond to gaps. This procedure excludes the injections that happen to lie within the excluded region. In this way we obtain better results, which are valid only in a sub-interval of the 1-Hz band. In Appendix C we report all the intervals excluded from the upper limit computations. In total they cover about 8.6 Hz (out of the 108 Hz analyzed).

For each 1-Hz band, and for each injected signal amplitude, once we have computed the fraction of candidates louder than the actual analysis candidate, the pair of amplitudes that are nearest, respectively from above and from below, to the 90% confidence level are used as starting point of a new round, in which a new set of signals with amplitudes between those values are injected until the final confidence level of 90% is reached. Overall, of the order of 2×10^5 signals have been injected in both VSR2 and VSR4 data. The final accuracy of the upper limit depends on the number of injections actually used and on the quality of the data in the considered 1-Hz band. By repeating of the order of 10 times the upper limit computation in a few selected bands, we have verified that the upper limits have an accuracy better than $\sim 5\%$ for “bad” bands and better than about 1% for “good” bands, in any case smaller than the amplitude uncertainty due to the data calibration error (Sec. III).

In Fig. 12 the upper limits on signal strain as a function of the frequency are plotted. The values that are

valid on just a portion of the corresponding 1-Hz band are labeled by a filled circle. A rough comparison with other all-sky searches result shows that our upper limits are better than those established, for instance, by the Einstein@Home pipeline on LIGO S5 data [4] for frequencies below ~ 80 Hz (however, in [4] upper limits are computed every 0.5 Hz, instead of 1 Hz). A useful plot to understand the astrophysical reach of the search is shown in Fig. 13. The various sets of points give the relation between the signal frequency derivative and the signal frequency for sources detectable at various distances, assuming their spin-down to be due solely to the emission of gravitational radiation (these types of sources are also referred to as *gravitars* [28]). For instance, considering a source at a distance of 100 parsecs and emitting a CW signal with a frequency of 80 Hz, it would be detectable if its frequency derivative was larger, in modulus, than about 2.8×10^{-12} Hz/s. On the same plot curves of constant ellipticity are also shown. The above source example would have an ellipticity larger than about 2×10^{-5} . From this plot we can conclude that our search would have been sensitive enough to detect all the gravitars within about 400 parsecs, emitting a signal with frequency above about 60 Hz and with spin-down age larger than about 4500 years. Also, all the gravitars within about 50 parsecs, emitting a signal with frequency above 20 Hz and with spin-down age larger than about 1500 years would have been detected.




FIG. 11. Peakmap peak distribution for two bands in VSR2 data. The left plot refers to the band 35-36 Hz, with gaps clearly visible. The right plot refers to a much cleaner band, ranging from 126 Hz to 127 Hz.




FIG. 12. Joint VSR2-VSR4 upper limit on dimensionless strain as a function of the frequency. Open circles refer to upper limits values valid over the full corresponding 1-Hz band, while filled circles refer to upper limit values valid only in a portion of the corresponding 1-Hz band.

XI. CONCLUSIONS

In this paper we have described a low-frequency all-sky search for CW signals, performed using data from the Virgo VSR2 and VSR4 runs. This is the first all-sky search for isolated stars with GW frequencies below 50 Hz. We have applied a hierarchical procedure, in which the incoherent step is based on the FrequencyHough transform. Particular care has been placed on cleaning the data of known and unknown instrumental disturbances, before selecting potentially interesting CW candidates. The criteria used to select candidates have been designed in such a way to avoid sensitivity to any single noise disturbance. For each candidate surviving a coincidence step, a follow-up procedure has been applied in order to reject or confirm it. Three candidates have been found to be most significant, but their potential GW

origin has been ruled out after a deeper analysis. Having found no evidence for CW signals, a 90% confidence level upper limit on signal strain has been computed over the full frequency band. Our upper limit improves upon prior results below ~ 80 Hz and the astrophysical reach of the search is interesting. Our search would have been able to detect all the *gravitars* within about 400 parsecs from the Earth, spinning at frequencies above 30 Hz and with spin-down age larger than about 4500 years. It would have been also able to detect all the *gravitars* within about 50 parsecs from the Earth, spinning at frequencies above 10 Hz and with spin-down age larger than about 1500 years. This analysis pipeline will be applied to data of Advanced LIGO and Virgo detectors. Given their improved sensitivities, the chance to detect CW signals by spinning neutron stars will be significantly better. Detection of such signals would provide insights into the

cont. from prev. page						
Order number	Frequency [Hz]	Spin-down $\times 10^{-11}$ [Hz/s]	λ [deg]	β [deg]	Rank $\times 10^{-4}$	Significance
13	110.8583	-5.484	332.6659	-37.837	0.008547	0.97310
14	55.8628	-8.2608	129.375	-51.6569	0.0093692	0.88753
15	123.3888	-2.9019	208.9888	-31.6621	0.009537	0.50855
16	68.9449	0.57005	186.5033	41.9996	0.0097457	0.09512
17	56.0031	-2.5206	335.2227	-38.9303	0.010498	0.21112
18	87.4609	-4.3538	252.5236	55.405	0.011137	0.22346
19	125.4042	1.7658	121.5085	18.4552	0.011196	0.65921
20	33.0337	-10.4675	31.4954	-49.4705	0.012961	0.55307
21	124.9926	-2.7093	240.8653	-20.0556	0.013249	0.31822
22	95.3279	-3.0965	209.6354	46.4176	0.013873	0.29095
23	67.0831	-8.7653	149.2214	-70.1448	0.014011	0.52417
24	119.6512	-7.3829	111.728	-55.7048	0.015368	0.76410
25	43.2998	-0.50451	238.3516	88.2667	0.016197	$< 2.45 \cdot 10^{-3}$
26	127.8655	-4.2248	135.7367	37.1047	0.016758	
27	114.0429	0.88388	270.2045	73.8248	0.016903	
28	76.2553	-4.2208	228.0401	45.2728	0.016994	0.37804
29	28.5502	-9.8955	108.4932	-63.7331	0.017029	0.21229
30	35.6011	-0.44095	221.1306	59.8507	0.017121	0.14525
31	61.1424	-7.3134	327.0508	45.8359	0.017213	0.97660
32	99.9012	-0.6336	212.2641	-17.2635	0.018999	0.33918
33	107.2998	1.2632	159.1845	-43.6778	0.019099	0.74695
34	113.7034	0.50451	178.7906	-48.3726	0.019143	0.74301
35	117.7723	-2.7073	80.7946	10.7556	0.023097	0.07017
36	47.851	-5.1722	27.665	-39.351	0.024202	0.46368
37	82.1015	1.2632	126.6622	-49.7879	0.02424	0.12849
38	86.6052	-10.0882	177.278	30.7229	0.02499	0.36871
39	83.0785	-10.4695	342.531	-32.6498	0.025006	0.70731
40	65.4443	-1.0726	304.134	-19.8682	0.025727	0.16759
41	96.2287	-7.6888	166.1909	22.8864	0.026087	0.67039
42	25.0585	-2.3378	241.7518	-18.2543	0.027399	0.47953
43	34.1145	-5.7998	260.6087	61.9021	0.028246	0.88333
44	75.7372	-10.0901	112.5	23.0135	0.029795	0.09669
45	92.1503	0.13108	172.16	63.8249	0.032852	0.42105
46	51.5143	2.024	29.8065	-50.9461	0.033757	0.10526
47	98.9983	-6.8724	315.3778	50.3757	0.035169	0.268949
48	109.1544	2.3954	321.2877	-15.2403	0.035476	0.29516
49	78.7986	-2.5206	44.8338	28.9471	0.035727	0.08720
50	81.1001	-3.0906	310.0948	-22.2304	0.035921	0.37222
51	88.2414	-4.2267	336.0401	-31.4333	0.036207	0.27933
52	71.2004	1.7678	162.6366	-55.2736	0.036235	0.01222
53	120.7367	0.94744	86.9381	-54.5046	0.036489	0.82222
54	63.8707	-4.7273	209.8977	46.7812	0.036588	0.44134
55	60.0709	-7.2538	190.8441	69.5008	0.036653	0.24022
56	48.5878	-3.2852	329.1029	44.5078	0.037649	0.42458
57	85.8599	-4.0361	355.2824	44.4263	0.039416	0.27374
58	72.5912	-9.6472	21.228	34.9257	0.039794	0.70244
59	69.598	-3.2157	309.3422	-24.3772	0.039822	0.90220
60	91.7067	-8.1297	180.7389	-35.0032	0.040292	0.66081
61	23.3888	-9.3274	254.1493	59.3324	0.040735	0.35195
62	41.9977	-10.1537	38.3415	-58.7027	0.041007	0.58343
63	106.3632	-10.6562	269.7816	-9.1791	0.041596	0.32824
64	112.503	-6.3719	223.3497	-15.9813	0.041966	0.03307
65	45.6924	-9.0176	40.9655	36.3556	0.044055	0.97206
66	36.2578	-9.3294	35.7818	37.7206	0.045444	0.93854
67	44.3556	-10.2788	109.623	14.9484	0.046831	0.42458
68	54.0477	-3.907	309.394	62.1078	0.046967	0.53801
69	30.8852	-1.2593	236.7692	51.7512	0.049419	0.01117
70	74.1304	-0.62964	330.9153	-31.5795	0.05038	0.59657
71	46.9325	-1.3864	218.8421	81.1703	0.051493	0.40350
72	53.1366	-5.488	342.5228	-34.7568	0.052257	0.29516
73	59.0207	1.9564	248.6587	-1.5716	0.053051	0.57457
cont. on next page						

cont. from prev. page						
Order number	Frequency [Hz]	Spin-down $\times 10^{-11}$ [Hz/s]	λ [deg]	β [deg]	Rank $\times 10^{-4}$	Significance
74	24.0609	-10.4715	108.5902	-64.411	0.053547	0.03352
75	122.1853	-4.4134	325.8941	48.4439	0.057407	0.50279
76	94.0253	0.44293	100.3213	10.2366	0.057537	0.21119
77	104.0544	-9.4585	96.6947	9.8744	0.057864	0.93048
78	105.7558	-8.8904	241.7306	-31.8533	0.058097	0.63636
79	80.5196	-6.0541	340.1432	45.3848	0.060008	0.21229
80	21.3803	-10.3404	324.3529	-34.1692	0.062657	0.22905
81	66.6324	-0.25423	124.8241	-57.8314	0.062913	0.29584
82	102.2397	-6.4275	358.6386	47.3049	0.063841	0.42458
83	126.5903	-10.8429	188.3702	51.0238	0.065495	0.05022
84	90.2319	-3.0906	324.5108	-24.7639	0.066156	0.69273
85	101.9708	-2.2703	19.0030	-61.613	0.06835	0.92178
86	73.5608	-0.50451	286.4738	-12.1638	0.069204	0.61124
87	64.0628	-4.475	246.9913	-16.5005	0.069771	0.81564
88	27.3621	-3.2157	228.9322	48.5627	0.070012	0.39766
89	58.5039	-9.7108	131.6129	-55.3565	0.070837	0.67039
90	84.5958	-2.3338	103.5869	-51.7004	0.072562	0.32960
91	20.2353	-3.9725	87.1304	-55.6842	0.075261	0.42222
92	37.7963	-4.0976	178.6723	44.0613	0.078217	0.30726
93	31.032	-4.6657	342.9863	-35.3718	0.081675	0.02545
94	29.902	-8.0741	243.7377	69.6141	0.086694	0.53216
95	111.1327	-6.1137	23.9462	27.6296	0.087627	0.89821
96	39.6152	-3.5931	229.8023	42.5234	0.087633	0.41899
97	57.1775	-8.952	22.4698	25.4575	0.095732	0.32163
98	70.0715	-7.7543	231.3566	43.4535	0.097824	0.46943
99	42.9996	-0.31977	72.6974	14.6357	0.098683	0.25917
100	49.1547	0.065538	286.3941	-14.3416	0.10996	0.67597
101	62.2854	-6.5625	155.5203	-86.2875	0.11363	0.07262
102	38.8918	-7.3769	229.5254	-17.7099	0.11676	0.44134
103	93.0512	-3.3409	297.0175	-21.1575	0.12164	0.40782
104	40.4971	-1.3268	123.7959	31.1944	0.13931	0.48603
105	79.9665	-11.3455	230.6667	-10.579	0.15052	0.46368
106	22.5768	-9.391	357.9813	33.7066	0.16427	0.38596
107	89.1945	-9.0811	278.9743	51.5108	0.16766	0.23463
108	50.8034	0.37739	84.3333	-40.3263	0.26302	0.92441

Appendix B: Outliers analysis

In this section we describe the further analysis steps applied to the three outliers found in the follow-up step (see Sec. VIII). The goal here is to reject or confirm each of them as potential gravitational wave signal candidates. The main parameters of the three outliers are listed in Tab. VI.

1. Outlier at 30.88 Hz

This candidate has a significance just above the 1% threshold. An important verification step consists in checking if the “signal characteristics” are consistent among the two detectors. For this purpose we have repeated the last part of the follow-up procedure (from the FrequencyHough stage on) separately for the two runs. The result is shown in Fig. 14, where the final peakmap histograms are plotted both for the two runs separately and for the joint analysis. We clearly see that the outlier is strong in VSR4 data but not in VSR2 data. Given the

Significance	f [Hz]	\dot{f} [Hz/s]	λ [deg]	β [deg]
0.0114	30.88541	-9.19×10^{-12}	235.40	50.99
<0.0024	43.29974	-5.04×10^{-12}	198.30	89.42
0.0112	71.20035	1.13×10^{-11}	162.83	-55.27

TABLE VI. Main parameters of the three outliers found in the follow-up. The significance is given in the second column; f and \dot{f} are the candidate frequency and spin-down; λ and β is the candidate position in ecliptic coordinates. They correspond, respectively, to candidates number 69, 25 and 52 in Tab. . Note that the reported parameter values are those after the follow-up step and then are slightly different from those in Tab. which, on the contrary, have been computed before the follow-up.

characteristics of the two runs this is not what we expect for a real GW signal. The expected ratio between the peakmap histogram heights for the two data runs can be estimated for nominal signal strengths [7] and is about 1.5. This is much lower than the observed ratio, which is about 13. The inconsistency of the apparent signals

in VSR2 and VSR4 is also indicated by the lower height of the joint-run peak map histogram in Fig. 14 with respect to that of VSR4 alone. This is confirmed by the Hough map, shown in Fig. 15 (left), where a single and wide stripe is clearly visible, suggesting that the outlier is produced by a rather short duration disturbance present in VSR4 data only (for comparison, Fig. 3 in [7] shows the Hough map for an HI in VSR2 data). This seems to be confirmed looking at the peakmap around the candidate [see Fig. 15 (right)], where a high concentration of peaks, of unidentified origin, is present at the beginning of VSR4, lasting for about 10 days. From this analysis we think the outlier can be safely ruled out as being inconsistent with a real GW signal.

2. Outlier at 43.30 Hz

This candidate has a very high significance (corresponding to the smallest possible p-value). In Fig. 16 the peakmap histogram for the full analysis, together to those for the single runs are shown. We see that basically the “signal” is present mainly in one of the two runs. This is also what we conclude looking at the Hough map and the peakmap in Fig. 17, where the main contribution coming from VSR2 is clearly visible. From the previous considerations it seems that also this outlier is associated to some disturbance in the data. In particular it happens in a frequency region (between 40 and 45 Hz) which is heavily polluted by noise produced by rack cooling fans. It must be noticed that outliers clearly associated to a disturbed region found at this step of the procedure should not be surprising. In fact, despite all the cleaning procedures and criteria used to select candidates, some instrumental disturbance can be still present into the data.

3. Outlier at 71.20 Hz

This third outlier has many characteristics in common with that at 30.88 Hz. Comparing the peakmap histograms of Figs. 14 and 18, we see again that it is mainly present in VSR4 data. In this case the expected ratio between the peakmap histogram heights, in presence of a signal, is about 0.7 while we observe a ratio of about 3. Moreover, as for the outlier at 30.88 Hz, the height of the joint-run peak map histogram is smaller than for VSR4 alone, which is also inconsistent with the signal hypothesis. This is confirmed by the Hough map shown in Fig. 19 (left), where basically only a wide stripe in the frequency/spin-down plane is visible, indicating that the outlier comes mainly from only one of the two runs. A further confirmation comes from the peakmap in Fig. 19 (right), in which a high concentration of peaks is visible at the right frequencies just at the beginning of VSR4, similarly to what happens for the first outlier. We conclude that this outlier is not due to a real GW signal.

Appendix C: Excluded bands

In the following table the noisy frequency intervals excluded from the computation of the upper limits are given. The two intervals 52-53 Hz and 108-109 Hz have been excluded because their most significant candidate is an hardware injected signal.




FIG. 14. Final peakmap histograms for the outliers at 30.88 Hz. The dark thick curve corresponds to the full VSR2/VSR4 analysis; the thin dashed (blue in color version) curve corresponds to VSR2 analysis while the thin continuous (red in color version) curve corresponds to the analysis of VSR4 alone.




FIG. 15. Hough map (left) and peakmap (right) for the outlier at 30.88 Hz. The right plot shows the combination of the peakmaps from VSR2 and VSR4 data. The empty region in the middle corresponds to the time separation between the end of VSR2 and the beginning of VSR4. The outlier at 30.88 Hz is due to the concentration of peaks clearly visible at the beginning of VSR4.

-
- [1] J. Abadie *et al.*, The Astrophysical Journal **737**, 93 (2011)
 - [2] J. Aasi *et al.*, The Astrophysical Journal **785**, 119 (2014)
 - [3] B. Abbott *et al.*, Phys. Rev. Lett. **102**, 111102 (2009)
 - [4] J. Aasi *et al.*, Phys. Rev. D **87**, 042001 (2013)
 - [5] J. Aasi *et al.*, Classical Quantum Gravity **31**, 165014 (2014)
 - [6] J. Aasi *et al.*, Classical Quantum Gravity **31**, 085014 (2014)
 - [7] P. Astone, A. Colla, S. D'Antonio, S. Frasca, and C. Palomba, Physical Review D **90**, 042002 (2014)
 - [8] A. Noutsos *et al.*, MNRAS **430**, 2281 (2013)
 - [9] A. P. Igoshev and S. B. Popov, MNRAS **432**, 967 (2013)
 - [10] P. Astone, S. D'Antonio, S. Frasca, and C. Palomba, Class. Q. Grav. **27**, 194016 (2010)
 - [11] P. Jaranowski, A. Krolak, and B. F. Schutz, Physical Review D **58**, 063001 (1998)
 - [12] C. J. Horowitz and K. Kadau, Phys. Rev. Lett. **102**, 191102 (2009)
 - [13] K. Hoffman and J. Heyl, MNRAS **426**, 2404 (2012)
 - [14] N. K. Johnson-McDaniel and B. J. Owen, Phys. Rev. D **88**, 044004 (2013)
 - [15] R. Ciolfi, V. Ferrari, and L. Gualtieri, MNRAS **406**, 2540 (2010)
 - [16] B. Abbott *et al.*, Rep. Prog. Phys. **72**, 076901 (2009)
 - [17] T. Accadia *et al.*, J. Instrum. **7**, P03012 (2012)

Band [Hz]	Excluded range [Hz]
20 - 21	[20,20.17505]
22 - 23	[22.34997,22.43506]
23 - 24	[23.68359,23.77637]
25 - 26	[25.10864,25.15381], [25.15967,25.24634]
26 - 27	[26.41052,26.43799]
28 - 29	[28.78516,28.82019]
29 - 30	[29.08484,29.09985], [29.10010,29.13440]
30 - 31	[30.82519,30.85510]
34 - 35	[34.54517,34.58508]
35 - 36	[34.99304,35.05713], [35.27398,35.30729], [35.63635,35.78503]
37 - 38	[37.05517,37.15515], [37.27429,37.31531], [37.38379,37.39734] [37.50513,37.62573], [37.70520,37.77563], [37.96509,37.99988]
38 - 39	[38,38.02795], [38.02820,38.04565], [38.16516,38.20740]
39 - 40	[39.95520,39.99988]
40 - 41	[40,40.07568]
41 - 42	[41.10522,41.13501], [41.50329,41.53833], [41.92456,41.98804]
42 - 43	[42.01514,42.04700], [42.67517,42.76758], [42.80493,42.85730]
43 - 44	[43.22302,43.30041], [43.30359,43.33606], [43.69189,43.71558] [43.85803,43.93787]
45 - 46	[45.96863,45.99939]
46 - 47	[46.00024,46.08508], [46.29724,46.31506], [46.36523,46.41504] [46.60522,46.69507], [46.81738, 46.89501]
47 - 48	[47,47.02551], [47.06518,47.10510]
48 - 49	[48.83484,48.87512]
49 - 50	[49.38806,49.403564], [49.40588,49.62561], [49.67956,49.75683] [49.83520,49.95080], [49.95105,49.99890]
50 - 51	[50,50.05041], [50.05066,50.07971], [50.07995,50.27893] [50.29407,50.60705]
51 - 52	[51.01477,51.07690], [51.18506,51.22534], [51.38513,51.41504] [51.59521,51.63525]
52 - 53	[52.0,53.0]
56 - 57	[56.13452,56.15942]
60 - 61	[60.25024,60.27502], [60.81225,60.82788]
61 - 62	[61.32324,61.35095], [61.47375,61.49219], [61.50574,61.51977] [61.52026,61.58581], [61.61938,61.63513], [61.67663,61.71472] [61.71496,61.97717]
62 - 63	[62,62.25500], [62.29517,62.31323], [62.31396,62.32690] [62.38623,62.47876], [62.48059,62.52502]
63 - 64	[63.19714,63.22656], [63.75146,63.77502]
82 - 83	[82.22692,82.24511]
84 - 85	[84.81482,84.89502]
90 - 91	[90.49523,90.52685]
92 - 93	[92.00756,92.02527], [92.02746,92.07629], [92.264523,92.27991] [92.30432,92.32837], [92.32861,92.36609], [92.39587,92.47815] [92.49524,92.73511], [92.93469,92.97558]
99 - 100	[99.92712,99.97461]
100 - 101	[100,100.04614], [100.21228,100.23535]
102 - 103	[102.78369,102.80505]
108 - 109	[108.0 109.0]
111 - 112	[111.76636,111.80127], [111.81116,111.87512]
112 - 113	[112.47681,112.49194], [112.50500,112.51843], [112.52331,112.59997] [112.61475,112.65307], [112.67578,112.83032], [112.83374,112.87512]
113 - 114	[113.00012,113.03137], [113.04187,113.08520], [113.11633,113.14355] [113.17065,113.28479], [113.28552,113.38379], [113.38513,113.47986] [113.48010,113.52502], [113.59570,113.62610], [113.63488,113.66504]
114 - 115	[114.11096,114.15332], [114.15515,114.22534]
127 - 128	[127.98523,127.99841]

TABLE VII. Frequency intervals excluded in the computation of the upper limit. The total vetoed frequency band amounts to about 8.6 Hz.




FIG. 16. Final peakmap histograms for the outliers at 43.30 Hz. The dark thick curve corresponds to the full VSR2/VSR4 analysis; the light continuous (red in color version) curve corresponds to VSR4 analysis while the light dashed (blue in color version) curve corresponds to the analysis of VSR2 alone.




FIG. 17. Hough map (left) and peakmap (right) for the outlier at 43.30 Hz.

- [18] H. Grote (The LIGO Scientific Collaboration), *Class. Quantum Grav.* **27**, 084003 (2010)
- [19] T. Accadia *et al.* (The Virgo Collaboration), *Class. Quantum Grav.* **28**, 025005 (2011)
- [20] L. Rolland, Virgo note VIR-0703A-11(2011)
- [21] P. R. Brady and T. Creighton, *Physical Review D* **61**, 082001 (2000)
- [22] S. Frasca, P. Astone, and C. Palomba, *Classical & Quantum Gravity* **22**, S1013 (2005)
- [23] P. Astone, S. Frasca, and C. Palomba, *Classical & Quantum Gravity* **22**, S1197 (2005)
- [24] F. Acernese *et al.*, *Classical & Quantum Gravity* **26**, 204002 (2009)
- [25] F. Antonucci *et al.*, *Classical & Quantum Gravity* **25**, 184015 (2008)
- [26] C. Palomba, P. Astone, and S. Frasca, *Classical & Quantum Gravity* **22**, S1255 (2005)
- [27] A. Sintes and B. Krishnan, *Journal of Physics: Conference Series* **32**, 206 (2006)
- [28] C. Palomba, *MNRAS* **359**, 1050 (2005)

ACKNOWLEDGEMENT

The authors gratefully acknowledge the support of the United States National Science Foundation (NSF) for the construction and operation of the LIGO Laboratory, the Science and Technology Facilities Council (STFC) of the United Kingdom, the Max-Planck-Society (MPS), and the State of Niedersachsen/Germany for support of the construction and operation of the GEO600 detector, the Italian Istituto Nazionale di Fisica Nucleare (INFN) and the French Centre National de la Recherche Scientifique (CNRS) for the construction and operation of the Virgo




FIG. 18. Final peakmap histograms for the outlier at 71.20 Hz. The dark thick curve corresponds to the full VSR2/VSR4 analysis; the light dashed (blue in color version) curve corresponds to VSR2 analysis while the light continuous (red in color version) curve corresponds to the analysis of VSR4 alone.




FIG. 19. Hough map (left) and peakmap (right) for the outlier at 71.20 Hz. As for the outlier at 30.88 Hz, this outlier is due to a concentration of peaks at the beginning of VSR4, barely visible in the right plot.

detector and the creation and support of the EGO consortium. The authors also gratefully acknowledge research support from these agencies as well as by the Australian Research Council, the International Science Linkages program of the Commonwealth of Australia, the Council of Scientific and Industrial Research of India, Department of Science and Technology, India, Science & Engineering Research Board (SERB), India, Ministry of Human Resource Development, India, the Spanish Ministerio de Economía y Competitividad, the Conselleria d'Economia i Competitivitat and Conselleria d'Educació, Cultura i Universitats of the Govern de les Illes Balears, the Foundation for Fundamental Research on Matter supported by the Netherlands Organisation for Scientific Research, the Polish Ministry of Science and Higher

Education, the FOCUS Programme of Foundation for Polish Science, the European Union, the Royal Society, the Scottish Funding Council, the Scottish Universities Physics Alliance, the National Aeronautics and Space Administration, the Hungarian Scientific Research Fund (OTKA), the Lyon Institute of Origins (LIO), the National Research Foundation of Korea, Industry Canada and the Province of Ontario through the Ministry of Economic Development and Innovation, the National Science and Engineering Research Council Canada, the Brazilian Ministry of Science, Technology, and Innovation, the Carnegie Trust, the Leverhulme Trust, the David and Lucile Packard Foundation, the Research Corporation, and the Alfred P. Sloan Foundation. The authors gratefully acknowledge the support of the NSF, STFC, MPS,

INFN, CNRS and the State of Niedersachsen/Germany for provision of computational resources.