

NOTES ON GEOGRAPHIC DISTRIBUTION

Reptilia, Squamata, Serpentes, Scolecophidia, Anomalepididae, *Liotyphlops* cf. *ternetzii* (Boulenger, 1896): first family record for the state of Ceará, Brazil

Daniel Loebmann

Universidade Estadual Paulista, Instituto de Biociências, Laboratório de Herpetologia.
Av. 24 A, 1515, Bairro Bela Vista. CEP 13506-900. Rio Claro, SP, Brazil.
E-mail: contato@danielloebmann.com

The Scolecophidia is a basal group in the phylogeny of snakes and includes three families of very odd creatures, collectively called Blind Snakes (Greene 1997). The family Anomalepididae (Dawn Blind Snakes) is considered the world's smallest snakes and they are represented currently by four genera comprising 17 species: *Anomalepis* Jan, 1860 (4 spp.), *Helminthophis* Peters, 1860 (3 spp.), *Liotyphlops* Peters, 1860 (8 spp.), and *Typhlophis* Fitzinger, 1843 (2 spp.) (Uetz et al. 2008), ranging from southern Middle America (Nicaragua, Costa Rica, and Panama) to South America (Argentina, Brazil, Colombia, Ecuador, Paraguay, Peru, and Uruguay) (Carreira 2004; Freire et al. 2007). *Liotyphlops* is the richest genus in the family with eight species currently recognized: *L. albirostris* (Peters, 1857), *L. anops* (Cope, 1899), *L. argaleus* Dixon and Kofron, 1984, *L. beui* (Amaral, 1924), *L. schubarti* Vanzolini, 1948, *L. ternetzii* (Boulenger, 1896), *L. wilderi* (Garman, 1883), and *L. trefauti* (Freire, Caramaschi and Argôlo, 2007) (Freire et al. op cit.).

During a field survey conducted on 10th July 2008 in the Ubajara National Park a juvenile specimen (total length = 121 mm) of *Liotyphlops* cf. *ternetzii* was collected (Figure 1). The specimens were found in the surroundings of Minas River (03°50'01.9" S, 40°54'15.6" W; 519 m above sea level), an area with a physiognomy of *Caatinga* (*sensu* Ab'Saber 1977). Although the external diagnostic characters analyzed combined with *Liotyphlops ternetzii* (see Dixon and Kofron 1984), I prefer to leave as confer status due to the fact that anomalepidids has a complex taxonomy and it is very difficult to determine precisely the species identification, such as in this case with a single juvenile found over a two years of expedition in the Plateau of Ibiapaba.

The specimen was deposited in the snakes collection of Institute Butantan, São Paulo, Brazil (IBSP 76856). Collecting permits was authorized by Ibama (License # 13571-1).



Figure 1. A juvenile of *Liotyphlops* cf. *ternetzii* (IBSP 76856) collected at Ubajara National Park, municipality of Ubajara, state of Ceará.

Liotyphlops ternetzii was known from the Brazilian states of Minas Gerais, Mato Grosso, Pará, São Paulo, and also for the countries of Argentina, Paraguay, and Uruguay (Cunha and Nascimento 1993; Carreira 2004; Recorder and Nogueira 2007). This finding represents the first record for the family Anomalepididae in state of Ceará and also the extension of the genus distribution as follows: ca. 720 km E from municipality of Capitão Poço, state of Pará, Brazil (*L. ternetzii*; Cunha and Nascimento 1993), 1,375 km N from the Grande Sertão Veredas National Park, state of Minas Gerais, Brazil (*L. ternetzii*; Recorder and Nogueira 2007), and ca. 780 km NW from municipality of São José da Lage, state of Alagoas, Brazil (*L. trefauti*; Freire et al. 2007).

NOTES ON GEOGRAPHIC DISTRIBUTION

Two hypotheses should be considered about the wide and fragmented distribution of *L. ternetzii*, based on available literature. Firstly, it is possible that *L. ternetzii* is, in fact, a complex of species. Second, the fossorial behavior of the species associated with its apparent low density becomes

very difficult find these snakes to in the wild and, perhaps, the distribution of the species is not so fragmented as actually considered. Therefore, the best way to try to solve this dilemma definitively is to conduct a taxonomic review for the currently recognized populations of *L. ternetzii*.

Acknowledgements

I am grateful to Valdir Germano (*Instituto Butantan*) for help in the specimen identification. Maria Cristina Oddone (Secretaria Especial de Aquicultura e Pesca) for reviewed the English grammar and style of the manuscript.

Literature cited

- Ab'Saber, A. N. 1977. Os Domínios Morfoclimáticos na América do Sul. Primeira aproximação. *Geomorfologia* 52: 1-159.
- Carreira, S. 2004. Geographic distribution: *Liotyphlops ternetzii*. *Herpetological Review* 35(4): 411-412.
- Cunha, O. R. and F. P. Nascimento. 1993. Ofídios da Amazônia. As cobras da região leste do Pará. *Boletim do Museu Paraense Emílio Goeldi (Série Zoologia)* 9(1): 1-191.
- Dixon, J. R. and Kofron, C. P. 1984. The Central and South American anomalepid snakes of the genus *Liotyphlops*. *Amphibia-Reptilia* 4(2-4): 241-264.
- Freire, E. M. X., U. Caramaschi, and A. J. S. Argolo. 2007. A new species of *Liotyphlops* (Serpentes: Anomalepididae) from the Atlantic Rain Forest of Northeastern Brazil. *Zootaxa* 1393: 19-26.
- Greene, H. W. 1997. Snakes: the evolution of mystery in nature. Berkeley: University of California Press. 351p.
- Recorder, R. and C. Nogueira 2007. Composição e diversidade de répteis na região sul do Parque Nacional Grande Sertão Veredas, Brasil Central. *Biota Neotropica* 7(3): 267-278.
- Uetz, P. 2008. TIGR. Reptile database. Electronic Database accessible at <http://www.reptile-database.org>. Captured on March 2008.

Received March 2009

Accepted April 2009

Published online May 2009