

Hisonotus acuen, a new and phenotypically variable cascudinho (Siluriformes, Loricariidae, Hypoptopomatinae) from the upper rio Xingu basin, Brazil

Gabriel S. C. Silva¹, Fábio F. Roxo¹, Claudio Oliveira¹

¹ Universidade Estadual Paulista, Departamento de Morfologia, Laboratório de Biologia e Genética de Peixes, Botucatu, SP, Brazil

Corresponding author: Fábio F. Roxo (roxoff@hotmail.com.br)

Academic editor: Carole Baldwin | Received 10 May 2014 | Accepted 13 September 2014 | Published 25 September 2014

<http://zoobank.org/763D2A89-717A-49A1-A7CC-6E41A15EE576>

Citation: Silva GSC, Roxo FF, Oliveira C (2014) *Hisonotus acuen*, a new and phenotypically variable cascudinho (Siluriformes, Loricariidae, Hypoptopomatinae) from the upper rio Xingu basin, Brazil. ZooKeys 442: 105–125. doi: 10.3897/zookeys.442.7870

Abstract

A new species of *Hisonotus* is described from the headwaters of the rio Xingu. The new species is distinguished from its congeners by having a functional V-shaped spinelet, odontodes not forming longitudinal aligned rows on the head and trunk, lower counts of the lateral and median series of abdominal plates, presence of a single rostral plate at the tip of the snout, absence of the unpaired platelets at typical adipose fin position, yellowish-tipped teeth, absence of conspicuous dark saddles and stripe on the body and higher number of teeth on the premaxillary and dentary. The new species, *Hisonotus acuen*, is restricted to headwaters of the rio Xingu basin, and is the first species of the genus *Hisonotus* described from the rio Xingu basin. *Hisonotus acuen* is highly variable in aspects of external body proportions, including body depth, snout length, and abdomen length. This variation is partly distributed within and among populations, and is not strongly correlated with body size. PCA of 83 adult specimens from six allopatric populations indicates the presence of continuous variation. Therefore, the available morphological data suggest that the individuals inhabiting the six localities of rio Xingu represent different populations of a single species. Low intraspecific variation in mitochondrial Cytochrome oxidase subunit I (COI) provides corroborative evidence.

Keywords

Biodiversity, Cascudinhos, Fresh-water, Neotropical fish, Taxonomy

Introduction

The subfamily Hypoptopomatinae is a monophyletic group of Loricariids (Schaefer, 2003) composed of 19 genera and 139 species (Eschmeyer and Fong 2014). Within this subfamily, *Hisonotus* Eigenmann & Eigenmann, 1889 comprises 33 valid species (Eschmeyer 2014). The genus *Hisonotus* was resurrected from the synonymy of *Otocinclus* by Schaefer (1998) based on the reduced or absent snout plates anterior to the nostril, rostrum with enlarged odontodes, and thickened plates forming the lateral rostral margin. However, the phylogenetic relationships in this genus are not well resolved (Britski and Garavello 2007) and, according to molecular (Chiachio et al. 2008; Cramer et al. 2011) and morphological (Martins et al. 2014) data, *Hisonotus* is a polyphyletic genus.

Although there is no definition of *Hisonotus* that supports its monophyly, many authors have considered this genus as valid. In the past decade, 18 species of *Hisonotus* have been described (Britski and Garavello 2007; Carvalho et al. 2008; Carvalho and Reis 2009; Carvalho and Reis 2011; Martins and Langeani 2012; Carvalho and Datovo 2012; Roxo et al. 2013; Roxo et al. 2014). Recently, during a collecting trip in tributaries of the rio Xingu basin, we found fish specimens that have the generally accepted characteristics of *Hisonotus* listed above but do not match any known species. Herein we describe the rio Xingu specimens as a new species.

Material and methods

All measurements and counts were taken on the left side of specimens. Measurements were taken point to point to the nearest 0.1 mm with a digital caliper. Body plate and osteology nomenclature followed Schaefer (1997) and measurements followed Carvalho and Reis (2009), as shown in Table 1. Abbreviations used in the text followed Carvalho and Reis (2009). Morphometrics are given as percentages of standard length (SL), except for subunits of the head region, which are expressed as percentages of head length (HL). Specimens were cleared and stained (c&s) according to the method of Taylor and Van Dyke (1985). Vertebral counts also include the five vertebrae that comprise the Weberian apparatus. Dorsal-fin ray counts include the spinelet as the first unbranched ray. All examined specimens were collected according to the Brazilian laws, and are deposited under permanent scientific collection licenses. After collection, specimens were euthanized using 1% benzocaine in water, fixed in 10% formaldehyde for morphological studies and preserved in 70% alcohol. For molecular studies specimens were fixed directly in 95% alcohol. Sequencing and molecular analysis followed Roxo et al. (2012). Institutional acronyms follow Fricke and Eschmeyer (2014). All samples are deposited at the LBP, Laboratório de Biologia e Genética de Peixes, Universidade Estadual Paulista, Botucatu; MZUSP, Museu de Zoologia, Universidade de São Paulo, São Paulo; NUP, Coleção Ictiológica do Nupelia, Universidade Estadual de Maringá, Maringá. Zoological nomenclature

follows the International Code of Zoological Nomenclature (4th Ed.). The GenBank accession numbers for Cytochrome oxidase subunit I (COI) sequences are: *H. acuen* – KM365043, KM365044, KM365045, KM365046, KM365047, KM365048, KM365049, KM365050, KM104473; *H. chromodontus* – KM104474, KM104475, JN998567, JN998566, JN998567, JN998565, JN998564, KM365054; *H. insperatus* – KM104485, GU701888, GU701749, GU701748, GU701747, GU701746, KM365055, KM365056, KM365057, KM365058, KM365059, KM365060, KM365061; *H. notatus* – JN998579, JN998581, JN998580; *H. oliveirai* – KM104486, KM365062, KM365063; *H. parsi* – KM104490, KM365042; *H. piracanjuba* – KM104487, KM104488, KM365051, KM365052, KM365053.

Principal component analysis (PCA)

Principal component analysis (PCA) was used to check overall variation among samples, including differences in morphometrics among species. PCA is a statistical procedure that uses orthogonal transformation to convert a set of observations of possibly correlated variables into a set of values of linearly uncorrelated variables called principal components (Jolliffe 2002). The analyses were made using all measurements listed above. Juvenile specimens below 18.0 mm SL were excluded from the analyses. PCA on covariances of base 10 logarithmically transformed measurements to reduce the influence of size were obtained using Past version 1.28 (Hammer et al. 2004). The PCA Loadings are presented in Table 2.

Results

Hisonotus acuen sp. n.

<http://zoobank.org/12454B4D-E9CA-4A89-A307-B77A703308C9>

Figs 1, 5, 6; Table 1

Holotype. MZUSP 115350, female, 25.9 mm SL, Brazil, Mato Grosso State, municipality of Querência, affluent of rio Toguro, rio Xingu basin, 13°00'26"S, 52°11'27"W, 01 Aug 2012, coll. C. Oliveira, M. Taylor, G.J.C. Silva & J.M. Henriques.

Paratypes. All from Brazil, Mato Grosso State, rio Xingu basin. LBP 15755, 16, 19.5–26.0 mm SL, municipality of Ribeirão Cascalheira, affluent of rio Suiá-Missu, 12°55'36"S, 51°53'27"W, 30 July 2012, coll. Oliveira C, Taylor M, Silva GJC, Henriques JM. LBP 16274, 27, 20.2–29.1 mm SL, 2 c&s 23.6–24.2 mm SL, municipality of Gaúcha do Norte, affluent of rio Culuene, 13°27'26"S, 53°09'36"W, 03 Aug 2012, coll. Oliveira C, Taylor M, Silva GJC, Henriques JM. LBP 16275, 29, 16.7–25.2 mm SL, 2 c&s 19.3–20.8 mm SL, municipality of Querência, affluent of rio Feio, 12°33'20"S, 52°16'16"W, 31 Sep 2012, coll. Oliveira C, Taylor M, Silva GJC, Henriques JM. LBP 16276, 9, 20.7–27.9 mm SL, 2 c&s 21.2–21.4 mm SL, munici-

Table 1. Morphometrics and meristic data for all analyzed specimens of *Hisonotus acuen* and for specimens by sub basins. SD = Standard deviation.

	<i>Hisonotus acuen</i> , n = 83					rio Toguero, n = 7					rio Culluene, n = 29					rio Suia Missu, n = 15					
	Holotype	Low	High	Mean	SD	Low	High	Mean	SD	Low	High	Mean	SD	Low	High	Mean	SD	Low	High	Mean	SD
SL	25.9	18.2	29.0	23.3	2.6	20.5	25.9	22.9	2.23	21.7	29.0	25.6	1.7	20.5	27.1	23.3	2.0				
Percents of SL																					
Head length	39.4	35.1	44.1	39.4	1.8	38.4	44.1	40.7	1.7	8.81	10.8	9.7	0.4	8.3	10.5	9.3	0.6				
Predorsal length	50.1	41.9	54.3	50.0	1.9	48.6	52.8	51.1	1.5	41.9	54.3	48.8	2.0	49.1	52.7	51.1	1.1				
Dorsal-fin spine length	20.6	19.4	25.8	21.9	1.4	19.5	22.8	20.7	1.06	4.9	6.3	5.6	0.3	4.0	6.2	5.0	0.6				
Anal-fin unbranched ray length	15.3	13.7	19.9	16.9	1.3	14.9	16.6	15.7	0.69	3.8	5.0	4.4	0.2	2.8	4.2	3.6	0.3				
Pectoral-fin spine length	24.0	15.9	28.8	25.0	1.9	21.6	26.6	23.9	1.68	22.7	28.8	25.6	1.4	4.9	6.6	5.8	0.5				
Pelvic-fin unbranched ray length	16.1	13.1	25.0	16.9	2.1	13.9	17.6	15.8	1.17	15.2	23.2	17.6	2.1	2.8	4.1	3.6	0.3				
Cleithral width	22.1	13.2	27.5	23.9	1.9	22.1	24.6	23.6	1.05	5.2	7.0	6.3	0.5	4.6	6.2	5.4	0.5				
Thoracic length	12.8	10.5	23.1	13.7	2.0	11.2	16.5	13.4	2.07	2.8	4.5	3.7	0.4	2.2	5.8	3.2	0.9				
Abdominal length	22.0	10.2	24.4	20.8	2.5	18.3	22.0	20.6	1.16	3.6	6.6	5.6	0.6	2.1	6.0	4.5	1.1				
Caudal-peduncle length	27.7	25.5	33.0	28.6	1.5	26.1	30.0	27.7	1.18	6.3	8.6	7.4	0.6	5.2	7.8	6.5	0.7				
Caudal-peduncle depth	9.5	8.6	11.1	9.6	0.4	8.7	10.3	9.5	0.54	8.9	11.1	9.9	0.4	9.0	10.3	9.5	0.4				
Percents of HL																					
Snout length	56.0	34.2	57.2	53.5	2.6	41.2	56.0	51.8	4.95	51.5	56.7	53.6	1.52	34.2	56.4	53.0	5.4				
Orbital diameter	13.4	11.2	16.2	13.1	0.9	11.6	13.8	12.7	0.83	11.7	15.0	13.4	0.8	1.0	1.3	1.1	0.1				
Interorbital width	36.2	15.6	41.8	36.3	3.4	33.1	53.0	37.1	1.05	3.1	4.1	3.7	0.2	2.7	3.9	3.3	0.3				
Head depth	51.7	35.1	53.1	44.2	4.6	40.0	56.5	45.5	6.1	3.9	5.3	4.6	0.3	3.2	4.8	4.0	0.5				
Suborbital depth	19.2	13.3	22.4	17.4	2.2	12.6	19.2	16.4	2.3	16.0	21.0	19.0	1.3	1.1	1.8	1.5	0.2				
Mandibular ramus	11.5	6.9	12.9	10.2	1.4	9.7	17.9	12.5	3.6	8.3	12.9	10.9	1.3	8.5	9.6	9.1	0.4				
Meristics																					
Left premaxillary teeth	23	14	27	14	4.1	14	20	-	3.87	22	24	23.3	0.8	14	21	17.2	2.5				
Left dentary teeth	18	12	23	21	3.5	13	18	-	2.38	12	23	21	3.5	13	17	15.0	1.5				
Left lateral scutes	23	22	24	23	0.7	23	24	23	0.82	20	23	21.3	1.0	24	24	24	-				

Table 1. Continued.

	córrego Xavante, n = 13				rio Coronel Vanick, n = 4				rio Feio, n = 15			
	Low	High	Mean	SD	Low	High	Mean	SD	Low	High	Mean	SD
SL	19.4	25.6	22.1	2.0	19.5	24.8	21.5	2.4	18.2	24.6	20.8	1.9
Percents of SL												
Head length	37.2	41.7	39.1	1.2	38.2	41.2	39.9	1.3	39.2	42.7	41.2	1.1
Predorsal length	47.5	51.5	49.2	1.1	49.1	52.9	51.1	1.6	49.0	52.9	51.3	1.3
Dorsal-fin spine length	20.2	24.7	22.2	1.1	19.4	21.6	20.5	0.7	20.0	24.4	22.3	1.1
Anal-fin unbranched ray length	16.4	19.9	17.6	0.8	13.9	16.7	15.7	1.2	15.5	19.7	17.7	1.0
Pectoral-fin spine length	15.9	27.7	24.5	3.6	23.2	25.6	24.8	1.0	23.3	26.8	24.9	1.0
Pelvic-fin unbranched ray length	15.5	20.7	17.8	1.7	15.6	16.3	15.9	0.3	13.1	25.0	16.8	2.8
Cleithral width	23.9	27.5	25.5	0.8	13.2	25.4	22.0	5.8	20.8	22.9	21.8	0.6
Thoracic length	10.5	16.6	12.8	1.5	11.6	13.5	12.7	0.8	11.2	19.0	12.9	2.0
Abdominal length	12.1	22.5	20.3	2.5	20.3	22.1	21.2	0.7	18.4	23.9	20.3	1.3
Caudal-peduncle length	27.1	32.1	29.2	1.3	26.2	27.6	26.7	0.6	26.8	30.9	28.8	1.4
Caudal-peduncle depth	9.1	10.3	9.7	0.3	9.1	9.7	9.4	0.2	8.6	9.9	9.2	0.3
Percents of HL												
Snout length	50.3	54.6	52.9	1.1	51.3	53.7	52.7	1.0	52.7	57.2	54.9	1.24
Orbital diameter	12.2	16.2	13.7	1.1	12.5	14.2	13.4	0.7	11.2	14.6	13.1	0.9
Interorbital width	35.8	41.8	38.5	1.6	33.8	36.8	35.1	1.2	15.6	37.6	32.9	5.1
Head depth	40.6	51.4	46.5	3.0	40.2	43.8	41.6	1.5	35.1	39.9	37.5	1.5
Suborbital depth	14.7	22.4	19.1	2.0	14.4	18.4	16.1	1.9	13.3	17.5	14.9	1.2
Mandibular ramus	7.3	10.1	8.8	1.0	7.2	8.9	8.1	0.4	6.9	11.5	9.3	1.2
Meristics												
Left premaxillary teeth	20	27	-	2.94	17	25	-	4.0	14	15	14	0.5
Left dentary teeth	16	22	-	2.75	14	17	-	1.5	12	13	12	0.5
Left lateral scutes	23	24	23	0.5	23	24	23	0.5	22	24	22	1.5

Table 2. Variable loadings in the first and second axes of size-free Principal Component Analysis (Axis 1 and Axis 2) of combined samples of *H. acuen*.

	Axis 1	Axis 2
Predorsal length	0.1871	-0.0346
Preanal length	0.2139	-0.0923
Head length	0.1631	-0.0090
Cleithral width	0.2370	0.1401
Dorsal-fin spine length	0.2015	0.0902
Base of dorsal-fin length	0.2838	-0.2170
Thorax length	0.2649	-0.1924
Pectoral-fin spine length	0.2234	0.0939
Abdomen length	0.2688	-0.5397
Pelvic-fin spine length	0.2155	0.4822
Anal-fin spine length	0.1747	0.2453
Lower cd spine	0.2088	0.0789
Caudal peduncle depth	0.1887	0.0911
Caudal peduncle length	0.2168	0.3770
Anal width	0.2543	-0.1638
Body depth	0.2608	-0.2413
Head depth	0.2506	0.0613
Snout length	0.1376	-0.1066
Interorbital width	0.2147	0.1441
Orbital diameter	0.1189	0.0778
Suborbital depth	0.2368	0.0426

pality of Ribeirão Cascalheira, affluent of rio Suiá-Missu, 12°53'04"S, 52°02'00"W, 30 Sep 2012, coll. Oliveira C, Taylor M, Silva GJC, Henriques JM. LBP 16277, 10, 18.9–23.3 mm SL, municipality of Querência, affluent of rio Feio, 12°31'55"S, 52°20'29"W, 31 Sep 2012, coll. Oliveira C, Taylor M, Silva GJC, Henriques JM. LBP 16278, 12, 18.8–25.1 mm SL, 2 c&s 26.8–27.1 mm SL, municipality of Primavera do Leste, córrego Xavante, 14°38'24"S, 53°55'38"W, 05 Aug 2012, coll. Oliveira C, Taylor M, Silva GJC, Henriques JM. LBP 16279, 10, 20.8–26.7 mm SL, municipality of Gaúcha do Norte, affluent of rio Culuene, 13°26'32"S, 53°08'45"W, 03 Aug 2012, coll. Oliveira C, Taylor M, Silva GJC, Henriques JM. LBP 16280, 11, 17.4–24.9 mm SL, municipality of Canarana, affluent of rio Culuene, 13°25'30"S, 52°16'47"W, 01 Aug 2012, coll. Oliveira C, Taylor M, Silva GJC, Henriques JM. LBP 16281, 4, 17.5–24.6 mm SL, same collection information as holotype. LBP 16282, 5, 17.5–23.9 mm SL, municipality of Canarana, rio Coronel Vanick, 13°31'34"S, 52°43'52"W, 02 Aug 2012, coll. Oliveira C, Taylor M, Silva GJC, Henriques JM. LBP 16283, 2, 21.3–24.2 mm SL, municipality of Canarana, affluent of rio Toguro, 13°16'52"S, 52°14'42"W, 01 Aug 2012, coll. Oliveira C, Taylor M, Silva GJC, Henriques JM. LBP 16284, 3, 20.2–24.3 mm SL, collected with holotype. LBP 18845, 1, 23.7 mm SL, municipality of Gaúcha do Norte, affluent of rio Culuene, 13°30'57"S, 53°06'39"W, 03 Aug



Figure 1. *Hisonotus acuen*, MZUSP 115350, female, 25.9 mm SL, holotype, from Mato Grosso State, municipality of Querência, affluent of rio Toguro, rio Xingu basin, 13°00'26"S, 52°11'27"W.

2012, coll. Oliveira C, Taylor M, Silva GJC, Henriques JM. NUP 16444, 5, 22.2–27.1 mm SL, municipality of Gaúcha do Norte, affluent of rio Culuene, 13°27'26"S, 53°09'36"W, 03 Aug 2012, coll. Oliveira C, Taylor M, Silva GJC, Henriques JM.

Diagnosis. *Hisonotus acuen* differs from all congeners except *H. bockmanni*, *H. chromodontus*, *H. insperatus*, *H. luteofrenatus*, *H. oliveirai* and *H. parsi* by having a functional V-shaped spinelet, Fig. 2a–f (*vs.* non-functional spinelet, a square ossification, or spinelet absent, Fig. 2g–h). It differs from *H. insperatus*, *H. parsi*, *H. luteofrenatus*, and




Figure 2. Spinelet variation among *Hisonotus* species. Figure a-f showing the extension of the bone forming a lock mechanism. In figure g-h, the bones lack the lock mechanism: **a** *H. acuen*, LBP 16276, 21.4 mm SL **b** *H. acuen*, LBP 16274, 21.1 mm SL **c** *H. oliveirai* LBP 13332, 23.7 mm SL **d** *H. chromodontus*, LBP 7964, 28.3 mm SL **e** *H. piracanjuba*, LBP 17256, 27.1 mm SL **f** *H. parestis*, NUP 10928, 23.6 mm SL **g** *H. cf. notatus*, LBP 3472, 25.8 mm SL **h** *H. depressicauda*, LBP 17474, 28.1 mm SL. Scale bar = 0.5 mm.

H. oliveirai by having odontodes not forming longitudinally aligned rows on head and trunk (*vs.* odontodes forming longitudinally aligned rows on head and trunk). *Hisonotus acuen* differs from *H. insperatus* and *H. luteofrenatus* by the lower counts of the lateral series of abdominal plates (4–5 *vs.* 6–8 and 7–8, respectively) and from *H. insperatus* by the lower counts of the lateral median plates (22–24 *vs.* 25–26). The new species can be distinguished from *H. luteofrenatus*, *H. oliveirai* and *H. parestis* by the presence of a single rostral plate at tip of snout (*vs.* presence of a pair of rostral plates at tip of snout); from *H. bockmanni* by the absence of unpaired platelets at typical adipose fin position (*vs.* presence of the unpaired platelets); from *H. chromodontus* by having yellowish-tipped teeth, Fig. 3a (*vs.* reddish-brown teeth, Fig. 3b), by having the caudal-fin color pattern mostly hyaline, except for dark blotch on origin of rays, and dark brown chromatophores largely concentrated on rays near lower caudal spine, Fig. 3c (*vs.* caudal-fin mostly dark brown with chromatophores largely concentrated on rays and membranes, and with two hyaline spots on middle of the fin, Fig. 3d); from *H. parestis* by the absence of conspicuous dark dorsal saddle and longitudinal stripe on the body (*vs.* inconspicuous dark saddles and stripe of the body) and from *H. insperatus* by the higher number of premaxillary (14–27 *vs.* 6–12) and dentary teeth (12–23 *vs.* 5–11).

Description. Morphometric and meristic data presented in Table 1. Maximum body length 29.0 mm SL. Dorsal profile of head in lateral view convex to straight from upper part of rostrum to posterior margin of nares, slightly curved from eyes to posterior margin of parieto supraoccipital, almost straight to dorsal-fin origin. Dorsal profile of trunk almost straight, descending from base of dorsal-fin origin to caudal peduncle. Ventral profile slightly concave from snout tip to anal-fin origin, slightly convex to caudal peduncle. Greatest body depth at dorsal-fin origin (13.5–22.8% SL). Greatest body width at cleithral region, gradually decreasing towards snout and caudal




Figure 3. **a** *H. acuen*, holotype, MZUSP 115350, 25.9 mm SL, showing the yellowish-tipped teeth **b** *H. chromodontus*, NUP 10924, 29.7 mm SL, showing the reddish-brown teeth **c** *H. acuen*, holotype, MZUSP 115350, 25.9 mm SL, showing the caudal-fin color pattern mostly hyaline, except for dark blotch on origin of rays, and dark brown chromatophores largely concentrated on rays near lower caudal spine **d** *H. chromodontus*, LBP7964, 27.3 mm SL, showing caudal-fin dark brown with chromatophores largely concentrated on rays and membranes, and with two hyaline spots on middle of the fin.

fin. Cross-section of caudal peduncle almost ellipsoid; rounded laterally and almost flat dorsally and ventrally.

Head rounded in dorsal view. Snout slightly pointed, its tip rounded, elongated (34.2–57.2% HL) and depressed in front of each nostril on dorsal surface. Dorsal and ventral series of odontodes completely covering anterior margin of snout; odontodes of snout similar in size to remaining ones found on head. Snout tip lacking band devoid of odontodes. Odontodes on head and trunk well defined and not forming longitudinal rows. Usually no tufts or crests of odontodes on head, in some juvenile specimens, a tiny tuft of odontodes at posterior tip of supraoccipital. Eyes small (11.2–16.2% HL), dorso-laterally positioned. Lips roundish and papillose; papillae uniformly distributed on base of dentary and premaxilla and slightly decreasing in size distally. Lower lip larger than upper lip; its border fringed. Maxillary barbel present. Teeth slender and bicuspid; mesial cusp larger than lateral cusp. Premaxillary teeth 14–27. Dentary teeth 12–23.

Dorsal-fin ii,7; dorsal-fin spinelet short and V-shaped; dorsal-fin lock functional; its origin slightly posterior to pelvic-fin origin. Tip of adpressed dorsal-fin rays slightly surpassing end of anal-fin base. Pectoral-fin i,6; tip of longest pectoral-fin ray almost reaching half of pelvic-fin length, when depressed. Pectoral axillary slit present be-




Figure 4. *Hisonotus acuen*, paratype, LBP 16278, 27.1 mm SL **a** Ventral view of abdominal plates **b** Cranial bones and dermal plates of the head in dorsal view. Scale bars: 1 mm.

tween pectoral-fin insertion and lateral process of cleithrum. Pectoral spine supporting odontodes anteroventrally. Pelvic-fin i,5; its tip not exceeding anal-fin origin when depressed in both sex. Pelvic-fin unbranched ray with dermal flap along its dorsal surface in males. Anal fin i,5; its tip reaching 7th and 8th plate from its origin. Caudal-fin i,14,i; distal margin forked. Adipose-fin absent. Total vertebrae 27.

Body covered with bony plates except on ventral part of head, around pectoral and pelvic-fin origin and on dorsal-fin base. Cleithrum and coracoid totally exposed. Arrector fossae partially enclosed by ventral lamina of coracoids. Abdomen entirely covered by plates in adults (about 23.0 mm SL); lateral plate series with elongate and large plates, formed by two lateral plate series, similar in size; median plates formed by four to five irregular plate series reaching anal shield (Fig. 4a). Lateral side of body entirely covered by plates; mid-dorsal plates poorly developed, reaching middle of dorsal-fin base; median plates not interrupted in median portion of body, but with 2 or 3 plates not perforated before end of series; mid-ventral plates exceed end of anal-fin base.

Parts of head osteology presented in Fig. 4b. Tip of snout formed by a single almost square rostral plate (r). Nasal (n) almost rectangular, forming anterior medial nostril margin in contact posteriorly with frontals (f), and anterior and lateral margins contacting pre-nasals (pn). Lateral surface of head formed by three posterior rostrum plates (pr1–pr3) similar in size. Complete infraorbital plate series, present just below posterior rostrum series, composed of five plates (io1–io5), fourth infraorbital expanded ventrally, all associated with latero-sensory canal system; first infraorbital (io1) largest and fifth smallest (io5). Preopercle (pop) present just under fifth infraorbital (io5); an elongated bone, covered by latero-sensory canal. Subocular cheek plates (cp1–cp2) present above preopercle plate (pop). Top of head composed of compound pterotic-supracleithrum (cpt), supraoccipital (soc), prefrontal (pf), frontal (f), and sphenotic (sp); cpt covered with fenestrae randomly distributed and with different sizes and shapes.

Color in alcohol. Large inconspicuous brown lateral stripe extending from tip of snout through inferior orbit to end of caudal peduncle Fig. 5a, e (very weak in some specimens, such as holotype Fig. 5c). Body ground color brown on dorsum, yellowish on ven-




Figure 5. Variation in external morphology and coloration of *H. acuen*: **a, b** LBP 16279, 26.86 mm SL, affluent of rio Culuene, municipality of Gaúcha do Norte **c, d** MZUSP 115350, 25.9 mm SL, holotype, affluent of rio Toguro, municipality of Querência **e (f)** LBP16275, 21.67 mm SL, affluent of rio Feio, municipality of Querência.



Figure 6. *Hisonotus acuen*, LBP 16284, live specimen, from affluent of rio Toguro, Querência, Mato Grosso State, Brazil. Photo: M. Taylor.

tral region under lateral stripe. Some specimens with dark saddle on mid-ventral to ventral portion of body (Fig. 5b). Dorsal, pectoral, pelvic and anal fins with brown dots on rays, varying in concentration of chromatophores from one individual to another; inter-radial membranes hyaline. Caudal fin hyaline, except for dark blotch on origin of rays, and dark brown chromatophores largely concentrated on rays near lower caudal spine (Fig. 5d). In some specimens, chromatophores forming two dark bands on middle of rays (Fig. 5b, e).

Color in life. Similar to pattern described for alcohol individuals, but with ground color light brown (Fig. 6).

Sexual dimorphism. Males bear a papilla posterior to urogenital opening and present the pelvic-fin unbranched ray with dermal flap along its dorsal surface. Both characters are absent in females.

Distribution. *Hisonotus acuen* is known from small to median-sized streams of the upper rio Xingu basin, Mato Grosso State in Brazil (Fig. 7a).

Habitat. *Hisonotus acuen* was collected on flat areas in creeks of headwaters of the rio Xingu basin in places of shallow clear waters with low current. The fishes are found associated with vegetation that covers the bottom and the border of the headwaters (Fig. 7b).

Etymology. The specific name “acuen” is in reference to the Xavante indigenous peoples, who in anthropological literature are known as “acuen”. These people are constituted by the natives inhabiting the east of the Mato Grosso State, living in the margins of the rivers Culuene, Xingu, Mortes and Araguaia.

Discussion

The new species has a functional V-shaped spinelet (Fig. 2a–f). Carvalho and Datovo (2012) first reported this structure in *H. bockmanni*, *H. chromodontus*, *H. insperatus* and *H. luteofrenatus*. Subsequently, Roxo et al. (2014) reported this character in *H. oliveirai* and *H. paresi*. The functional V-shaped spinelet (Fig. 2a–f) is a putative apomorphic character within *Hisonotus*, and may distinguish a monophyletic group within the genus (Carvalho and Datovo 2012). However, Martins et al. (2014) have a different interpretation, in which the spinelet in *H. chromodontus*, *H. luteofrenatus* and




Figure 7. a Map of the distribution of *H. acuen*. Red star = holotype locality, affluent of rio Toguro. Red circles = paratype localities, affluent of rio Culuene, affluent of rio Suiá-Missu, affluent of rio Feio, córrego Xavante, rio Coronel Vanick. All are tributaries of rio Xingu, Mato Grosso State, Brazil **b** Affluent of rio Feio, municipality of Querência, 12°33'20"S, 52°16'16"W, habitat where the specimens of *H. acuen* were found. Photo: M. Taylor.

H. piraicanjuba is reduced, and the locking mechanism is not functional (Martins et al. 2014, Fig. 11A), the same character state found in *H. armatus*, *H. depressicauda*, *H. francirochai* and *H. notatus*. Martins et al. (2014) also suggested that in *H. insperatus*




Figure 8. Scatter plot of Principal Component Analysis (PCA) of six allopatric populations of *H. acuen* ($n = 83$) indicating the presence of continuous external morphology variation. Purple circle = affluent of rio Toguro; Red cross = affluent of rio Culuene; Dark blue star = affluent of rio Suiá-Missu; Blue square = affluent of rio Feio; Yellow triangle = córrego Xavante; Green diamonds = rio Coronel Vanick. All are tributaries of rio Xingu, Mato Grosso State in Brazil.

the spinelet is absent (Martins et al. 2014, Fig. 11b). However, in our interpretation, *H. acuen*, *H. chromodontus*, *H. bockmanni*, *H. insperatus*, *H. luteofrenatus*, *H. oliveirai*, *H. parsi*, and *H. piracanjuba* exhibit a functional V-shaped spinelet, which is not present in *H. depressicauda* and *H. notatus* (Fig. 2). Therefore, despite the fact that the genus *Hisonotus* may not represent a monophyletic unit, we include *H. acuen* within *Hisonotus* pending a formal phylogenetic analysis of Hypoptopomatinae, and the species-level composition is established.

Hisonotus acuen exhibits a large amount of variation in external body proportions across its range (Fig. 5), especially in body depth at dorsal-fin origin (13.5–22.8% of SL), snout length (34.2–57.2% of HL), and abdomen length 10.2–24.4% of SL). This variation is partly distributed within populations, and partly between populations, and is not strongly correlated with body size. We performed a PCA to evaluate the morphometric variation within this new species. We compared the morphometric data of six populations found in different tributaries of the rio Xingu, and our results suggest that the range in morphology has a continuous distribution. The lack of phenotypic discontinuities among populations suggests they are not different species (Fig. 8). Additionally, we found that the genetic variation of the Cytochrome Oxidase I (COI) gene within the populations of *H. acuen* is 1%, and that variation among closely related congeners (i.e. *H. chromodontus*, *H. insperatus*, *H. oliveirai*, *H. parsi* and *H. piracanjuba*) is more than 17% (see Table 3 and Fig. 9; sequences can




Figure 9. Phenogram constructed using Neighbor-Joining (NJ) method, based on the COI gene (581 pb). Numbers above branches are bootstrap values obtained from 1,000 pseudoreplicates. Values below 90% are not shown.

Table 3. Genetic distance (and standard deviation) between *Hisonotus* species and specimens of the same species (main diagonal). This analysis was performed using Kimura 2-parameter substitution model, Gamma distribution and 1,000 bootstrap pseudoreplicates.

	1	2	3	4	5	6	7
1. <i>H. acuen</i>	1.0±0.2						
2. <i>H. parsi</i>	17.0%±2.3	0±0					
3. <i>H. piracanjuba</i>	21.2%±2.6	6.8%±1.3	0±0				
4. <i>H. chromodontus</i>	19.3%±2.5	19.5%±2.6	20.0%±2.6	0±0			
5. <i>H. notatus</i>	20.5%±2.8	19.0%±2.7	18.5%±2.4	22.3%±2.9	0±0		
6. <i>H. insperatus</i>	21.4%±2.7	8.0%±1.4	5.1%±0.9	21.6%±2.7	16.1%±2.1	0.1±0.1	
7. <i>H. oliveirai</i>	20.2%±2.5	6.8%±1.2	4.1%±0.9	21.2%±2.7	18.1%±2.3	3.0%±0.7	0.6±0.3

be downloaded from GenBank using the accession numbers provided in Methods). Therefore, the available morphological and molecular data support the recognition of the individuals inhabiting the six localities of rio Xingu and representing different populations as a single species.

The new species *H. acuen* is the first described species of *Hisonotus* from the rio Xingu basin, and is externally very similar to *H. chromodontus*, a species from the rio Tapajos basin. The coloration of the caudal fin and the tip of the teeth distinguish these species that also are very different genetically (i.e. 19.3% of genetic divergence; Table 3 and Fig. 9). Britski and Garavello (2003) discussed the coloration of the teeth of *H. chromodontus*, reporting that in more than one hundred specimens examined, varying from 12.0 to 32.2 mm SL, all tooth-tips have a reddish-brown color. We analyzed more than one hundred specimens of *H. chromodontus* from the museum collections of LBP and NUP, and found the same reddish-brown tooth tips. This tooth features appears to be unique within the genus *Hisonotus*. A very similar external morphology, as well as the presence of the functional V-shaped spinelet among *H. acuen* and *H. chromodontus*, could suggest a close relationship between these species.

Comparative material

All from Brazil, except when stated otherwise: *Hisonotus aky* (Azpelicueta, Casciotta, Almirón & Koerber, 2004): MHNG 2643.039, 2, 33.1–34.2 mm SL, paratypes, arroyo Fortaleza, Argentina; *Hisonotus bocaiúva* Roxo, Silva, Oliveira & Zawadzki, 2013: MZUSP 112204, male, 24.2 mm SL, holotype, córrego Cachoeira, Bocaiúva, Minas Gerais; LBP 9817, 9, 3 c&s, 18.3–23.2 mm SL, paratypes, córrego Cachoeira, Bocaiúva, Minas Gerais; *Hisonotus carreiro* Carvalho & Reis, 2011: MCP 40943, 3, 33.6–35.8 mm SL, arroyo Guabiju, Guabiju, Rio Grande do Sul; *Hisonotus charrua* Almirón, Azpelicueta, Casciotta & Litz, 2006: LBP 4861, 1, 35.9 mm SL, arroyo Guaviyú, Artigas, Uruguay; MHNG 2650.051, 1, 34.2 mm SL, paratype, arroyo Aspinillar, Uruguay; *Hisonotus chromodontus* Britski & Garavello, 2007: LBP 7964, 25, 24.0–28.3 mm SL, 3 females c&s, 26.5–28.9 mm SL, 1 male c&s 24.9 mm SL, rio dos

Patos, Nova Mutum, Mato Grosso; LBP 7974, 26, 17.7–24.8 mm SL, rio dos Patos, Nova Mutum, Mato Grosso; LBP 12278, 2, 26.7–28.7 mm SL, 1 unsexed c&s, 26.7 mm SL, rio Sumidouro, Tangará da Serra, Mato Grosso; MZUSP 45355, 25.9 mm SL, holotype, affluent of rio Preto, Diamantino, Mato Grosso; MZUSP 70758, 7, 19.4–23.9 mm SL, paratype, riacho Loanda, Sinop, Mato Grosso; NUP 10924, 24, 19.5–31.5 mm SL, rio Preto, Diamantino, Mato Grosso; *Hisonotus depressicauda* (Miranda Ribeiro, 1918): MZUSP 5383, 24.4 mm SL, paralectotype, Sorocaba; LBP 17474, 5 c&s, 18.1–24.0 mm SL, rio Araquá, Botucatu, São Paulo; *Hisonotus francirochaei* (Ihering, 1928): LBP 13923, 22, 25.7–35.7 mm SL, córrego sem nome, Capitinga, Minas Gerais; MZUSP 3258, 29.4 mm SL, lectotype, rio Grande, São Paulo; *Hisonotus heterogaster* Carvalho & Reis, 2011: LBP 3335, 39, 20.8–30.1 mm SL, arroio sem nome, rio Grande, Rio Grande do Sul; *Hisonotus insperatus* Britski & Garavello, 2003: LBP 1299, 3, 23.5–29.6 mm SL, 1 female c&s, 24.8 mm SL, rio Araquá, Botucatu, São Paulo; LBP 1316, 2, 24.1–27.4 mm SL, 1 female c&s, 24.7 mm SL, 1 male c&s, 23.9 mm SL, rio Araquá, Botucatu, São Paulo; LBP 1344, 2, 22.9–24.9 mm SL, rio Araquá, Botucatu, São Paulo; LBP 1373, 1, 25.8 mm SL, rio Araquá, Botucatu, São Paulo; LBP 1405, 2, 22.2–27.3 mm SL, rio Araquá, Botucatu, São Paulo; LBP 4699, 17, 19.6–26.9 mm SL, 4 females c&s, 20.3–26.8 mm SL, 3 males c&s, 24.3–26.1 mm SL, ribeirão Cubatão, Marapoama, São Paulo; LBP 4945, 5, 27.3–28.5 mm SL, 2 females c&s, 28.2–29.9 mm SL, Botucatu, São Paulo; LBP 6770, 5, 25.1–28.2 mm SL, 3 females c&s, 20.0–27.0 mm SL, ribeirão Cubatão, Marapoama, São Paulo; LBP 13336, 1 female c&s, 26.0 mm SL, rio Capivara, Botucatu, São Paulo; LBP 13337, 2 females c&s, 27.4–28.6 mm SL, rio Araquá, Botucatu, São Paulo; MZUSP 22826, 1, 25.4 mm SL, paratype, córrego Água Tirada, Três Lagoas, Mato Grosso; MZUSP 24832, 1, 23.8 mm SL, paratype, rio Corumbataí, Corumbataí, São Paulo; MZUSP 78957, 29.6 mm SL, holotype, rio Capivara, Botucatu, São Paulo; MZUSP 78960, 31, 12.6–26.0 mm SL, paratypes, 5 c&s, 22.7–24.7 mm SL, rio Pardo, Botucatu, São Paulo; MZUSP 78965, 10, 15.6–28.6 mm SL, paratypes, 3 c&s, not measured, rio Araquá, Botucatu, São Paulo; MZUSP 78968, 5, 24.1–27.3 mm SL, paratypes, córrego da Figueira, Lins, São Paulo; *Hisonotus iota* Carvalho & Reis, 2009: LBP 13072, 5, 32.3–33.0 mm SL, rio Chapecó, Coronel Freitas, Santa Catarina; *Hisonotus laevior* Cope, 1894: LBP 3377, 1, 25.2 mm SL, arroio dos Corrientes, Pelotas, Rio Grande do Sul; LBP 6037, 8, 33.4–47.0 mm SL, rio Maquiné, Osório, Rio Grande do Sul; LBP 13187, 7, 19.4–45.8 mm SL, córrego sem nome, Camaquá, Rio Grande do Sul; *Hisonotus leucofrenatus* (Miranda Ribeiro, 1908): LBP 2085, 7, 38.3–50.6 mm SL, rio Sagrado, Morretes, Paraná; LBP 6837, 36, 35.1–43.5 mm SL, rio Fau, Miracatu, São Paulo; *Hisonotus leucophrys* Carvalho & Reis, 2009: LBP 13065, 6, 17.2–33.6 mm SL, rio Ariranhas, Xavantina, Santa Catarina; LBP 13073, 1, 36.8 mm SL, rio Guarita, Palmitinho, Rio Grande do Sul; *Hisonotus luteofrenatus* Britski & Garavello, 2007: MZUSP 62593, 28.6 mm SL, holotype, córrego Loanda, Cláudia, Mato Grosso; MZUSP 62594, 8, 22.4–30.5 mm SL, paratype, riacho Selma, Sinop, Mato Grosso; MZUSP 95940, 3, 26.1–28.5 mm SL, affluent of rio Teles Pires, Itaúba, Mato Grosso; *Hisonotus megaloplax* Carvalho & Reis, 2009: LBP 13108, 6, 36.4–37.8 mm SL, cór-

rego sem nome, Saldanha Marinho, Rio Grande do Sul; *Hisonotus montanus* Carvalho & Reis, 2009: LBP 13051, 3, 26.4–27.2 mm SL, rio Goiabeiras, Vargem, Santa Catarina; LBP 13055, 5, 24.8–31.9 mm SL, rio Canoas, Vargem, Santa Catarina; *Hisonotus nigricauda* (Boulenger, 1891): LBP579, 16, 34.1–40.1 mm SL, rio Guaíba, Eldorado do Sul, Rio Grande do Sul; *Hisonotus notatus* Eigenmann & Eigenmann, 1889: LBP 3472, 20, 21.0–34.3 mm SL, 2 males c&s 25.8–26.5 mm SL, 1 female c&s, 25.0 mm SL, rio Aduelas, Macaé, Rio de Janeiro; LBP 10742, 25, 24.4–43.3 mm SL, rio Macabu, Conceição de Macabu, Rio de Janeiro; *Hisonotus oliveirai* Roxo, Zawadzki & Troy, 2014: MZUSP 115061, female, 26.4 mm SL, holotype, ribeirão Cambira, affluent of rio Ivaí, Cambira, Paraná; LBP 13332, 1 male, 23.2 mm SL, 1 unsexed c&s, 23.7 mm SL, paratype, rio Mourão, rio Ivaí basin, Campo Mourão, LBP 17578, 3 females, 27.7–30.4 mm SL, 2 males, 25.4–26.1 mm SL, paratypes, rio Mourão, rio Ivaí basin, boundary between Engenheiro Beltrão and Quinta do Sol; NUP 3578, 7 females, 27.8–28.1 mm SL, 8 males, 24.7–26.8 mm SL, 1 female c&s, 27.6 mm SL, 1 male c&s, 25.5 mm SL, paratypes, ribeirão Salto Grande, rio Ivaí basin, Maria Helena; *Hisonotus parsi* Roxo, Zawadzki & Troy, 2014: MZUSP 115062, female, 26.2 mm SL, holotype, riacho Águas Claras, affluent of rio Sepotuba, Santo Afonso; LBP 13351, 9, 14.7–24.3 mm SL, paratype, riacho Águas Claras, Santo Afonso; LBP 13352, 1, 23.7 mm SL, paratype, riacho Águas Claras, Santo Afonso; NUP 10928, 2 males, 23.2–24.2 mm SL, paratype, 2 c&s, 23.6–24.2 mm SL, riacho Águas Claras, affluente of rio Sepotuba, Santo Afonso; NUP 10976, 3 unsexed, 16.7–20.5 mm SL, paratype, riacho São Jorge, Santo Afonso; *Hisonotus piraicanjuba* Martins & Langeani, 2012: LBP 17256, 9, 17.2–26.3 mm SL, 1, c&s 27.1 mm SL, córrego sem nome, Morrinhos, Goiás; NUP 5059, 1, 24.7 mm SL, córrego Posse, Anápolis, Goiás; NUP 10979, 3, 21.4–21.8 mm SL, ribeirão Bocaina, Piraicanjuba, Goiás; *Hisonotus prata* Carvalho & Reis, 2011: MCP 40492, 18, 19.5–33.2 mm SL, rio da Prata, Nova Prata, Rio Grande do Sul; LBP 9918, 14, 21.7–32.6 mm SL, Laguna dos Patos system, Nova Prata, Rio Grande do Sul; *Hisonotus ringueleti* Aquino, Schaefer & Miquelarena, 2001: FMNH 108806, 2, 25.7–32.2 mm SL, rio Quaraí basin, Uruguay; LBP 13148, 1, 24.5 mm SL, arroio Putiá, Uruguiana, Rio Grande do Sul; *Hisonotus* sp.: LBP 8276, 1 c&s, 25.6 mm SL, rio Verde Grande, Jaíba, Minas Gerais; *Microlepidogaster arachis* Martins, Calegari & Langeani, 2013: LBP 10882, 3, 22.8–35.3 mm SL, rio Paraná basin, Araxás, Minas Gerais; *Microlepidogaster dimorpha* Martins & Langeani, 2011: LBP 10683, 2, 28.8–35.6 mm SL; rio Paraná basin, Uberaba, Minas Gerais; *Otothyris travassosi* Garavello, Britski & Schaefer, 1998: LBP 1971, 13, 14.0–27.2 mm SL; coastal drainage, Canavieiras, Bahia; *Otothyropsis marapoama* Ribeiro, Carvalho & Melo, 2005: LBP 4698, 6, 23.9–36.3 mm SL; rio Tietê basin, Marapoama, São Paulo; *Parotocinclus aripuanensis* Garavello, 1988: LBP 10981, 33, 15.63–18.47 mm SL, rio Lageado, Guajará Mirim, Rondônia; *Parotocinclus* aff. *spilurus* (Fowler, 1941): LBP 5624, 21.5–22.0 mm SL, rio Maravilha, Balsas, Maranhão; *Parotocinclus* cf. *bahiensis* (Miranda Ribeiro, 1918): LBP 7182, 3, 27.9–35.6 mm SL; rio Paraguaçu basin, Lençóis, Bahia; *Parotocinclus maculicauda* (Steindachner, 1877): LBP 2869, 15, 20.2–44.7 mm SL, rio Ribeira do Iguape basin, Miracatu, São Paulo; *Parotocinclus prata* Ribeiro,

Melo & Pereira, 2002: LIRP 1136, 38, 19.8–41.9 mm SL; rio São Francisco basin, Presidente Olegário, Minas Gerais; *Parotocinclus* sp.: LBP 1572, 3, 19.0–24.0 mm SL, ribeirão Ínsula, Barra do Garça, Mato Grosso; LBP 2414, 19, 16.7–20.8 mm SL, 1 c&s 23.6 mm SL, córrego Fundo, Barra do Garça, Mato Grosso.

Acknowledgments

We wish to thank Guilherme José da Costa Silva, Jefferson M. Henriques, and Martin Taylor for their help during the collection expeditions. Martin Taylor, for the live fish and type locality photos. James S. Albert and Ricardo Britzke, for reading the manuscript and giving valuable suggestions. Maria Thereza for English review. This research was supported by the Brazilian agencies FAPESP (Fundação de Amparo à Pesquisa do Estado de São Paulo, proc. 2010/01610-9 to FFR and 2012/01622-2 to GSCS) and CNPq (Conselho Nacional de Desenvolvimento Científico e Tecnológico).

References

- Almirón AE, Azpelicueta MM, Casciotta JR, Litz T (2006) A new species of *Hisonotus* (Siluriformes, Loricariidae, Otothyriini) from the República Oriental del Uruguay. *Revue suisse de Zoologie* 113(1): 87–94.
- Aquino AE, Schaefer SA, Miquelarena AM (2001) A new species of *Hisonotus* (Siluriformes, Loricariidae) of the Upper Río Uruguay Basin. *American Museum Novitates* 3333: 1–12. doi: 10.1206/0003-0082(2001)333<0001:ANSOHS>2.0.CO;2
- Azpelicueta MM, Casciotta JR, Almirón AE, Koerber S (2004) A new species of Otothyriini (Siluriformes: Loricariidae: Hypoptopomatinae) from the Río Uruguay basin, Argentina. *Verhandlungen der Gesellschaft für Ichthyologie* 4: 81–90.
- Boulenger GA (1891) An account of the siluroid fishes obtained by Dr. H. von Ihering and Herr Sebastian Wolff in the Province Rio Grande do Sul, Brazil. *Proceedings of the Zoological Society of London* 2: 231–235.
- Britski HA, Garavello JC (2003) *Hisonotus insperatus*: new species, from the upper rio Paraná basin (Pisces: Ostariophysi: Loricariidae). *Copeia* 2003(3): 588–593. doi: 10.1643/CI-02-23R
- Britski HA, Garavello JC (2007) Description of two new sympatric species of the genus *Hisonotus* Eigenmann and Eigenmann, 1889, from upper rio Tapajós, Mato Grosso state, Brazil (Pisces: Ostariophysi: Loricariidae). *Brazilian Journal of Biology* 67(3): 413–420. doi: 10.1590/S1519-69842007000300005
- Carvalho M, Datovo A (2012) A new species of cascudinho of the genus *Hisonotus* (Siluriformes: Loricariidae: Hypoptopomatinae) from the upper rio Tapajós basin, Brazil. *Copeia* 2012(2): 266–275. doi: 10.1643/CI-11-016
- Carvalho TP, Lehmann PA, Pereira EHL, Reis RE (2008) A New Species of *Hisonotus* (Siluriformes: Loricariidae: Hypoptopomatinae) from the Laguna dos Patos basin, Southern Brazil. *Copeia* 3(3): 510–516. doi: 10.1643/CI-07-130

- Carvalho TP, Reis RE (2009) Four new species of *Hisonotus* (Siluriformes: Loricariidae) from the upper rio Uruguay, southeastern South America, with a review of the genus in the rio Uruguay basin. *Zootaxa* 2113: 1–40.
- Carvalho TP, Reis RE (2011) Taxonomic review of *Hisonotus* Eigenmann and Eigenmann (Siluriformes: Loricariidae: Hypoptopomatinae) from the laguna dos Patos system, southern Brazil. *Neotropical Ichthyology* 9(1): 1–48. doi: 10.1590/S0074-02762011000900001
- Chiachio MC, Oliveira C, Montoya-Burgos JI (2008) Molecular systematic and historical biogeography of the armored Neotropical catfishes Hypoptopomatinae and Neoplecostominae (Siluriformes: Loricariidae). *Molecular Phylogenetic and Evolution* 49(2): 606–617. doi: 10.1016/j.ympev.2008.08.013
- Cope ED (1894) On the fishes obtained by the Naturalist Expedition in Rio Grande do Sul. *Proceedings of the American Philosophical Society* 33: 84–108, Pls. 4–9.
- Cramer CA, Bonatto SL, Reis R (2011) Molecular Phylogeny of the Neoplecostominae and Hypoptopomatinae (Siluriformes: Loricariidae) using Multiple Genes. *Molecular Phylogenetic and Evolution* 59(1): 43–52. doi: 10.1016/j.ympev.2011.01.002
- Eigenmann CH, Eigenmann RS (1889) Preliminary notes on South American Nematognathi. *Proceedings of the California Academy of Sciences* 1: 119–172.
- Fowler HW (1941) A collection of fresh-water fishes obtained in eastern Brazil by Dr. Rodolpho von Ihering. *Proceedings of the Academy of Natural Sciences of Philadelphia* 93: 123–199.
- Garavello JC (1988) Three new species of *Parotocinclus* Eigenmann & Eigenmann, 1889 with comments on their geographical distribution (Pisces, Loricariidae). *Naturalia* (São Paulo) 13: 117–128.
- Garavello JC, Britski HA, Schaefer SA (1998) Systematics of the genus *Otothyris* Myers 1927, with comments on geographic distribution (Siluriformes: Loricariidae: Hypoptopomatinae). *American Museum Novitates* 3222: 1–19.
- Hammer O, Harper DAT, Ryan PD (2004) *Past - Palaeontological Statistics*, ver. 1.32. Oslo: University of Oslo.
- Ihering R von (1928) Uma nova espécie de *Otocinclus* (Pisces. Nematognatha) “cascudinho” de S. Paulo. *Boletim Biologia, Trabalho Laboratório de Parasitologia Faculdade de Medicina, São Paulo* 11(42): 1–3.
- Jolliffe IT (2002) *Principal Component Analysis*. Springer Series in Statistics, 2nd ed, Springer, New York, 487 pp.
- Martins FO, Calegari BB, Langeani F (2013) *Microlepidogaster arachas*, a new species of hypoptopomatine catfish (Siluriformes: Loricariidae) from the upper rio Paraná basin, Brazil. *Zootaxa* 3608(5): 379–388. doi: 10.11646/zootaxa.3608.5.6
- Martins FO, Britski HA, Langeani F (2014) Systematics of *Pseudotothyris* (Loricariidae: Hypoptopomatinae). *Zoological Journal of the Linnean Society* 170: 822–874. doi: 10.1111/zoj.12107
- Martins FO, Langeani F (2011) *Microlepidogaster dimorpha*, a new species of Hypoptopomatinae (Siluriformes: Loricariidae) from the upper Rio Paraná system. *Neotropical Ichthyology* 9(1): 79–86. doi: 10.1590/S1679-62252011000100005
- Martins FO, Langeani F (2012) *Hisonotus piracanjuba*, a new species of Hypoptopomatinae (Siluriformes: Loricariidae) from the rio Paranaíba, upper rio Paraná system, central Brazil. *Ichthyological Exploration of Freshwaters* 23: 29–36.

- Miranda Ribeiro A de (1908) Peixes da Ribeira. Resultados de excursão do Sr. Ricardo Krone, membro correspondente do Museu Nacional do Rio de Janeiro. Kosmos, Rio de Janeiro 5(2): 5 unnum. pp.
- Miranda Ribeiro A de (1918) Três gêneros e dezessete espécies novas de peixes Brasileiros. Revista do Museu Paulista 10: 631–646.
- Ribeiro AC, Carvalho M, Melo ALA (2005) Description and relationships of *Otothyropsis marapoama*, a new genus and species of Hypoptopomatinae catfish (Siluriformes: Loricariidae) from rio Tietê basin, southeastern Brazil. Neotropical Ichthyology 3(4): 489–498. doi: 10.1590/S1679-62252005000400006
- Ribeiro AC, Melo ALA, Pereira EHL (2002) A new species of *Parotocinclus* (Siluriformes: Loricariidae) from the rio São Francisco basin, southeastern Brazil. Ichthyological Exploration of Freshwaters 13(3): 217–224.
- Roxo FF, Oliveira C, Zawadzki CH (2012) Three new species of *Neoplecostomus* (Teleostei: Siluriformes: Loricariidae) from the Upper Rio Paraná basin of southeastern Brazil. Zootaxa 3233: 1–12.
- Roxo FF, Silva GSC, Oliveira C, Zawadzki CH (2013) *Hisonotus bocaiuva*, a new species from the rio São Francisco basin, Brazil (Teleostei: Loricariidae). Ichthyological Exploration of Freshwaters 23(4): 319–326.
- Roxo FF, Zawadzki CH, Troy WP (2014) Description of two new species of *Hisonotus* Eigenmann & Eigenmann, 1889 (Ostariophysi, Loricariidae) from the rio Paraná-Paraguay basin, Brazil 395: 57–78. doi: 10.3897/zookeys.395.6910
- Schaefer SA (1997) The Neotropical cascudinhos: Systematics and biogeography of the *Otocinclus* catfishes (Siluriformes: Loricariidae). Proceedings of the Academy of Natural Sciences of Philadelphia 148: 1–120.
- Schaefer SA (1998) Conflict and resolution: impact of new taxa on phylogenetic studies of the Neotropical cascudinhos (Siluroidei: Loricariidae). In: Malabarba LR, Reis RE, Vari RP, Lucena ZMS, Lucena CAS (Eds) Phylogeny and classification of neotropical fishes. Edipucrs, Porto Alegre, RS, 375–400.
- Steindachner F (1877) Die Süßwasserfische des südöstlichen Brasilien. (IV). Sitzungsberichte der Kaiserlichen Akademie der Wissenschaften. Mathematisch-Naturwissenschaftliche Classe 76(1): 217–230.
- Taylor WR, Van Dyke GC (1985) Revised procedures for staining and clearing small fishes and other vertebrates for bone and cartilage study. Cybium 9: 107–109.

Electronic references

- Eschmeyer W (2014) Catalog of Fishes. Electronic publication in “World Wide Web”. <http://research.calacademy.org/ichthyology/catalog> [accessed January 2014]
- Eschmeyer WN, Fong JD (2014) Species by Family/Subfamily. <http://research.calacademy.org/research/ichthyology/catalog/SpeciesByFamily.asp/> [accessed January 2014]
- Fricke R, Eschmeyer WN (2014) Catalog of fishes. <http://research.calacademy.org/research/ichthyology/catalog/collections.asp> [accessed January 2014]