

Research

Dendrochemical analysis of *Brachystegia longifolia* Benth as an indicator of Sulphur deposition in a copper smelting area of Mufulira, Zambia

Charles Mulenga¹ · Daigard Ricardo Ortega-Rodriguez² · Hermes Dias Brito³ · Darius Phiri⁴ · Abubakar Sadiq Mohammed⁵ · Chisala A. Lwali⁴ · James H. Speer⁶

Received: 20 December 2024 / Accepted: 9 May 2025

Published online: 27 May 2025

© The Author(s) 2025 **OPEN**

Abstract

High levels of Sulphur dioxide (SO₂) emanating from a copper smelter in Mufulira, Zambia has been reported around the year 2000, exceeding environmental quality guidelines. This resulted in the overhauling and upgrading of an obsolete smelter, the principal source of pollution. However, there is limited environmental evidence on the comparative variations in SO₂ emissions in Mufulira before and after the smelter upgrade. This study investigated the dendrochemical potential of *Brachystegia longifolia* in detecting environmental change induced by variations in industrial pollution load. *B. longifolia* wood discs were extracted at breast height, air-dried and sanded until the transverse surface and annual rings were clearly visible. μ -XRF spectrometer was used to determine the distribution of Sulphur by scanning and exposing samples along the bark-pith transverse surface to the X-ray beam, obtaining X-ray transmission intensity values of counts per second (cps) for Sulphur. Finally, the same radial slices were then scanned with a cellulose acetate calibration scale using an X-ray densitometry chamber. Digital images were then analysed with Windendro® program to determine wood density. The mean Sulphur distribution in tree rings at the beginning (2007) and completion (2014) of the smelter upgrades were 15.41 cps and 17.51 cps, respectively. By 2019, the highest Sulphur content in *B. longifolia* wood had significantly ($p < 0.05$) reduced to 13.28 cps. Furthermore, a significant reduction in wood density with increasing tree age was observed after the installation of the new smelter. 0.78 g/cm³ and 0.75 g/cm³ mean wood density were recorded in annual rings at the beginning and completion of the upgrades, respectively. However, the mean wood density had significantly ($p < 0.05$) reduced to 0.67 g/cm³ by 2019. The significant variations in the distribution of Sulphur and wood density in tree rings formed before and after upgrading an obsolete copper smelter demonstrates the dendrochemical potential of *B. longifolia* and possibly other Miombo woodland species, offering an opportunity to biomonitor the impact of industrial activities in the Miombo landscapes which are extensively distributed across sub-Saharan Africa.

Article Highlights

- Mining processes emit potentially toxic elements, including SO₂ that are dispersed across nearby landscapes.

✉ Charles Mulenga, charles.mulenga@cbru.ac.zm | ¹Department Biomaterial Science and Technology, School of Natural Resources, The Copperbelt University, Kitwe, Zambia. ²Departamento de Ciências Florestais, Escola Superior de Agricultura Luiz de Queiroz, Universidade de São Paulo (USP), Piracicaba, SP, Brazil. ³Centro de Ciências Naturais Aplicadas (UNESPetro – IGCE), Universidade Estadual Paulista, Rio Claro, SP, Brazil. ⁴Department of Plant and Environmental, Sciences School of Natural Resources, The Copperbelt University, Kitwe, Zambia. ⁵Department of Wood Technology, Tamale Technical University, Tamale, Ghana. ⁶Department of Earth and Environmental Systems, Indiana State University, Terre Haute, USA.



- Our study demonstrates that *B. longifolia* absorbs and sequesters Sulphur in a trend that reflects the generation of SO₂ from a point source.
- These results suggest that *B. longifolia* and possibly other miombo species have the potential to biomark and bio-monitor environmental change.

Keywords Copper smelter · Sulphur dioxide · *Brachystegia longifolia* · Annual rings · Sulphur distribution · Wood density

1 Background

Mufulira town (Zambia) is one of the most polluted mining towns in the World [1]. A number of reports [2, 3] highlight the high levels of Sulphur dioxide (SO₂) around the year 2000 emanating from a copper smelter located in Kankoyo township dispersed across the nearby landscape, exceeding environmental quality guidelines-EQG. This resulted in overhauling and upgrading the smelter in 2007 and by 2014, the facility had been upgraded to a stage where it could capture 97% of SO₂ produced at the facility [3], converting it to Sulphuric acid-H₂SO₄. However, there is limited environmental evidence on the comparative variations in SO₂ emissions in Mufulira before and after the smelter upgrade.

Mineral beneficiation processes in the mining industry emit flue gas and particulate matter comprising SO₂, among other toxic substances [4]. Wind dispersion aids the mobilization of these potentially toxic elements (PTEs) from mining operations into the environment and leads to the contamination of air, surface soil and water bodies [5, 6]. Some plants take up these PTEs in excess, resulting in deleterious effects on the health and productivity of vegetation [7, 8].

SO₂ enters plant tissues through the lenticels, leaves, and roots [9]. In the plant tissues (mesophyll cells), it reacts with water molecules and converts to Sulphate (SO₄²⁻) or Sulphite (SO₃²⁻) ions [10, 11]. However, excess SO₃²⁻ is highly toxic to plants, affecting plant growth and productivity [12]. The detoxification process of SO₃²⁻ to SO₄²⁻ increases the generation of reactive oxygen species (ROS) such as superoxide radicals (O₂⁻) and hydrogen peroxide-H₂O₂ [13]. In addition, stomatal closure triggers an insufficient concentration of intercellular CO₂, instigating the formation of singlet atoms (¹O₂) and hydroxyl radicals (OH[•]), other family members of destructive ROS [14–16]. According to Mansoor et al. [14], ROS are highly toxic and destructive to biomolecules such as lipids, proteins, nucleic acids, and carbohydrates.

¹O₂ severely damages nucleic acids, proteins, pigments, and lipids [17], jeopardizing photosynthetic and photosystems machinery [16]. O₂⁻ transforms into the more reactive and toxic OH[•] and ¹O₂, causing lipid peroxidation [18]. OH[•] is the most toxic and reactive among the members of the ROS family [15], destroying cellular compartments through, lipid peroxidation, membrane damage and protein destruction [17, 18]. Zentgraf et al. [19] noted that the longer half-life of H₂O₂ enables it to traverse long distances and cross cell membranes through aquaporins, covering significant lengths within the cell and causing oxidative damage. They further argued that H₂O₂ is responsible for programmed cell death and 50% loss of enzymatic activities at a high cellular concentration. Therefore, extensive SO₂ exposure affects morphological and biochemical activities resulting in reduced plant growth and forest productivity [20].

Field-based studies have investigated the potential of using SO₂ and plants in detecting industrial pollution-instigated environmental change by evaluating the concentration of Sulphur in different tree species and its associated effect on tree growth and wood cell characteristics [21, 22]. Sensula [23] observed an increased concentration of Sulphur in *Pinus sylvestris* growing near industrial areas and reported reduced annual ring width attributed to SO₂. In addition, Likus-Cieslik et al. [21] recorded a higher concentration of Sulphur in the leaves of *P. sylvestris* growing near an industrial site and reported a significant correlation between the concentration of Sulphur in needles and stand defoliation degrees.

Thus far, there is limited information on the dendrochemical potential of Miombo woodland species in detecting changes in industrial pollution. This study seeks to unravel the dendrochemical potential of *Brachystegia longifolia* in detecting environmental change induced by variations in industrial copper mining activities around the Miombo woodlands. This was achieved through (i) determining the Sulphur concentration in tree rings, representing annual deposition of SO₂ in *B. longifolia*, (ii) dating the Sulphur concentration in annual tree rings, (iii) identifying variations in SO₂ deposition before and after the installation of a modern smelter and iv) determining the effect of SO₂ variations on the wood density of *B. longifolia*.

Characterized by *Julbernardia*, *Isoberrlinia* and *Brachystegia* species, Miombo woodlands extend widely in the Sub-Saharan Africa covering 2.7 million km² of Africa's landmass, representing 9% of its total area [24]. This forest type is extensively distributed in Angola, Mozambique, Malawi, the Democratic Republic of Congo, Zambia, Zimbabwe and Tanzania. Most of these countries are endowed with mineral resources where mineral extraction and beneficiation

activities have been carried out commercially for close to a century. In Zambia, at least 80% of forest cover is classified as Miombo woodland [25]. However, the potential impacts of mining activities on the Miombo woodland species and forest productivity are not well established.

2 Materials and methods

2.1 Study area description

Mufulira is a copper mining town in Zambia, highly polluted with SO_2 [3] and copper [26]. The principal source of pollution is a copper smelter (Fig. 1) located in Kankoyo township with its associated facilities such as the converter slag blow, converter copper blow, and matte setting furnace [27].

Commissioned in 1936, Mufulira smelter is the oldest and largest in Zambia and treats concentrates from the Copperbelt region, Northwestern Zambia, and in-house mining operations [26]. This facility was replaced with a modern Isasmelter in 2014 [28]. The new smelter converts more than 97% of SO_2 produced at this mining facility into H_2SO_4 . However, there is limited environmental evidence on the comparative variations in SO_2 emissions in Mufulira before and after the smelter upgrade. Therefore, this study investigated the impact of copper mining activities on the Miombo woodlands by assessing the potential of *B. longifolia* in detecting industrial pollution dynamics through dendrochemistry and analyzing Sulphur distribution in tree rings formed before and after the replacement of the obsolete smelter. *B. longifolia* was preferred among the characteristic Miombo species because of previous studies demonstrating that *Brachystegia* species form annual growth rings [29], which is critical in Dendrochronological studies.




Fig. 1 Study area Map showing the location of the copper smelter and the sampling plot in Mufulira, Zambia

The study area recorded 21 °C mean annual temperature and 1147 mm mean annual precipitation between 1960 and 2016 [30].

2.2 Sampling and justification

Wood discs were collected from two *B. longifolia* trees (NNA and NNC) naturally growing from the nearest forest patch, 1.8 km from the copper smelter (Fig. 1). Each disc then produced two samples (NNA-A, NNA-B, NNC-A and NNC-B) for Sulphur and wood density analysis, giving a total of four samples. The limited number of sampled trees was because of the destructive nature of the sampling method. Furthermore, Cherubini et al. [31] and Mifsud et al. [32] guided that sample location and sample extraction height along the tree trunk as well as the number of samples per tree are the most important parameters to consider when sampling for dendrochemical studies. This is because the correlation between environmental change drivers and elemental concentrations in tree rings offers great annually and intra-annually resolved proxy potential as the average concentration of the studied element(s) is drawn from hundreds of data points for a single year depending on the width of the growth band/ring (Fig. 2) and trees are statically available overtime [31, 33–35].

Therefore, studies [35–38] that have sampled discs using destructive sampling for dendrochemistry usually aim at sampling the bare minimum to reduce a contribution to deforestation. Cobanoglu et al. [37], Scharnweber et al. [35] and Battipaglia & Cherubini [36] sampled one tree each to study the effect of traffic density and mineral extraction on the concentration of trace elements in tree rings. Conversely, Fornasaro et al. [38] and Luong et al. [39] sampled three trees for similar objectives, without any reported significant differences in the overall results between and among studies with different sample sizes.

2.3 Sample preparation and analysis of Sulphur distribution and wood density in tree rings

Sample preparation: Wood discs were extracted at breast height (≈ 1.3 m) of the selected trees (Fig. 2), air dried and sanded progressively using 80 to 600 grits until the transverse surface and annual rings were clearly visible. Two radial samples of each disc were cut for the analysis of Sulphur and wood density. The radial samples were glued on a wood support and slides were cut transversely (1.7 mm thick) with a double circular saw. Radial wood slices were then kept at 20 °C and 60% relative humidity until reaching a stable moisture content of 12% (~ 24 h) [40].




Fig. 2 Sample preparation and analysis of Sulphur distribution and wood density in *B. longifolia* trees

Sulphur distribution analysis: Radial slices were linearly scanned along the bark-pith transverse surface. Analyses were performed with a μ -XRF spectrometer (Bruker, M4 Tornado, USA), with an Rh tube (50 kV, 600 μ A) and a silicon drift detector. Samples were exposed to the X-ray beam with 25 μ m width for 100 ms at each measurement point (dead time of 15%) in the radial direction (25 μ m step intervals) and in vacuum condition (< 0.5 Torr). X-ray transmission intensity values of counts per second (cps) for Sulphur were analysed to distinguish the detected signal from the background with reasonable certainty for the analytical quantification and the Sulphur profiles in cps were finally built.

Wood density analysis: The same radial slices were then scanned with a cellulose acetate calibration scale using an X-ray densitometry chamber (Faxitron MX20-DC12, Faxitron X-Ray, IL, USA) [41]. Digital images (TIFF, 513 dpi) were analysed with the Windendro[®] program, obtaining wood density profiles in g/cm³ with 15 μ m step intervals. The position of the latewood-earlywood boundary was defined by Mulenga et al. [29]. Then annual time series of average Sulphur and wood density for each radial sample were produced.

2.4 Dating of Sulphur distribution and wood density

The sampling year (November 2020) served as the basis for dating tree rings. Annual growth rings were counted and the latest complete growth band was allocated to the previous full growth season (2019 winter plus summer), before the sampling year. Dating of wood samples was carried out systematically starting from the bark towards the pith. Thereafter, Sulphur distribution and wood density in *B. longifolia* trees were dated by allocating respective values according to the year when that tree ring was formed.

2.5 Statistical analysis

A simple linear regression analysis was conducted to determine whether the average wood density and average Sulphur concentration in tree rings significantly varied with time, after satisfying the requirements for a regression test. The test was carried out in two segments. The first analysed data before 2014 while the second one was for data after 2014. This was done to compare trends and the associated significance levels at $p < 0.05$ between the two time periods. All the graphics and statistics were produced using an Integrated Development Environment for R (RStudio IDE) which is a programming language for statistical computing and graphics [42].

3 Results and discussion

3.1 Sulphur distribution in *B. longifolia* tree rings

The number of annual growth rings demonstrated that the studied trees were aged 25 and 11 years old. Trees NNA and NNC started growing in 1994 and 2008, respectively and were both sampled at the same time in November 2020. The mean concentration of Sulphur in tree rings that developed in 2014 and 2019 were 17.51 cps and 13.28 cps, respectively (Table 1).


The distribution of Sulphur in tree rings formed before 2014 increased significantly ($p < 0.05$) with tree age or time (Fig. 3). Conversely, significant ($p < 0.05$) reduction of Sulphur concentration in *B. longifolia* tree rings was observed after 2014. These trends suggest significant differences in the concentration of Sulphur or Sulphur compounds in the atmosphere or soil before and after 2014. This can be attributed to the SO₂ emissions from the copper smelter as this is the only smelter within the radius of approximately, 30 km. Chambishi copper smelter is the nearest located in the downwind

Table 1 Concentration of Sulphur (cps) in *B. longifolia* tree rings in 2014 and 2019

Sample ID	2014	2019
NNC_A	15.14	14.01
NNC_B	12.52	14.48
NNA_A	22.26	16.64
NNA_B	20.11	8.00
Mean	17.51	13.28

Bold shows the average sulphur content and wood density, respectively

Fig. 3 Dated mean Sulphur distribution in *B. longifolia* trees naturally growing near a copper smelter in Mufulira, Zambia. The dating provides significant milestones for the copper smelter upgrade project



direction. Muma et al. [3], reported that SO_2 and heavy metals from stack emissions at the copper mine heavily pollute Mufulira and the surrounding landscapes. SO_2 levels exceeding the acceptable limit of $50 \mu\text{g}/\text{m}^3$ annual exposure were reported within a radius of 12 km downwind and 4 km upwind. These exposure levels far exceed the EQG to protect fauna and flora.

The project to replace and upgrade a copper smelter aimed at minimizing environmental footprint by abetting the dispersion of SO_2 emissions into the atmosphere was completed around this period [28]. Briefly, the smelter upgrade project started in 2007 and was completed in 2014 [28]. The facility was by 2009 able to capture up to 51% of SO_2 from the mineral beneficiation processes and up to 97% at the completion of the project in 2014 [3, 28]. Therefore, the reduction of Sulphur distribution and trend in *B. longifolia* tree rings after 2014 can be attributed to a decrease in SO_2 emissions. Conversely, the increasing trend of Sulphur in tree rings before 2014 could have been instigated by increasing SO_2 dispersion triggered by increased copper production [43].

The mean Sulphur concentration in tree rings at the beginning and completion of the smelter upgrades were 15.41 cps and 17.51 cps, respectively. By 2019, the mean Sulphur content in *B. longifolia* wood had significantly reduced to 13.28 cps. A slight lag and downward trend in the distribution of Sulphur after the completion of the smelter upgrade can be attributed to residual (carry-over) Sulphur in the soil which was ultimately absorbed through the roots in the subsequent years.

Muma et al. [3] argued that although the smelting activities increased in the last two decades, atmospheric SO_2 recorded at approximately 1 km downwind of the smelter, reduced significantly between 2014 and 2018 from 247 to $105 \mu\text{g}/\text{m}^3$. It would be expected that if the production capacity of the smelter increased after the upgrades, then SO_2 emissions would increase as well, potentially increasing the concentration of Sulphur in tree rings. In contrast, Muma et al. [3] observed decreased levels of atmospheric SO_2 a kilometre away, downwind of the copper smelter. Our study shows that the SO_2 footprint in tree rings is reduced, suggesting that a significant amount of this pollutant is being effectively captured by the modern smelter. Therefore, its emission into the atmosphere has significantly reduced, achieving the aim of the copper mining operator, which was to reduce the emission of SO_2 by converting it into H_2SO_4 and capturing it for industrial use [43].

Similar results have been reported elsewhere in different plant species, including *P. monophylla* [44], *P. pinea* [45], *P. elderica* [46], *Lax occidentalis* [47], *Picea abies* and *P. sylvestris* [48], and *Quercus pubescens* [49]. These species recorded increased concentrations of Sulphur in different biomass samples including tree rings at the peaks of SO_2 emanating from varying industrial activities.

3.2 Wood density in *B. longifolia* tree rings

On average, $0.78 \text{ g}/\text{cm}^3$ and $0.75 \text{ g}/\text{cm}^3$ wood density were recorded in annual rings at the beginning and completion of the upgrades, respectively (Table 2). However, the mean wood density had significantly reduced to $0.67 \text{ g}/\text{cm}^3$ by 2019.

Table 2 Density (g/cm³) of *B. longifolia* wood formed in 2014 and 2019

Sample ID	2014	2019
NNC_A	0.71	0.67
NNC_B	0.71	0.61
NNA_A	0.84	0.74
NNA_B	0.72	0.66
Mean	0.75	0.67

Bold shows the average sulphur content and wood density, respectively

A significant ($p < 0.05$) increase in wood density was observed in tree NNA with increasing age, before 2014 and thereafter decreased, significantly ($p < 0.05$), Fig. 4. Conversely, wood density for samples extracted from tree NNC decreased significantly ($p < 0.05$) before and after 2014.


Wood density increased significantly with increasing tree age before the upgrading of the smelter in tree NNA (Fig. 4). Conversely, a significant reduction in wood density with increasing tree age was observed from 2009 when the smelter upgrades could capture 51% of SO₂. This downward trend in wood density is observed in both trees, NNA and NNC. The observed changes in the density of *B. longifolia* trees before and after the upgrading of the copper smelter can be attributed to variations in the copper mining-induced toxic emissions. Emissions such as SO₂ may induce variations in tree growth rate [21, 22, 29], resulting in the changes in wood density of *B. longifolia* trees [29, 50–54].

A number of studies have evaluated the effect of tree growth rate on the wood density of several tree species including *Quercus petraea* [51] and *Acacia auriculiformis* [53]. All these studies reported a gradually decreasing wood density with an increasing tree growth rate. A higher growth rate induces increased cell lumen dimensions and increased cell wall thickness [55–57] critical for wood density. Werdin et al. [58] reported that wood density is significantly influenced by wood cell characteristics. They observed higher wood density in species with thinner cell walls and greater fibre diameters but recorded lower density in those with thicker cell walls and smaller fibre diameters. Therefore, the observed wood density in these studies was attributed to increased cell lumen diameters with significantly increased cell wall thickness formed at a higher growth rate.

Other studies have examined the impact of SO₂ phytotoxicity and reported significant increments in wood cell lumen diameter and reduced cell wall thickness in *Calendula officinalis* [59], *Pinus elderica* [46], *Mangifera indica* [60], *Prosopis cineraria* [61], *Syzygium cumini* [61], and *Ailanthus excelsa* [62].

Therefore, the observed changes in wood density before the smelter upgrades in our site can be attributed to the low tree growth rate instigated by copper mining-induced SO₂ phytotoxicity. Conversely, the reduction in wood density after 51% and 97% capture of SO₂ can be attributed to increased tree growth. It is assumed that the growth rate picks up

Fig. 4 Dated mean wood density of *B. longifolia* prior to and after replacement and upgrading of the copper smelter. The dating provides significant milestones for the copper smelter upgrade project



resulting in the formation of reduced cell lumen diameters and increased cell wall thickness after the significant capture of SO₂, leading to the demonstrated reduction in the density of *B. longifolia* wood (Fig. 4).

3.3 Study implication on the utilization potential of *B. longifolia* wood formed in a SO₂ polluted environments

B. longifolia is one of the most commercially valuable Miombo woodland species in Zambia and Southern Africa. Species of the genera *Brachystegia*, *Isoberlinia*, and *Julbernardia* are characteristic trees of the Miombo [63] and extensively spread across all mining countries, including Zambia, Angola, Malawi, the Democratic Republic of Congo (DRC), Tanzania, Zimbabwe, and Mozambique [24]. These tree species are widely used for furniture manufacturing, housing construction and bioenergy production [64]. The Miombo woodland is the most extensive vegetation type in Zambia covering over 80% of forests and accounts for 60% of Zambia's land mass [25]. This vegetation type covers 9% of Africa's total land area, covering over 2.7 million km² [65].

In Zambia, most mining activities are surrounded by forests characterized by Miombo species. Given the impact of environmental pollution on tree growth and forest productivity [21, 22, 29], there is a need to examine and exploit the biomass value chain to maximize the utilization potential of forests to sustain the bioeconomy. Potent economic opportunities are feasible for high-value products including bioenergy products, biopolymers, biocomposites, pulp, and chemical products. However, the utilization potential of biomass resources depends on characteristics such as density, chemical composition, machinability, and mechanical properties [66]. Therefore, the variations in wood density and Sulphur concentration observed in this study suggest that the utilization potential of *B. longifolia* trees and possibly other Miombo woodland species naturally growing in mining environments is limited.

Generally, this study suggests opportunities for multidisciplinary efforts exploring how to exploit and maximize the utilization potential of biomass generated from polluted environments, considering the expected high ash content instigated by high concentrations of Sulphur as demonstrated in this study. For instance, contaminated biomass with high-quality cellulose can be used to produce liquid-based biofuels [67, 68] and then utilize biomass remnants to manufacture lignocellulosic composites, among other innovative bio-based products. These multidisciplinary collaborations would unlock viable economic opportunities through the utilization of biomass produced in polluted environments advancing the cause of a circular economy.

Additionally, this study demonstrates that it is technically feasible for mining firms to take initiatives that minimize and mitigate the impact of mining activities on the environment. Furthermore, our study shows that the Miombo woodlands are vulnerable to environmental stresses instigated by anthropogenic activities such as mining and other industrial activities.

4 Conclusion

This study investigated the dendrochemical potential of *B. longifolia* in detecting environmental change induced by variations in industrial pollution load. The results show a significant ($p < 0.05$) reduction in the distribution of Sulphur after the completion of smelter upgrades. The mean Sulphur concentration in tree rings at the beginning (2007) and completion (2014) of the smelter upgrades were 15.41 cps and 17.51 cps, respectively. By 2019, the highest Sulphur content in *B. longifolia* wood had significantly reduced to 13.28 cps. Furthermore, a significant reduction in wood density with increasing tree age was observed after the installation of the new smelter. 0.78 g/cm³ and 0.75 g/cm³ mean wood density were recorded in annual rings at the beginning and completion of the upgrades, respectively. However, the mean wood density had significantly reduced to 0.67 g/cm³ by 2019. The significant variations in the distribution of Sulphur and wood density in tree rings formed before and after upgrading an obsolete copper smelter demonstrates the dendrochemical potential of *B. longifolia* and possibly other miombo woodland species, offering an opportunity to biomonitor the impact of industrial activities in the miombo landscapes which are extensively distributed across sub-Saharan Africa.

Author contributions C.M. and C.A.L conceptualized the study and collected samples. D.R.O.R and HDB carried out laboratory sample analysis. C.M, D.R.O.R, HDB, D.P, C.A.L and J.H.S designed the methodology. S.A.M. designed Fig. 3. D.P. designed Fig. 1. C.M, D.R.O.R, H.D.B, D.P, C.A.L and J.H.S analyzed data and interpreted the results. C.M, DROR, HDB, D.P, C.A.L and J.H.S drafted the manuscript. JHS edited the final manuscript. All authors reviewed the manuscript.

Funding This research work did not receive any form of funding.

Data availability The data used during the current study available from the corresponding author on reasonable request.

Declarations

Ethics approval and consent to participate Not applicable.

Plant guidelines The authors confirm that the use of plants in the present study complies with international, national, and/or institutional guidelines.

Wild plants *Brachystegia longifolia* Benth was collected in Zambia by Royal Botanical Gardens. The plant material was identified by Boume. R, and a voucher specimen was deposited at Royal Botanic Gardens, Kew—Herbarium Specimens with ID K Herbarium K000417827.

Consent for publication Not applicable.

Competing interests The authors declare no competing interests.

Open Access This article is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License, which permits any non-commercial use, sharing, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if you modified the licensed material. You do not have permission under this licence to share adapted material derived from this article or parts of it. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by-nc-nd/4.0/>.

References

1. Mulenga C. Soil governance and the control of mining pollution in Zambia. *Soil Secur.* 2022;6:100039. <https://doi.org/10.1016/j.soisec.2022.100039>.
2. Dusengemungu L, Mubemba B, Gwanama C. Evaluation of heavy metal contamination in copper mine tailing soils of Kitwe and Mufulira, Zambia, for reclamation prospects. *Sci Rep.* 2022;12:11283. <https://doi.org/10.1038/s41598-022-15458-2>.
3. Muma D, Besa B, Manchisi J, Banda W. Effects of mining operations on air and water quality in Mufulira district of Zambia: a case study of Kankoyo Township. *J South Afr Inst Min Metall.* 2020;120:287–98.
4. Kasongo J, Alleman LY, Kanda JM, Kaniki A, Riffault V. Metal-bearing airborne particles from mining activities: a review on their characteristics, impacts and research perspectives. *Sci Total Environ.* 2024;951:175426.
5. Lotfi S, Chakit M, Belghyti D. Groundwater quality and pollution index for heavy metals in Sais plain, Morocco. *J Heal Pollut.* 2020;10(26):1–12.
6. Bouzekri S, El Hachimi ML, Kara K, El Mahi M, Lotfi EM. Metal pollution assessment of surface water from the abandoned Pb mine Zaida, high Moulouya-Morocco. *Geosystem Eng.* 2020;23(4):226–33. <https://doi.org/10.1080/12269328.2020.1772125>.
7. Song J, Sun Z, Saud S, Fahad S, Nawaz T. Exploring the deleterious effects of heavy metal cadmium on antioxidant defense and photosynthetic pathways in higher plants. *Plant Stress.* 2025;15:100716.
8. Sperdouli I. Heavy metal toxicity effects on plants. *Toxics.* 2022;10(12):1–4.
9. Bethke PC. Potato tuber lenticels: a review of their development, structure, function, and disease susceptibility. *Am J Potato Res.* 2023;100(4):253–64.
10. Li ZG, Li XE, Chen HY. Sulfur dioxide: an emerging signaling molecule in plants. *Front Plant Sci.* 2022;13:1–9.
11. Ren Z, Wang RY, Huang XY, Wang Y. Sulfur compounds in regulation of stomatal movement. *Front Plant Sci.* 2022;13:2016–23.
12. Sharma U, Bekturova A, Ventura Y, Sagi M. Sulfite oxidase activity level determines the sulfite toxicity effect in leaves and fruits of tomato plants. *Agronomy.* 2020;10(5):694.
13. Afzal S, Abdul Manap AS, Attiq A, Albokhadaim I, Kandeel M, Alhojaily SM. From imbalance to impairment: the central role of reactive oxygen species in oxidative stress-induced disorders and therapeutic exploration. *Front Pharmacol.* 2023;14:1–22.
14. Mansoor S, Wani OA, Lone JK, Manhas S, Kour N, Alam P, et al. Reactive oxygen species in plants: from source to sink. *Antioxidants.* 2022;11(2):1–14.
15. Sachdev S, Ansari SA, Ansari MI, Fujita M, Hasanuzzaman M. Abiotic stress and reactive oxygen species: generation, signaling, and defense mechanisms. *Antioxidants.* 2021;10(2):1–37.
16. Dumanović J, Nepovimova E, Natić M, Kuča K, Jačević V. The significance of reactive oxygen species and antioxidant defense system in plants: a concise overview. *Front Plant Sci.* 2021;11:552969.
17. Dogra V, Kim C. Singlet oxygen metabolism: from genesis to signaling. *Front Plant Sci.* 2020;10:1640.
18. Andrés Juan C, Manuel Pérez de la Lastra J, Plou FJ, Pérez-Lebeña E, Reinbothe S. Molecular sciences the chemistry of reactive oxygen species (ROS) revisited: outlining their role in biological macromolecules (DNA, lipids and proteins) and induced pathologies. *Int J Mol Sci.* 2021;22:4642. <https://doi.org/10.3390/ijms>.

19. Zentgraf U, Andrade-Galan AG, Bieker S. Specificity of H₂O₂ signaling in leaf senescence: is the ratio of H₂O₂ contents in different cellular compartments sensed in *Arabidopsis* plants? *Cell Mol Biol Lett*. 2022;27(1):4.
20. Daldoul G, Souissi R, Tili H, Elbahri D, El Hamiani O, Chebbi N, et al. Assessment of heavy metal toxicity in soils contaminated by a former Pb–Zn mine and tailings management using flotation process, Jebel Ghazlane, Northern Tunisia. *Environ Earth Sci*. 2019;78(24):703.
21. Likus-Cieślak J, Socha J, Gruba P, Pietrzykowski M. The current state of environmental pollution with sulfur dioxide (SO₂) in Poland based on Sulfur concentration in scots pine needles. *Environ Pollut*. 2020;258:113559. <https://doi.org/10.1016/j.envpol.2019.113559>.
22. Padbhushan R, Kumar U, Sharma S, Rana DS, Kumar R, Kohli A, et al. Impact of land-use changes on soil properties and carbon pools in India: a meta-analysis. *Front Environ Sci*. 2022;9:1–15.
23. Sensula BM. The impact of climate, sulfur dioxide, and industrial dust on δ18O and δ13C in glucose from pine tree rings growing in an industrialized area in the Southern part of Poland. *Water Air Soil Pollut*. 2016;227(4):1–13.
24. Jimu L, Mataruse L, Musemwa L, Nyakudya IW. The miombo ecoregion up in smoke: the effect of tobacco curing. *World Dev Perspect*. 2017;5:44–6. <https://doi.org/10.1016/j.wdp.2017.03.007>.
25. Chidumayo EN. Is charcoal production in *Brachystegia-Julbernardia* woodlands of Zambia sustainable? *Biomass Bioenergy*. 2019;125:1–7.
26. Mulenga C, Clarke C, Meincken M. Bioaccumulation of Cu, Fe, Mn and Zn in native *Brachystegia longifolia* naturally growing in a copper mining environment of Mufulira, Zambia. *Environ Monit Assess*. 2022;194(1):1–13. <https://doi.org/10.1007/s10661-021-09656-0>.
27. Mwaanga P, Silondwa M, Kasali G, Banda PM. Preliminary review of mine air pollution in Zambia. *Heliyon*. 2019;5(9):e02485. <https://doi.org/10.1016/j.heliyon.2019.e02485>.
28. Kunda J. Mopani mine smelter upgrade almost done. *Times of Zambia*. 2014.
29. Mulenga C, Clarke C, Meincken M. Effect of copper mining pollution-induced heavy metal toxicities on *B. longifolia* Benth wood cell characteristics. *Eur J For Res*. 2022. <https://doi.org/10.1007/s10342-022-01524-x>.
30. Musonda B, Jing Y, Iyakaremye V, Ojara M. Analysis of long-term variations of drought characteristics using standardized precipitation index over Zambia. *Atmosphere (Basel)*. 2020;11(12):1–20.
31. Cherubini P, Gargano A, Grob M, Nievergelt D, Passardi S. Dendrochronologia's tutoring recipes how to take samples for small basic dendroecological studies. *Dendrochronologia*. 2020;64:125774.
32. Mifsud DV, Stüeken EE, Wilson RJS. A preliminary study into the use of tree-ring and foliar geochemistry as bio-indicators for vehicular NO_x pollution in Malta. *Isotopes Environ Health Stud*. 2021;57(3):301–15. <https://doi.org/10.1080/10256016.2021.1902319>.
33. Ayan S, Sarsekova D, Kenesaryuly G, Yilmaz E, Gülseven O, Şahin İ. Accumulation of heavy metal pollution caused by traffic in forest trees in the park of Kerey and Janibek Khans of the city of Nur-Sultan, Kazakhstan. *J For Sci*. 2021;67(7):357–66.
34. Gavrikov VL, Fertikov AI, Vidus VE, Sharafutdinov RA, Vaganov EA. Elemental variability in stems of *Pinus sylvestris* L.: whether a single core can represent all the stem. *Diversity*. 2023;15(2):281.
35. Scharnweber T, Rocha E, González Arrojo A, Ahlgrimm S, Gunnarson BE, Holzkämper S, et al. To extract or not to extract? Influence of chemical extraction treatment of wood samples on element concentrations in tree-rings measured by X-ray fluorescence. *Front Environ Sci*. 2023;11:1–10.
36. Battipaglia G, Cherubini P. Stable Isotopes in tree rings of Mediterranean forests. *Cham: Springer*; 2022. p. 605–29.
37. Cobanoglu H, Sevik H, Koç İ. Do annual rings really reveal Cd, Ni, and Zn pollution in the air related to traffic density? An example of the cedar tree. *Water Air Soil Pollut*. 2023;234(2):1–10.
38. Fornasaro S, Ciani F, Nannoni A, Morelli G, Rimondi V, Lattanzi P, et al. Tree rings record of long-term atmospheric Hg pollution in the monte Amiata mining district (Central Italy): lessons from the past for a better future. *Minerals*. 2023;13(5):688.
39. Luong TH, Jang KS, Lim HW, Choi WJ, Lee KH. Correlation of tree ring growths of four major species with climate changes in South Korea. *Forest Sci Technol*. 2013;9(4):180–6. <https://doi.org/10.1080/21580103.2013.839278>.
40. Teodorescu I, Erbasu R, Branco JM, Tapusi D. Study in the changes of the moisture content in wood. *IOP Conf Ser Earth Environ Sci*. 2021;664(1):012017.
41. Quintilhan MT, Santini L, Ortega-Rodriguez DR, Guillemot J, Cesilio GHM, Chambi-Legoas R, et al. Growth-ring boundaries of tropical tree species: aiding delimitation by long histological sections and wood density profiles. *Dendrochronologia*. 2021;69:125878.
42. R Core Team. R: a language and environment for statistical computing. Vienna: R Core Team; 2021.
43. Auditor General. Management of environmental degradation caused by mining activities in Zambia. 2014.
44. Thompson MA. Tree rings and air pollution: a case study of *Pinus monophylla* growing in east-central Nevada. *Environ Pollut Ser A Ecol Biol*. 1981;26(4):251–66.
45. Battipaglia G, Marzaioli F, Lubritto C, Altieri S, Strumia S, Cherubini P, et al. Traffic pollution affects tree-ring width and isotopic composition of *Pinus pinea*. *Sci Total Environ*. 2010;408(3):586–93. <https://doi.org/10.1016/j.scitotenv.2009.09.036>.
46. Safdari V, Ahmed M, Devall MS, Bayramzadeh V. Effects of air pollution on morphological and anatomical characteristics of *Pinus eldarica* wood. *Fuust J Biol*. 2012;2(2):5–12.
47. Fox CA, Kincaid WB, Nash TH III, Young DL, Fritts HC. Tree-ring variation in western larch (*Larix occidentalis*) exposed to Sulfur dioxide emissions. *Can J For Res*. 1986;16(2):283–92. <https://doi.org/10.1139/x86-049>.
48. Seftigen K, Moldan F, Linderholm HW. Radial growth of Norway spruce and scots pine: effects of nitrogen deposition experiments. *Eur J For Res*. 2013;132(1):83–92.
49. Perone A, Cocozza C, Cherubini P, Bachmann O, Guillon M, Lasserre B, et al. Oak tree-rings record spatial-temporal pollution trends from different sources in Terni (Central Italy). *Environ Pollut*. 2018;233:278–89. <https://doi.org/10.1016/j.envpol.2017.10.062>.
50. Mulenga C. Growth response of *Brachystegia longifolia* to copper mining pollution-induced heavy metal toxicity [Internet]. Stellenbosch University; 2022. Available from: <https://scholar.sun.ac.za/handle/10019.1/125906>
51. Arsić J, Stojanović M, Petrovičová L, Noyer E, Milanović S, Světlík J, et al. Increased wood biomass growth is associated with lower wood density in *Quercus petraea* (Matt.) Liebl. saplings growing under elevated CO₂. *PLoS ONE*. 2021;16:1–20.
52. Diao S, Sun H, Forrester DI, Soares AAV, Protásio TP, Jiang J. Variation in growth, wood density, and stem taper along the stem in self-thinning stands of *Sassafras tzumu*. *Front Plant Sci*. 2022;13:1–11.
53. Féty Tonouéwa JFM, Biaou SSH, Assédé ESP, Langbour P, Balagueman OR. Influence of growth parameters on wood density of acacia auriculiformis. *Maderas Cienc Tecnol*. 2022;24(24):1–14.

54. Zhang SY, Ren H, Jiang Z. Wood density and wood shrinkage in relation to initial spacing and tree growth in black spruce (*Picea mariana*). *J Wood Sci*. 2021;67(1):1–10. <https://doi.org/10.1186/s10086-021-01965-9>.
55. Kašpar J, Šamonil P, Krůček M, Vašíčková I, Daněk P. Hillslope processes affect vessel lumen area and tree dimensions. *Front Plant Sci*. 2021;12:1–11.
56. Silvestro R, Zeng Q, Buttò V, Sylvain JD, Drolet G, Mencuccini M, et al. A longer wood growing season does not lead to higher carbon sequestration. *Sci Rep*. 2023;13(1):1–12. <https://doi.org/10.1038/s41598-023-31336-x>.
57. Vieira J, Carvalho A, Campelo F. Tree growth under climate change: evidence from Xylogenesis timings and kinetics. *Front Plant Sci*. 2020;11:1–11.
58. Werdin J, Fletcher TD, Rayner JP, Williams NSG, Farrell C. Biochar made from low density wood has greater plant available water than biochar made from high density wood. *Sci Total Environ*. 2020;705:135856. <https://doi.org/10.1016/j.scitotenv.2019.135856>.
59. Wali B, Iqbal M, Mahmooduzzafar. Anatomical and functional responses of *Calendula officinalis* L. to SO₂ stress as observed at different stages of plant development. *Flora Morphol Distrib Funct Ecol Plants*. 2007;202(4):268–80.
60. Iqbal M, Mahmooduzzafar, Aref IM, Khan PR, Iqbal M, Aref IM, et al. Behavioral responses of leaves and vascular cambium of *Prosopis cineraria* (L.) Druce to different regimes of coal-smoke pollution. *J Plant Interact*. 2010;5(2):117–33.
61. Hegazy SS, Aref IM, Iqbal M, Mahmooduzzafar, Hegazy SS, Aref IM, et al. Anatomical changes in the wood of *Syzygium cumini* exposed to coal-smoke pollution. *J Food Agric Environ*. 2010;8(3–4 Part 2):959–64.
62. Rajput KS, Rao KS. Cambial periodicity and formation of wood in *Ailanthus excelsa* growing under the influence of combined air pollutants. *Phyt Ann Rei Bot*. 2005;45(1):51–64.
63. Gonçalves FMP, Revermann R, Gomes AL, Aidar MPM, Finckh M, Juergens N. Tree species diversity and composition of miombo woodlands in south-central Angola: a chronosequence of forest recovery after shifting cultivation. *Int J For Res*. 2017;2017:6202093.
64. Montfort F, Nourtier M, Grinand C, Maneau S, Mercier C, Roelens JB, et al. Regeneration capacities of woody species biodiversity and soil properties in miombo woodland after slash-and-burn agriculture in Mozambique. *For Ecol Manag*. 2021;488:119039.
65. Sikuzani YU, Muteya HK, Bogaert J. Miombo woodland, an ecosystem at risk of disappearance in the Lufira Biosphere Reserve (Upper Katanga, DR Congo)? A 39-years analysis based on Landsat images. *Glob Ecol Conserv*. 2020;24:e01333.
66. Kusiak W, Majka J, Ratajczak I, Górka M, Zborowska M. Evaluation of environmental impact on selected properties of lime (*Tilia cordata* Mill.) wood. *Forests*. 2020;11(11):746.
67. Edgar VN, Fabián FL, Julián Mario PC, Ileana VR. Coupling plant biomass derived from phytoremediation of potential toxic-metal-polluted soils to bioenergy production and high-value by-products-a review. *Appl Sci*. 2021;11(7):2982.
68. Lee J, Kaunda RB, Sinkala T, Workman CF, Bazilian MD, Clough G. Phytoremediation and phytoextraction in Sub-Saharan Africa: addressing economic and social challenges. *Ecotoxicol Environ Saf*. 2021;226:112864. <https://doi.org/10.1016/j.ecoenv.2021.112864>.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.