

**UNIVERSIDADE ESTADUAL PAULISTA “JÚLIO DE MESQUITA
FILHO” FACULDADE DE CIÊNCIAS AGRÁRIAS E VETERINÁRIAS
CAMPUS JABOTICABAL**

**DIFERENÇAS SEXUAIS REFERENTES À PARTICIPAÇÃO DO
CANAL CATIÔNICO ASIC1a NO *LOCUS COERULEUS* NA
RESPOSTA COMPORTAMENTAL INDUZIDA POR CO₂**

Letícia Rodrigues Pinheiro

Bióloga

2024

**UNIVERSIDADE ESTADUAL PAULISTA - UNESP CÂMPUS DE
JABOTICABAL**

**DIFERENÇAS SEXUAIS REFERENTES À PARTICIPAÇÃO DO
CANAL CATIÔNICO ASIC1a NO *LOCUS COERULEUS* NA
RESPOSTA COMPORTAMENTAL INDUZIDA POR CO₂**

Letícia Rodrigues Pinheiro

Orientadora: Profa. Dra. Luciane Helena Gargaglioni Batalhão

Coorientadora: Dra. Alana Tercino Frias

Dissertação apresentada ao Programa Institucional de Pós-Graduação em Ciência Animal da Universidade Estadual Paulista “Júlio de Mesquita Filho” como parte dos requisitos para obtenção do título de Mestre em Ciência Animal.

P654d Pinheiro, Letícia Rodrigues
Diferenças sexuais referentes à participação do canal catiônico ASIC1a no
Locus coeruleus na resposta comportamental induzida por CO2 / Letícia
Rodrigues Pinheiro. – Jaboticabal, 2024
47 p.

Dissertação (mestrado) - Universidade Estadual Paulista (UNESP),
Faculdade de Ciências Agrárias e Veterinárias, Jaboticabal
Orientadora: Luciane Helena Gargaglioni Batalhão
Coorientadora: Alana Tercino Frias

1. Fisiologia. 2. Ventilação. 3. Comportamento. I. Título.

Sistema de geração automática de fichas catalográficas da Unesp. Biblioteca da Universidade Estadual Paulista (UNESP), Faculdade de Ciências Agrárias e Veterinárias, Jaboticabal. Dados fornecidos pelo autor(a).

Essa ficha não pode ser modificada.



UNIVERSIDADE ESTADUAL PAULISTA

Câmpus de Jaboticabal



CERTIFICADO DE APROVAÇÃO

TÍTULO DA DISSERTAÇÃO: DIFERENÇAS SEXUAIS REFERENTES À PARTICIPAÇÃO DO CANAL CATIÔNICO ASIC1a NO *LOCUS COERULEUS* NA RESPOSTA COMPORTAMENTAL INDUZIDA POR CO₂


AUTORA: LETÍCIA RODRIGUES PINHEIRO


ORIENTADORA: LUCIANE HELENA GARGAGLIONI BATALHÃO

COORIENTADORA: ALANA TERCINO FRIAS

Aprovada como parte das exigências para obtenção do Título de Mestra em Ciência Animal, área: Fisiologia e Bem Estar Animal pela Comissão Examinadora:

Profa. Dra. LUCIANE HELENA GARGAGLIONI BATALHÃO (Participação Presencial)
Departamento de Morfologia e Fisiologia Animal / FCAV UNESP Jaboticabal


Prof. Dr. HÉLIO ZANGROSSI JUNIOR (Participação Virtual)
Departamento de Farmacologia / FMRP USP Ribeirão Preto


Prof. Dr. LUIZ GUILHERME DE SIQUEIRA BRANCO (Participação Virtual)
Departamento de Morfologia, Fisiologia e Patologia Básica / FORP USP Ribeirão Preto/SP

Jaboticabal, 29 de fevereiro de 2024



Documento assinado digitalmente
LUCIANE HELENA GARGAGLIONI
Data: 13/03/2024 09:57:42-0300
Verifique em <https://validar.it.gov.br>

DADOS CURRICULARES DO AUTOR

Letícia Rodrigues Pinheiro, nascida em 12 de novembro de 1996 na cidade de Tangará da Serra - MT, filha de Lídia Bezerra Rodrigues Pinheiro e Paulo César, ingressou no curso de Ciências Biológicas da Universidade Federal de Uberlândia (UFU), Campus Pontal em março de 2017. Durante o período de graduação, a aluna realizou uma iniciação científica sendo bolsista do Programa de Educação Tutorial-PET BIO. O estágio curricular foi realizado no Laboratório de Fisiologia da Universidade Estadual Paulista “Júlio de Mesquita Filho” Campus Jaboticabal. Colou grau em outubro de 2021 e recebeu o título de Bacharel em Ciências Biológicas. Em janeiro de 2022 iniciou o curso de Mestrado pelo programa de Pós-graduação em Ciência Animal no Departamento de Morfologia e Fisiologia Animal na Universidade Estadual Paulista- UNESP Campus Jaboticabal, onde foi bolsista CAPES.

“Quem quer algo de grande, como diz Goethe, deve saber limitar-se. Quem, pelo contrário, tudo quer, nada, em verdade, quer e nada consegue.” Georg Wilhelm Friedrich Hegel.

Dedico à minha mãe Lída, meu pai Paulo César (*in memoriam*) e aos meus tios, a quem amo profundamente.

AGRADECIMENTOS

Agradeço inicialmente à força que tem me guiado na forma de fé e esperança ao início de cada dia. Agradeço a mim, por todas as vezes que desistir não foi uma opção, apesar de todas as dificuldades encontradas nestes dois últimos anos.

À minha família, que sempre me apoiou e incentivou a minha trajetória acadêmica, em especial minha mãe Lídia e meus tios Daniel, Gerusa e Miriam. Vocês foram e são minha base e referencial de bondade, de força e de transformação.

Agradeço à minha orientadora Profa. Dra. Luciane Gargaglioni, a qual nutro imensa e profunda admiração, tanto no quesito profissional quanto pessoal, pela capacidade de nos inspirar e saber demandar nosso melhor. Obrigada por confiar em mim este trabalho e por ter me recebido de braços abertos no laboratório, sempre serei grata.

Agradeço à minha coorientadora e amiga Alana, por toda a paciência e disponibilidade em ensinar e sanar as dúvidas, e principalmente por me acalmar quando tudo parecia desmoronar, você foi e é imprescindível para a realização deste trabalho.

Aos meus amigos, que me acompanharam e me auxiliaram em todos os momentos, independente da distância ou da dificuldade, em especial minhas queridas amigas Amanda e Maria Emanuelle. Agradeço aos amigos que Jaboticabal me presenteou, a Paloma e Felipe, sem vocês a vida não teria o mesmo brilho. Aos meus amigos e colegas de laboratório, Ane, Ângela, Beatriz, Bianca, Camila, Geni, Gleyce, Isa, João, Lívia, Luana, Mariana, Sofia e Thaís, por todos os momentos que compartilhamos.

Agradeço ao auxílio do Dr. Luís Gustavo em diversas etapas deste estudo.

À Profa. Kênia por toda a contribuição no decorrer desse trabalho e em meu amadurecimento profissional.

Ao técnico Euclides pelo cuidado com os animais e a Ângela por todo apoio e carinho presente no cotidiano do laboratório, possibilitando que pudéssemos realizar as atividades da melhor forma.

Aos animais que cederam a vida para a realização deste estudo, ofereço todo o respeito e gratidão.

O presente trabalho foi realizado com apoio da Coordenação de Aperfeiçoamento de Pessoal de Nível Superior - Brasil (CAPES) - Código de Financiamento 001.

SUMÁRIO

Página

Resumo.....	01
Abstract	02
CAPÍTULO 1- CONSIDERAÇÕES GERAIS	03
Introdução	03
Objetivos	05
Revisão da Literatura.....	05
CAPÍTULO 2- Sexual differences regarding the participation of cationic channel ASIC1a within the <i>Locus coeruleus</i> in behavioral responses induced by CO ₂	25
Introduction	26
Methods	27
Results	31
Discussion.....	33
References.....	36
Figure Legends.....	41
Figures.....	42


CEUA – COMISSÃO DE ÉTICA NO USO DE ANIMAIS

CERTIFICADO

Certificamos que o projeto de pesquisa intitulado "Participação do canal cationico ASIC1a nas alterações cardiorrespiratórias e comportamentais induzidas por CO₂", protocolo nº 3451/22, sob a responsabilidade da Profª Drª Luciane Helena Gargaglioni Batalhão, que envolve a produção, manutenção e/ou utilização de animais pertencentes ao Filo Chordata, subfilo Vertebrata (exceto o homem), para fins de pesquisa científica (ou ensino) - encontra-se de acordo com os preceitos da lei nº 11.794, de 08 de outubro de 2008, no decreto 6.899, de 15 de julho de 2009, e com as normas editadas pelo Conselho Nacional de Controle de Experimentação Animal (CONCEA), e foi aprovado pela COMISSÃO DE ÉTICA NO USO DE ANIMAIS (CEUA), da FACULDADE DE CIÊNCIAS AGRÁRIAS E VETERINÁRIAS, UNESP - CÂMPUS DE JABOTICABAL-SP, em reunião ordinária de 15 de junho de 2022.

Vigência do Projeto	01/07/2022 a 03/02/2024
Espécie / Linhagem	C57BL/6
Nº de animais	64
Peso / Idade	25-30g/9 semanas
Sexo	Machos e Fêmeas
Origem	Biotério de Roedores do Departamento de Morfologia e Fisiologia Animal – FCAV/UNESP

Jaboticabal, 15 de junho de 2022.


Profª Drª Fabiana Pilarski
Coordenadora – CEUA

DIFERENÇAS SEXUAIS REFERENTES À PARTICIPAÇÃO DO CANAL CATIÔNICO ASIC1a NO LOCUS COERULEUS NA RESPOSTA COMPORTAMENTAL INDUZIDA POR CO₂

Resumo: O transtorno de pânico (TP) é uma condição marcada por intensa angústia e ansiedade, na qual o paciente desenvolve rapidamente sintomas respiratórios como falta de ar, hiperventilação e sensação de sufocamento. Apesar dos mecanismos dessa desordem permanecerem desconhecidos, estudos recentes sugerem provável conexão entre os desafios respiratórios e o TP, uma vez que episódios de hipóxia ou hipercapnia podem desencadear transtornos relacionados à ansiedade. O locus coeruleus (LC) é uma região quimiossensível capaz de gerar respostas emocionais efíacas durante episódios de estresse. Além disso, o LC possui neurônios noradrenérgicos que participam na resposta ventilatória ao CO₂, e a inalação de ar enriquecido com CO₂ pode desencadear ataques de pânico. Estudos recentes também sugerem a participação do canal iônico sensível a ácido do tipo 1a (ASIC1a) localizados na amígdala na resposta panicogênica induzidas por CO₂. Apesar disso, ainda não se conhece totalmente o envolvimento desse canal no LC na resposta ventilatória e às alterações comportamentais induzidas por concentrações elevadas de CO₂, que é um modelo utilizado para indução de ataques de pânico. O presente estudo teve como objetivo avaliar a participação dos canais ASIC1a do LC nas respostas respiratórias, metabólicas e comportamentais à exposição a 20% de CO₂ em camundongos machos e fêmeas após injeção intra-LC de Psalmotoxina-1 (Pstx-1, antagonista seletivo de canais ASIC1a). Os experimentos foram realizados em camundongos C57BL/6 machos e fêmeas, que foram submetidos à cirurgia estereotáxica para implante de cânulas-guia direcionadas ao LC e sensor de temperatura na cavidade abdominal. Sete dias depois, a dose de 50ng/0,1uL de Pstx1 ou salina foi administrada intra-LC e os animais foram expostos ao desafio respiratório. Parâmetros ventilatórios e metabólicos foram medidos por pletismografia de corpo inteiro em sistema fechado. A análise comportamental foi gravada por uma câmera e as respostas de fuga foram analisadas seguindo os parâmetros: número de saltos, *freezing*, corrida e levantamento de patas. Os resultados encontrados em camundongos machos e fêmeas indicam que os canais ASIC1a não parecem estar envolvidos no controle da respiração, mas participam das respostas comportamentais. Em machos, o antagonista promoveu uma diminuição no número de saltos (Salina: $9,33 \pm 5,85$ vs Pstx-1: $3,40 \pm 3,85$), já em fêmeas, não foi observado resultado significativo referente ao número de saltos nos animais que receberam Pstx-1 intra-LC (Salina: 1 ± 1 vs Pstx-1: $9 \pm 4,51$). Nossos dados sugerem que os canais ASIC1a no LC não participam do controle respiratório sob desafio de CO₂, mas estão envolvidos no comportamento de pânico induzido por CO₂, demonstrando uma resposta sexo específica.

Palavras-chave: Locus coeruleus, diferenças sexuais, ventilação, comportamento, ASIC1a

SEXUAL DIFFERENCES REGARDING THE PARTICIPATION OF CATIONIC CHANNEL ASIC1a WITHIN THE *LOCUS COERULEUS* IN BEHAVIORAL RESPONSES INDUCED BY CO₂

ABSTRACT: Panic disorder (PD) is a condition marked by intense anxiety, in which the patient quickly develops respiratory symptoms such as dyspnea, hyperventilation and a feeling of suffocation. Although the mechanisms of this disorder remain unknown, recent studies suggest a possible connection between respiratory challenges and PD, since episodes of hypoxia or hypercapnia can trigger anxiety-related disorders. The locus coeruleus (LC) is a chemosensitive region capable of generating emotional and physical responses during stress episodes. This region has noradrenergic neurons that participate in the ventilatory response to CO₂. The acid-sensitive ion channel type 1a (ASIC1a) has been suggested to participate in the panicogenic response induced by CO₂. Despite this, it is not yet fully known the involvement of this channel in the LC in the ventilatory response and behavioral changes induced by hypercapnia. We studied the effects of injection of a ASIC1a antagonist [Psalmostoxin -1 (Pstx-1- 50ng/0.1uL)] into the LC of C57BL/6 male and female mice on respiratory, metabolic and behavioral responses to exposure to 20% CO₂. We observed that Pstx-1 did not change ventilation and metabolism under normocapnic and hypercapnic conditions in both male and female mice. As to CO₂-behavioral responses, intra-LC Pstx-1 injection decreased the number of jumps in males. Therefore, our data suggest that ASIC1a channels in LC do not participate in respiratory control under CO₂ challenge, but are involved in CO₂-induced panic behavior, demonstrating a specific sex-dependent response.

Keywords: Locus coeruleus, sexual differences, ventilation, behavior, ASIC1a

CAPÍTULO 1- CONSIDERAÇÕES GERAIS

Introdução

O Transtorno do Pânico (TP) caracteriza-se por episódios de ataques de pânico súbitos e recorrentes, manifestando sintomas respiratórios, cardiovasculares, gastrointestinais, autonômicos e cognitivos de início agudo (Grant et al., 2006). No TP, o indivíduo vivencia ataques de pânico inesperados e recorrentes, seguidos de momentos de apreensão e/ou medo de sofrer novos ataques. Estes ataques se caracterizam por momentos de medo intenso e desconforto, que alcançam o ápice em poucos minutos, sendo acompanhados de sintomas físicos e/ou cognitivos (American Psychiatry Association, 2013).

Uma ferramenta diagnóstica relevante para o TP é o uso de agentes químicos, denominados de panicogênicos, (Kessler et al., 2006; Esquivel et al., 2010), que estimulam a respiração, sendo esta função fortemente regulada pelo pH. A inalação de CO₂ eleva a PCO₂ e reduz o pH sanguíneo, resultando em alterações que podem ser percebidas pelo encéfalo. Algumas regiões encefálicas que são sensíveis ao H⁺ e ao CO₂ estão relacionadas tanto ao controle da ventilação, quanto aos comportamentos defensivos, incluindo os relacionados com o pânico (Esquivel et al., 2010).

Dentre essas estruturas, o Locus coeruleus (LC) desempenha uma importante participação. O desequilíbrio na modulação do sistema LC/noradrenérgico está implicado em inúmeras desordens psiquiátricas, como transtorno de déficit de atenção e hiperatividade, doenças neurodegenerativas, transtorno de estresse pós- traumático, depressão e transtornos de ansiedade, incluindo o TP (Bangasser e Valentino, 2014; Fortress et al., 2015; Isingrini et al., 2016; Weinshenker, 2018). Em estudos clínicos mostraram, que uma regulação noradrenérgica alterada está presente em pacientes com TP e durante ataques de pânico (Charney et al., 1990; Bremner et al., 1996; Balaban, 2002; Dierssen et al., 2002). A elevada atividade noradrenérgica aumenta as respostas comportamentais relacionadas à ansiedade. Além disso, a ativação inadequada do LC pode participar da responsividade exagerada ao estímulo e do aumento da emotividade

observada em pacientes com estresse ou transtornos de ansiedade. Por outro lado, o tratamento antidepressivo, que é eficaz em pacientes com transtorno de pânico, diminui o disparo no LC e a expressão da tirosina hidroxilase, assim como as manipulações comportamentais que diminuem o estresse, como o manejo pós-natal, modulam as respostas do sistema noradrenérgico (Escorihuela et al., 1995; Baamonde et al., 2002).

Os mecanismos pelos quais as células do LC detectam as alterações causadas pelo CO₂ ainda não são completamente compreendidos. Dentre as possibilidades, os canais iônicos sensíveis a ácidos (ASIC1a) merecem destaque. Os ASIC1a são particularmente abundantes na amígdala e outras estruturas do circuito de medo, sugerindo uma contribuição para distúrbios comportamentais como ansiedade e depressão (Wemmie et al., 2003; Coryell et al., 2009). De fato, diversos comportamentos induzidos por inalação de CO₂, incluindo congelamento, aversão e evitação, dependem do ASIC1a (Ziemann et al., 2009). Esses canais são controlados por prótons e mediadores-chave das respostas à lesão neuronal. Eles exibem padrões únicos de distribuição no sistema nervoso central (SNC), com alta expressão em neurônios e baixa expressão em células gliais (Cegielski et al., 2022). As correntes provenientes do canal ASIC1a podem sofrer inibição por moduladores seletivos, como a toxina inibitória do nó de cisteína Psalmotoxina-1 (Pstx-1) (Escoubas et al., 2003; Dawson et al., 2012). A Pstx-1 inibe especificamente ASIC1a, não inibindo nenhum outro canal ASIC, ligando-se a esses canais de modo a alterar a sua conformação, fazendo com que sua afinidade ao H⁺ se altere (Escoubas et al., 2000; Chen et al., 2006).

É bastante conhecido que o sexo do indivíduo pode afetar o controle da respiração e a suscetibilidade a algumas doenças respiratórias (Gargaglioni et al., 2019; Holley et al., 2012; Jiang et al., 2017). Alguns estudos mostraram que há um dimorfismo sexual em humanos em anormalidades do sistema respiratório, como apnéias repetidas, hipopneias e oscilações respiratórias, que têm uma incidência três vezes maior em homens (Behan e Kinkead, 2011; Gargaglioni et al., 2019). Além disso, a síndrome da morte súbita em neonatos (SIDS) também é mais prevalente em indivíduos do sexo masculino na proporção de 60:4 (Moon et al., 2007), assim como a apneia obstrutiva do sono afeta predominantemente os homens (Wenninger et al., 2009). Por outro lado, as mulheres têm probabilidade

duas a três vezes maior de desenvolver transtorno do pânico do que os homens durante a vida (Kessler et al., 2005; Sheikh et al., 2002; Yonkers et al., 1998).

Apesar das evidências apresentadas, pouco se sabe sobre o envolvimento dos canais ASIC1a expressos no LC nas respostas respiratórias e comportamentais relacionadas ao TP, em animais machos e fêmeas, uma vez que diversos distúrbios podem ser prevalentes em um determinado sexo.

Objetivos

Os objetivos deste estudo foram avaliar a participação dos canais ASIC1a do LC nas respostas respiratórias, metabólicas e comportamentais à exposição a 20% CO₂ em camundongos machos e fêmeas após a microinjeção intra-LC de Pstx-1.

Revisão de literatura

Transtorno do pânico

O transtorno do pânico (TP) é caracterizado como um transtorno de ansiedade, sendo um dos transtornos psiquiátricos mais comuns, com prevalência média ao longo da vida de 4,5% da população (Kim et al., 2012). O TP é conhecido pela recorrência de ataques de pânico (APs) inesperados, que consistem em episódios de medo intenso acompanhados de alterações autonômicas e respiratórias como taquicardia, sudorese e falta de ar (Grant et al., 2006; American Psychiatry Association, 2013).

De acordo com o DSM-V (American Psychiatry Association, 2013), os APs recorrentes no TP são categorizados como espontâneos (inesperados) ou sinalizados (esperados), e são definidos como um período de intensa angústia ou desconforto caracterizado por pelo menos quatro dos seguintes sintomas típicos: palpitações, tremores, sudorese, asfixia, aperto no peito, dispneia, náusea, tontura, parestesia ou dormência, calafrios ou ondas de calor, descaracterização, enlouquecimento e sensação de morte (Guan e Cao, 2023). A recorrência de APs leva à ansiedade antecipatória, ao medo condicionado e à evitação de contextos de pânico, que levam ao comprometimento da vida cotidiana e à incapacidade (Klein, 1993; Smoller e Tsuang, 1998; Craske et al., 2010), sendo sua maior incidência em mulheres do que em homens (Battelan et al., 2007; Carta et al., 2015).

Diversos estudos buscando compreender o comportamento defensivo associados à estratégia de sobrevivência mediante a inúmeras ameaças, principalmente em roedores, têm sido úteis para entender as bases fisiopatológicas dos transtornos de ansiedade, em especial o TP (Guan e Cao, 2023). Desta maneira, é feita a avaliação referente às ameaças ambientais e a partir disso, é gerado respostas apropriadas para mitigar ou evitar danos. Durante este processo, diversas emoções podem ser desencadeadas, entre elas a ansiedade, medo e pânico (Livermore et al., 2021).

Estudos buscaram investigar o circuito neuronal que sustenta o TP para fornecer informações sobre as origens dos estados de ansiedade patológica. Dentre as hipóteses estudadas, tem sido proposto que os APs surgem quando o sensor de sufocamento do encéfalo detecta erroneamente a falta de ar, fazendo com que um sistema evoluído de alarme de sufocamento seja ativado. Um indivíduo com funcionamento incorreto está sujeito a “falsos alarmes de sufocamento” (Klein, 1993; Nardi, 2022). Os desafios respiratórios têm sido úteis no desenvolvimento de teorias sobre o TP, indicando que a sensibilidade ao dióxido de carbono (CO₂) seria um componente dos APs (Nardi, 2022).

Como metabólito do corpo humano, o CO₂ desempenha um papel insubstituível nos processos fisiológicos. A inalação de concentrações elevadas de CO₂, a hiperventilação voluntária e outros métodos para desencadear distúrbios ácido-base, como a infusão de lactato de sódio, podem induzir sintomas panicogênicos semelhantes em alguns pacientes com TP (Nardi et al., 2006; 2007; 2008). Diversos estudos demonstram que desafios respiratórios, como a exposição a uma alta concentração de CO₂ ou a uma baixa concentração de O₂, provocam ataques de pânico (Gorman et al., 1993; Beck et al., 1999; Perna et al., 2004; Fernandes et al., 2019), pois estimulam a respiração, sendo esta função fortemente regulada pelo pH (Esquivel et al., 2010). A inalação de CO₂ eleva a pressão parcial de CO₂ (PaCO₂) e reduz o pH sanguíneo, resultando em alterações no encéfalo. Algumas regiões encefálicas que são sensíveis ao H⁺ e ao CO₂ estão sendo relacionadas tanto ao controle da ventilação, quanto ao desencadeamento de comportamentos defensivos e respostas autonômicas, relacionadas ao pânico (Esquivel et al., 2010).

Ainda, pode-se evidenciar que a probabilidade de experimentar os efeitos panicogênicos do CO₂ são intensificados em pacientes com TP do que em indivíduos que não são acometidos por este transtorno (Coryell et al., 2006; Esquivel et al., 2010; Vollmer et al., 2015). Em camundongos, a inalação de CO₂ provoca comportamento de fuga, evitação e congelamento (Ziemann et al., 2009; D'Amato et al., 2011; Leibold et al., 2016; Vollmer et al., 2015; Spiacci et al., 2018; Taugher et al., 2020) e mudanças endócrinas, que sugerem que os animais estão passando por uma situação altamente aversiva, que se assemelha ao desencadeamento de um AP (Spiacci et al., 2018).

Quimiorrecepção

Uma das funções mais importantes e essenciais para os seres vivos, é a respiração. No contexto fisiológico, a respiração diz respeito à difusão e transporte de metabólitos entre o organismo e o ambiente externo. As células necessitam constantemente de oxigênio (O₂) para executar a respiração celular, de modo a gerar energia para o corpo realizar suas funções necessárias. A respiração em mamíferos é contínua, cuja função é adequar a ventilação alveolar dos pulmões para manter dentro de uma faixa constante a pressão parcial de oxigênio (Pa O₂) e PaCO₂, bem como os valores de pH. Apesar de aparentar simplicidade, a respiração é composta por um conjunto de elementos que a tornam de grande complexidade (Del Negro et al., 2018). Dessa maneira, as trocas gasosas são equilibradas em função da demanda do metabolismo (Gilmour, 2001).




Figura 1: Principais estruturas envolvidas na respiração (Criado com BioRender®).

Uma rede neural no tronco encefálico gera o ritmo respiratório e regula o padrão motor de cada respiração para atender às demandas metabólicas, biomecânicas e comportamentais dos indivíduos. No sentido rostro-caudal, os principais núcleos são o Complexo Bötzing (BötC), pré-BötC, grupo respiratório ventral rostral (rVRG) e caudal (cVRG), os quais formam a coluna respiratória ventral. Outros núcleos como o complexo pós-inspiratório (PiCo) e o parafacial (pF) encontram-se adjacentes à coluna respiratória ventral e participam do processo respiratório, sendo importantes para geração da fase de pós-inspiração e expiração ativa (Del Negro et al., 2018). O ritmo respiratório é gerado no pré-BötC (Smith et al., 1991). A respiração é dependente de aferências que chegam à essa região referentes às informações sobre as pressões parciais dos gases sanguíneos, o pH dos líquidos corporais, os gases inspirados, o estado de estiramento pulmonar e

informações detectadas por quimiorreceptores e mecanorreceptores. Buscando o pleno funcionamento do organismo, o sistema respiratório necessita estar apto para a geração de um ritmo que permita as trocas gasosas e integre os outros comportamentos relacionados à respiração. O sistema respiratório desempenha grande participação em atividades que requerem um fluxo de ar, como vocalização, deglutição, reflexos das vias aéreas como tossir, espirrar e até mesmo a locomoção (Greer et al., 2006; Klein e Codd, 2010).

Além de garantir a ritmogênese, processo pelo qual há a garantia do ritmo respiratório, outra grande função do sistema de controle respiratório é a quimiorrecepção. Os quimiorreceptores sensíveis a alterações de CO₂/pH são fundamentais para o controle da respiração, pelo fato de auxiliarem na geração do ritmo e na modulação do padrão respiratório, visando prevenir grandes variações das concentrações de CO₂ e pH, uma vez que estas alterações podem ocasionar em danos irreversíveis ao indivíduo. Os quimiorreceptores sensíveis ao CO₂/pH são classificados como periféricos (localizados nos corpos carotídeos e no arco aórtico) e centrais (localizados no SNC), sendo os centrais com predominância sobre a resposta ventilatória ao CO₂ (Blain et al., 2010; Foster e Smith, 2010; Patrone, 2018). Segundo Guyenet et al. (2010), os quimiorreceptores centrais são células sensíveis a variações na pCO₂ e no pH encefálico e contribuem para a estimulação da ventilação causada por hipercapnia ou acidose metabólica. Estes receptores são banhados pelo líquido cefalorraquidiano e isolados do sistema circulatório pela barreira hematoencefálica, que é pouco permeável ao íon H⁺, protegendo o encéfalo de variações no pH sistêmico, porém, altamente permeável ao CO₂ (Hlastala e Berger, 2001).

Os neurônios intrinsecamente sensíveis ao CO₂/pH estão localizados em muitas estruturas no encéfalo (Biancardi et al., 2008; Mulkey et al., 2004; Putnam et al., 2004; Ziemann et al., 2009). Algumas dessas estruturas provavelmente desempenham funções diferentes, como respiração e comportamento defensivo (Esquivel et al., 2009). A contribuição destas estruturas encefálicas na sensibilidade ao CO₂, detectando mudanças no pH e provocando uma resposta comportamental relacionada ao medo e ao pânico, ainda é indefinida.

O locus coeruleus (LC) é o maior conjunto de neurônios noradrenérgicos do

SNC, formando um par de núcleos adjacentes ao quarto ventrículo, localizados no tegmento pontino dorsolateral superior. Estima-se que metade de todas as projeções noradrenérgicas no SNC originam-se no LC (Aston-Jones et al., 1985; Berridge e Waterhouse, 2003). É uma estrutura importante para a quimiorrecepção central, possuindo mais de 80% dos neurônios altamente sensíveis às variações de CO₂/pH (Pineda e Aghajanian, 1997; Oyamada et al., 1998; Hilaire et al., 2004; Biancardi et al., 2008; Gargaglioni et al., 2010). A acidificação local do LC por meio de ácidos ou CO₂ resulta em aumentos da atividade do nervo frênico, demonstrando sua participação no controle ventilatório (Elam et al., 1981; Coates et al., 1993). De acordo com alguns estudos, a atividade dos neurônios do LC está relacionada à atividade respiratória (Oyamada et al., 1998; Andrzejewski et al., 2001), e alterações do sistema noradrenérgico na fase perinatal afetam a atividade dos neurônios respiratórios do tronco encefálico responsáveis pela geração do ritmo respiratório. O LC está relacionado a um grande papel nos processos fisiológicos e comportamentais, entre eles o ciclo sono-vigília, alimentação, nocicepção, termorregulação e aprendizagem (Viemari et al., 2004; Hilaire, 2004).

Ele também está implicado em diversas condições fisiopatológicas, como a síndrome de Rett (RTT) e os transtornos de ansiedade/pânico, apresentando sua atividade aumentada na presença de tais distúrbios. Adicionalmente, a quimiorrecepção central do LC ao CO₂ é um fator que torna essa região de alta relevância no estudo sobre o TP, uma vez que a inalação de ar enriquecido com CO₂ é um dos modelos estabelecidos para indução de ataques de pânico (Klein, 1993; Hasler et al., 2005; Goodwin et al., 2010).

Canais ASICs

A manutenção da homeostase do pH é essencial para as células de indivíduos como os mamíferos, pois os íons de hidrogênio desempenham papéis importantes na regulação do metabolismo (Serova et al., 2020). O pH extracelular do encéfalo é um parâmetro essencial que regula muitas variáveis e entre elas comportamento e a respiração. Em um encéfalo *in vivo* o pH intersticial geralmente varia de 7,1 a 7,25, mas esse equilíbrio pode ser comprometido por inúmeras condições, incluindo acidose metabólica, inflamação e hipoventilação, nas quais o pH pode cair para até 6,5 (Kintner et al., 2000).

Os canais iônicos sensíveis a ácidos (ASICs) desempenham um papel crítico na percepção de alterações de pH em condições relacionadas à acidose tecidual. Estes canais catiônicos controlados por prótons, insensíveis à voltagem, são ativados pela diminuição do pH extracelular e estão expressos em todo o SNC e periférico (Figura 2).




Figura 2: Mudanças conformacionais induzidas por prótons nos canais ASICs
(Fonte: Rook et al., 2020).

Os ASICs pertencem à superfamília de canais iônicos ENaC (Canal epitelial para Na^+ Epitelial)/DEG (Degenerina) e existem 6 subunidades identificadas até o presente momento 1a, 1b, 2a, 2b, 3 e 4, codificadas por 4 genes (ACCN1-4) (Waldmann et al., 1997; Gründer et al., 2000; Ortega-Ramírez et al., 2017). Essas subunidades possuem diferentes sensibilidades de pH, que possibilitam detectar uma ampla gama de pH fisiológico. Os membros da superfamília ASIC compartilham a mesma topologia, consistindo em dois domínios transmembranas hidrofóbicos, uma grande alça extracelular rica em cisteína e terminais N e C intracelulares curtos (Kweon & Suh, 2013). O ASIC1a é um complexo proteico e pode ser ativado em um ambiente extracelular ácido (de la Rosa et al., 2003), o que por sua vez leva a um grande influxo de sódio (Na^+) e cálcio (Ca^{2+}), promovendo a liberação de neurotransmissores relacionados (Xiong et al., 2004; Yermolaieva et al., 2004; Zha et al., 2006). Pesquisas recentes

mostraram que o ASIC1a desempenha um papel importante em certas condições patológicas, como inflamação, isquemia, tumor e outras doenças com dano tecidual provindo de acidificação (Bychkov et al., 2020; Gao et al., 2019; Wang et al., 2019; Xiong et al., 2004).

Diferentes subtipos de ASIC têm diferentes sensibilidades à acidose extracelular, na qual ASIC1a e ASIC3 são os subtipos mais sensíveis (Sutherland et al., 2001), e a dessensibilização de ASIC1a ocorre mais rapidamente (Waldmann et al., 1997). Estes canais geralmente são impermeáveis ao Ca^{2+} , no entanto, o ASIC1a homomérico é o único membro da família relatado como permeável ao Ca^{2+} (Yermolaieva et al., 2004). O aumento na concentração intracelular de Ca^{2+} é crucial para muitos processos fisiológicos e patológicos dos neurônios, como sinalização sináptica e metabolismo energético (Brini et al., 2014). A condutância de Ca^{2+} do ASIC1a homomérico revela as funções específicas que este complexo desempenha em processos fisiológicos e patológicos relacionados ao Ca^{2+} . Os canais ASIC1a são amplamente encontrados no hipocampo, amígdala, córtex cingulado, córtex somatossensorial, estriado e LC (Wemmie et al., 2003; Coryell et al., 2009; Prince et al., 2014). Nos neurônios, o ASIC1a está localizado na região do soma e nas espinhas dendríticas (Zha et al., 2006), sendo expresso principalmente na membrana celular e no citoplasma. No entanto, nos astrócitos, o ASIC1a é distribuído principalmente no núcleo (Huang et al., 2010) e na superfície das células gliais do hipocampo NG2 (Lin et al., 2010).

Os canais ASIC1a estão intimamente relacionados à plasticidade sináptica em diferentes regiões do encéfalo, ou seja, refere-se à eficácia da transmissão sináptica em resposta à atividade, que inclui duas formas principais: potenciação de longo prazo (LTP) e depressão de longo prazo (LTD) (Mango e Nisticò, 2020). No estriado, o ASIC1a está envolvido na regulação da plasticidade sináptica dos neurônios espinhosos médios, o que é crítico para a memória dependente dessa estrutura e o aprendizado processual (Yu et al., 2018). No circuito hipocampo-pré-frontal, o ASIC1a contribui para a plasticidade cortical relacionada à dor (Wang et al., 2018). No córtex cingulado anterior, o ASIC1a participa na modulação da hipersensibilidade à dor, mediando criticamente a potenciação sináptica (Li et al., 2019).

Uma ferramenta relevante para o estudos dos ASIC1a é a Psalmotoxina-1 (Pstx-1), um polipeptídeo extraído do veneno da toxina da tarântula *Psalmopoeus cambridgei*, que inibe especificamente os ASIC1a homoméricos (Escoubas et al., 2000), não inibindo nenhum outro ASIC, mesmo aqueles contendo a subunidade ASIC1a na forma heterodímera. A Pstx-1 liga-se a esses canais de modo a alterar a sua conformação, fazendo com que sua afinidade ao H^+ se altere (Escoubas et al., 2000; Chen et al., 2006).

Diferenças sexuais

Na história da ciência, a maior parte de todo o conhecimento disponível em fisiologia é centrado nas pesquisas em machos, consequentemente, muitos dados disponíveis são voltados aos machos e há uma tendência em se ignorar as fêmeas (Gargaglioni et al., 2019). É possível observar em diversas estruturas encefálicas diferenças morfológicas entre os sexos, como o LC (Luque et al., 1992; Pendergast et al., 2008; Thanky et al., 2002), área tegmental ventral (Brown et al., 2015), área pré-óptica (Gorski et al., 1980; Raisman e Field, 1971), substância negra (Ma et al., 2007), neocórtex, hipocampo e corpo caloso (Juraska, 1991), sendo necessário investigar as respostas provindas dessa diferenciação.

O sexo pode afetar o controle respiratório e a suscetibilidade a algumas doenças respiratórias (Block et al., 1979; Holley et al., 2012; Jiang et al., 2017; Lin e Winer, 2008; Wenninger et al., 2017). Estudos demonstram que as mulheres apresentam melhor recuperação respiratória do que os homens após um episódio grave de hipóxia (Garcia III et al., 2013). Ainda, sabe-se que existe um dimorfismo sexual em humanos relacionados a anormalidades do sistema respiratório, como apneias, hipopneias e oscilações respiratórias repetidas, no qual a incidência é 3 vezes maior em homens do que em mulheres (Behan et al., 2011).

Diferenças entre machos e fêmeas, podem ser encontradas também nas propriedades morfofisiológicas do LC. A partir de estudos realizados, foi possível evidenciar que em ratas adultas tem o núcleo LC maior que os machos (Busch et al., 1997; Garcia-Falgueras et al., 2005; Guillamón et al., 1988; Hormigo et al., 2015; Luque et al., 1992; Ohm et al., 1997; Pinos et al., 2001). Em mulheres, a região rostraldeste núcleo apresenta maior número de neurônios, enquanto nos homens

essa prevalência ocorre na região mais caudal do núcleo (Hormigo et al., 2015). Essa diferença estrutural sexo-dependente compreende um número maior de células noradrenérgicas e dendritos do LC mais longos e mais complexos em ratas adultas do que nos machos, mais especificamente na região peri-LC (Bangasser et al., 2016, Bangasser et al., 2011; Luque et al., 1992; Pinos et al., 2001). Com maior densidade dendrítica, foram encontrados mais contatos sinápticos no peri-LC ventromedial de fêmeas (Guillamón et al., 1988; Luque et al., 1992; Bangasser et al., 2011), e a capacidade de sintetizar e liberar noradrenalina pelos neurônios LC é maior nas fêmeas do que nos machos (Bangasser et al., 2019).

Portanto, em vista das evidências apresentadas, nossa hipótese é que os canais ASIC1a localizados no LC podem participam das respostas respiratórias, metabólicas e comportamentais à altas concentrações de CO₂ e essas resposta diferem de acordo com o sexo.

Referência Bibliográfica

AMERICAN PSYCHIATRIC ASSOCIATION. DSM 5. 2013. Washington, DC: American Psychiatric Association. 2013.

Andrzejewski M, Mückenhoff K, Scheid P, Ballantyne D (2001) Synchronized rhythms in chemosensitive neurons of the locus coeruleus in the absence of chemical synaptic transmission. **Respiration physiology** 129:123-140.

Aston-Jones G, Foote SL, Segal M (1985) Impulse conduction properties of noradrenergic locus coeruleus axons projecting to monkey cerebrocortex. **Neuroscience** 15:765-777.

Baamonde C, Lumbreras MA, Martinez-Cue C, Vallina IF, Florez J, Dierssen M (2002) Postnatal handling induces long-term modifications in central β -noradrenergic signalling in rats. **Stress** 5:137-147.

Balaban CD (2002) Neural substrates linking balance control and anxiety. **Physiology & behavior** 77:469-475.

Bangasser DA, Eck SR, Ordoñez Sanchez E (2019) Sex differences in stress reactivity in arousal and attention systems. **Neuropsychopharmacology** 44:129-139.

Bangasser DA, Valentino RJ (2014) Sex differences in stress-related psychiatric disorders: neurobiological perspectives. **Frontiers in neuroendocrinology** 35:303-319.

Bangasser DA, Wiersielis KR, Khantsis S (2016) Sex differences in the locus coeruleus-norepinephrine system and its regulation by stress. *Brain research* 1641:177-188.

Bangasser DA, Zhang X, Garachh V, Hanhauser E, Valentino RJ (2011) Sexual dimorphism in locus coeruleus dendritic morphology: a structural basis for sex differences in emotional arousal. **Physiology & behavior** 103:342-351.

Batelaan N, De Graaf R, Van Balkom A, Vollebergh W, Beekman A (2007) Thresholds for health and thresholds for illness: panic disorder versus subthreshold panic disorder. **Psychological medicine** 37:247-256.

Behan M, Kinkead R (2011) Neuronal control of breathing: sex and stress hormones. **Comprehensive Physiology** 1:2101-2139.

Behan M, Zabka AG, Thomas CF, Mitchell GS (2003) Sex steroid hormones and the neural control of breathing. **Respiratory physiology & neurobiology** 136:249-263.

Beck JG, Ohtake PJ, Shipherd JC (1999) Exaggerated anxiety is not unique to CO₂ in panic disorder: A comparison of hypercapnic and hypoxic challenges. **Journal of Abnormal Psychology** 108:473.

Berridge CW, Waterhouse BD (2003) The locus coeruleus–noradrenergic system: modulation of behavioral state and state-dependent cognitive processes. **Brain research reviews** 42:33-84.

Biancardi V, Bícigo KC, Almeida MC, Gargaglioni LH (2008) Locus coeruleus noradrenergic neurons and CO₂ drive to breathing. *Pflügers Archiv-European Journal of Physiology*, 455, 1119-1128.

Blain GM, Smith CA, Henderson KS, Dempsey JA (2010) Peripheral chemoreceptors determine the respiratory sensitivity of central chemoreceptors to CO₂. **The Journal of physiology** 588:2455-2471.

Block AJ, Boysen PG, Wynne J W, Hunt LA (1979) Sleep apnea, hypopnea and oxygen desaturation in normal subjects: a strong male predominance. **New England Journal of Medicine** 300:513-517.

Bremner JD, Krystal JH, Southwick SM, Charney DS (1996) Noradrenergic mechanisms in stress and anxiety: I. Preclinical studies. **Synapse** 23:28-38.

Brini M, Calì T, Ottolini D, Carafoli E (2014) Neuronal calcium signaling: function and dysfunction. **Cellular and molecular life sciences** 71:2787-2814.

Brown RM, Kim AK, Khoo SYS, Kim JH, Jupp B, Lawrence AJ (2016) Orexin -1 receptors signalling in the prelimbic cortex and ventral tegmental area regulates cue-induced reinstatement of ethanol-seeking in iP rats. **Addiction biology** 21:603-612.

Busch C, Bohl J, Ohm T (1997) Spatial, temporal and numeric analysis of Alzheimer changes in the nucleus coeruleus. **Neurobiology of aging** 18:401-406.

Bychkov M, Shulepko M, Osmakov D, Andreev Y, Sudarikova A, Vasileva V, Lyukmanova E (2020) Mambalgin-2 induces cell cycle arrest and apoptosis in glioma cells via interaction with ASIC1a. **Cancers** 12:1837.

Carta MG, Moro MF, Aguglia E, Balestrieri M, Caraci F, Dell’Osso L, Faravelli C (2015) The attributable burden of panic disorder in the impairment of quality of life in a national survey in Italy. **International Journal of Social Psychiatry**, 61:693-699.

Cegielski V, Chakrabarty R, Ding S, Wacker MJ, Monaghan-Nichols P, Chu XP (2022) Acid-sensing ion channels in glial cells. **Membranes** 12:119.

Charney DS, Woods SW, Nagy LM, Southwick SM, Krystal JH, Heninger GR (1990) Noradrenergic function in panic disorder. **The Journal of clinical psychiatry** 51:5-11.

Chen X, Kalbacher H, Grunder S (2006) Interaction of acid-sensing ion channel

(ASIC)1 with the tarantula toxin psalmotoxin 1 is state dependent. **The Journal of general physiology** 127:267-276.

Coates EL, Li AIHUA, Nattie EE (1993) Widespread sites of brain stem ventilatory chemoreceptors. **Journal of Applied Physiology** 75:5-14.

Coryell MW, Wunsch AM, Haenfler JM, Allen JE, Schnizler M, Ziemann AE, Wemmie JA (2009) Acid-sensing ion channel-1a in the amygdala, a novel therapeutic target in depression-related behavior. **Journal of Neuroscience** 29:5381-5388.

Coryell W, Pine D, Fyer A, Klein D (2006) Anxiety responses to CO₂ inhalation in subjects at high-risk for panic disorder. **Journal of affective disorders** 92:63-70.

Craske MG, Kircanski K, Epstein A, Wittchen HU, Pine DS, Lewis-Fernández R, Hinton D (2010) Panic disorder: a review of DSM-IV panic disorder and proposals for DSM-V. **Depression and anxiety** 27:93-112.

D'Amato FR, Zanettini C, Lampis V, Coccurello R, Pascucci T, Ventura R, Battaglia M (2011) Unstable maternal environment, separation anxiety, and heightened CO₂ sensitivity induced by gene-by-environment interplay. **PLoS One** 6:e18637.

Dawson RJ, Benz J, Stohler P, Tetaz T, Joseph C, Huber S, Ruf A (2012) Structure of the acid-sensing ion channel 1 in complex with the gating modifier Psalmotoxin 1. **Nature communications** 3:936.

de la Rosa DA, Krueger SR, Kolar A, Shao D, Fitzsimonds RM, Canessa CM (2003) Distribution, subcellular localization and ontogeny of ASIC1 in the mammalian central nervous system. **The Journal of physiology** 546:77-87.

Del Negro CA, Funk GD, Feldman JL (2018) Breathing matters. **Nature Reviews Neuroscience** 19:351-367.

Dierssen M, Fotaki V, de Lagrán MM, Gratacos M, Arbones M, Fillat C, Estivill X (2002) Neurobehavioral development of two mouse lines commonly used in transgenic studies. **Pharmacology Biochemistry and Behavior** 73:19-25.

Elam M, Yao T, Thore P, Svensson TH (1981) Hypercapnia and hypoxia: chemoreceptor-mediated control of locus coeruleus neurons and splanchnic, sympathetic nerves. **Brain research** 222:373-381.

Escorihuela RM, Fernandezteruel A, Tobena A, Vivas NM, Marmol F, Badia A, Dierssen M (1995) Early environmental stimulation produces long-lasting changes on β -adrenoceptor transduction system. **Neurobiology of learning and memory**, 64:49- 57.

Escoubas P, De Weille JR, Lecoq A, Diochot S, Waldmann R, Champigny G, Lazdunski M (2000) Isolation of a tarantula toxin specific for a class of proton-gated Na⁺ channels. **Journal of Biological Chemistry** 275:25116-25121.

Escoubas P, Bernard C, Lambeau G, Lazdunski M, Darbon H (2003) Recombinant production and solution structure of PcTx1, the specific peptide inhibitor of ASIC1a proton-gated cation channels. **Protein Science** 12:1332-1343.

Esquivel G, Schruers KR, Maddock RJ, Colasanti A, Griez EJ (2010). Acids in the brain: a factor in panic?. **Journal of psychopharmacology** 24:639-647.

Fernandes GG, Frias AT, Spiacci Jr A, Pinheiro LC, Tanus-Santos JE, Zangrossi Jr H (2019) Nitric oxide in the dorsal periaqueductal gray mediates the panic-like escape response evoked by exposure to hypoxia. **Progress in Neuro-Psychopharmacology and Biological Psychiatry** 92:321-327.

Fortress AM, Hamlett ED, Vazey EM, Aston-Jones G, Cass WA, Boger HA, Granholm ACE (2015) Designer receptors enhance memory in a mouse model of Down syndrome. **Journal of Neuroscience** 35:1343-1353.

Forster HV, Smith CA (2010) Contributions of central and peripheral chemoreceptors to the ventilatory response to CO₂/H⁺. **Journal of Applied Physiology** 108:989-994.

Gao WF, Xu YY, Ge JF, Chen FH (2019) Inhibition of acid-sensing ion channel 1a attenuates acid-induced activation of autophagy via a calcium signaling pathway in articular chondrocytes. **International Journal of Molecular Medicine** 43:1778-1788

Garcia-Falgueras A, Pinos H, Collado P, Pasaro E, Fernandez R, Segovia S, Guillaumon A (2005) The expression of brain sexual dimorphism in artificial selection of rat strains. **Brain research** 1052:130-138.

Garcia III AJ, Rotem-Kohavi N, Doi A, Ramirez JM (2013) Post-hypoxic recovery of respiratory rhythm generation is gender dependent. **PloS one** 8:e60695.

Gargaglioni LH, Hartzler LK, Putnam RW (2010) The locus coeruleus and central chemosensitivity. **Respiratory physiology & neurobiology** 173:264-273.

Gargaglioni LH, Marques DA, Patrone LGA (2019) Sex differences in breathing. **Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology** 238:110543.

Gorman JM, Papp LA, Coplan J, Martinez J, Liebowitz MR, Klein DF (1993) The effect of acetazolamide on ventilation in panic disorder patients. **The American journal of psychiatry** 150:1480-1484.

Grant BF, Hasin DS, Stinson FS, Dawson DA, Goldstein RB, Smith S, Saha TD (2006) The epidemiology of DSM-IV panic disorder and agoraphobia in the United States: results from the National Epidemiologic Survey on Alcohol and Related Conditions. **The Journal of clinical psychiatry** 67: 6517.

Gilmour KM (2001) The CO₂/pH ventilatory drive in fish. **Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology** 130:219-240.

Goodwin RD, Pagura J, Cox B, Sareen J (2010) Asthma and mental disorders in Canada: impact on functional impairment and mental health service use. **Journal of psychosomatic research** 68:165-173.

Gorski RA, Harlan RE, Jacobson CD, Shryne JE, Southam AM (1980) Evidence for the existence of a sexually dimorphic nucleus in the preoptic area of the rat. **Journal of Comparative Neurology** 193:529-539.

Gründer S, Geissler HS, Bässler EL, Ruppertsberg JP (2000) A new member of acid-sensing ion channels from pituitary gland. **Neuroreport** 11:1607-1611.

Greer JJ, Funk GD, Ballanyi K (2006) Preparing for the first breath: prenatal maturation of respiratory neural control. **The Journal of physiology** 570:437-444.

Guan X, Cao, P (2023) Brain Mechanisms Underlying Panic Attack and Panic Disorder. **Neuroscience Bulletin** 1-20.

Guillamón A, de Blas MR, Segovia S (1988) Effects of sex steroids on the development of the locus coeruleus in the rat. **Developmental Brain Research** 40:306-310.

Guyenet PG, Bayliss DA (2015) Neural control of breathing and CO₂ homeostasis. **Neuron**, 87:946-961.

Guyenet PG, Mulkey DK (2010) Retrotrapezoid nucleus and parafacial respiratory group. **Respiratory physiology & neurobiology** 173:244-255.

Hasler G, Gergen PJ, Kleinbaum DG, Ajdacic V, Gamma A, Eich D, Angst J (2005) Asthma and panic in young adults: a 20-year prospective community study. **American journal of respiratory and critical care medicine** 171:1224-1230.

Hilaire G, Viemari JC, Coulon P, Simonneau M, Bévengut M (2004) Modulation of the respiratory rhythm generator by the pontine noradrenergic A5 and A6 groups in rodents. **Respiratory physiology & neurobiology** 143:187-197.

Hlastala MP, Berger AJ (2001) Physiology of respiration. Oxford University Press.

Holley HS, Behan M, Wenninger JM (2012) Age and sex differences in the ventilatory response to hypoxia and hypercapnia in awake neonatal, pre-pubertal and young adults. **Respiratory physiology & neurobiology** 180:79-87.

Hormigo S, Gómez-Nieto R, Castellano O, Herrero-Turrión MJ, López DE, de Anchieta de Castro e Horta-Júnior J (2015) The noradrenergic projection from the locus coeruleus to the cochlear root neurons in rats. **Brain Structure and Function** 220:1477-1496.

Huang C, Hu ZL, Wu WN, Yu DF, Xiong QJ, Song JR, Chen JG (2010) Existence and distinction of acid-evoked currents in rat astrocytes. **Glia** 58:1415-1424.

Isingrini E, Perret L, Rainer Q, Amilhon B, Guma E, Tanti A, Giros B (2016) Resilience to chronic stress is mediated by noradrenergic regulation of dopamine neurons. *Nature neuroscience*, 19(4), 560-563.

Jiang C, Cui N, Zhong W, Johnson CM, Wu Y (2017) Breathing abnormalities in animal models of Rett syndrome a female neurogenetic disorder. **Respiratory physiology & neurobiology** 245:45-52.

Juraska JM (1991) Sex differences in “cognitive” regions of the rat brain. **Psychoneuroendocrinology** 16:105-119.

Kessler RC, Chiu WT, Demler O, Walters EE (2005) Prevalence, severity, and comorbidity of 12-month DSM-IV disorders in the National Comorbidity Survey Replication. **Archives of general psychiatry** 62:617-627.

Kessler RC, Chiu WT, Jin R, Ruscio AM, Shear K, Walters EE (2006) The epidemiology of panic attacks, panic disorder, and agoraphobia in the National Comorbidity Survey Replication. **Arch Gen Psychiatry** 63:415– 24.

Kim JE, Dager SR, Lyoo IK (2012) The role of the amygdala in the pathophysiology of panic disorder: evidence from neuroimaging studies. **Biology of mood & anxiety disorders** 2:1-17.

Klein DF (1993) False suffocation alarms, spontaneous panics, and related conditions: an integrative hypothesis. **Archives of general psychiatry** 50:306-317.

Klein W, Codd JR (2010) Breathing and locomotion: Comparative anatomy, morphology and function. **Respiratory physiology & neurobiology** 173:S26-S32.

Kintner DB, Anderson MK, Fitzpatrick JH, Sailor KA, Gilboe DD (2000) ³¹P-MRS-based determination of brain intracellular and interstitial pH: its application to in vivo H⁺ compartmentation and cellular regulation during hypoxic/ischemic conditions. **Neurochemical research** 25:1385-1396.

Kweon HJ, Suh BC (2013) Acid-sensing ion channels (ASICs): therapeutic targets for neurological diseases and their regulation. **BMB reports** 46:295.

Leibold NK, Van Den Hove DLA, Viechtbauer W, Buchanan GF, Goossens L, Lange I, Schruers KRJ (2016) CO₂ exposure as translational cross-species experimental model for panic. **Translational psychiatry** 6:e885-e885.

Li HS, Su XY, Song XL, Qi X, Li Y, Wang RQ, Xu TL (2019) Protein kinase C lambda mediates acid-sensing ion channel 1a-dependent cortical synaptic plasticity and pain hypersensitivity. **Journal of Neuroscience** 39:5773-5793.

Lin NU, Winer EP (2008) Advances in adjuvant endocrine therapy for postmenopausal women. **Journal of Clinical Oncology** 26:798-805.

Lin YC, Liu YC, Huang YY, Lien CC (2010) High-density expression of Ca^{2+} -permeable ASIC1a channels in NG2 glia of rat hippocampus. **PloS one** 5:e12665.

Livermore JJ, Klaassen FH, Bramson B, Hulsman AM, Meijer SW, Held L, Roelofs K (2021) Approach-avoidance decisions under threat: the role of autonomic psychophysiological states. **Frontiers in Neuroscience** 15:621517.

Luque JM, de Blas MR, Segovia S, Guillamón A (1992) Sexual dimorphism of the dopamine- β -hydroxylase-immunoreactive neurons in the rat locus ceruleus. **Developmental brain research** 67:211-215.

Ma YY, Kong SZ, Yang LJ, Meng JL, Lv LC, He M (2007) Sexual dimorphisms of dopaminergic neurons in rat substantia nigra. **Sheng li xue bao:[Acta Physiologica Sinica]** 59:753-758.

Mango D, Nisticò R (2020) Role of ASIC1a in normal and pathological synaptic plasticity. **Reviews of physiology, biochemistry and pharmacology**:83-100.

Moon RY, Horne RS, Hauck FR (2007) Sudden infant death syndrome. **The Lancet** 370:1578-1587.

Mulvey B, Bhatti DL, Gyawali S, Lake AM, Kriaucionis S, Ford CP, Dougherty JD (2018) Molecular and functional sex differences of noradrenergic neurons in the mouse locus coeruleus. **Cell Reports** 23:2225-2235.

Nardi AE (2022) The amygdala might be the main target for Donald Klein's "Real" False Suffocation Alarm hypothesis for triggering panic attacks. **Biological Psychology** 108306-108306.

Nardi AE, Valença AM, Lopes FL, Nascimento I, Veras AB, Freire RC, Zin WA (2006) Psychopathological profile of 35% CO₂ challenge test-induced panic attacks: a comparison with spontaneous panic attacks. **Comprehensive psychiatry**, 47:209- 214.

Nardi AE, Valença AM, Lopes FL, Nascimento I, Veras AB, Freire RC, Zin WA (2006) Psychopathological profile of 35% CO₂ challenge test-induced panic attacks: a comparison with spontaneous panic attacks. **Comprehensive psychiatry** 47:209- 214.

Nardi AE, Valença AM, Lopes FL, de-Melo-Neto VL, Freire RC, Veras AB, Zin WA (2007) Caffeine and 35% carbon dioxide challenge tests in panic disorder. **Human Psychopharmacology: Clinical and Experimental** 22:231-240.

Nardi AE, Lopes FL, Valença AM, Freire RC, Veras AB, de-Melo-Neto VL, Zin WA (2007) Caffeine challenge test in panic disorder and depression with panic attacks. **Comprehensive psychiatry** 48:257-263.

Nardi AE, Valença AM, Nascimento I, Freire RC, Veras AB, de-Melo-Neto VL, Zin WA(2008) A caffeine challenge test in panic disorder patients, their healthy first-degree relatives, and healthy controls. **Depression and anxiety** 25:847-853.

Ohm TG, Busch C, Bohl J (1997) Unbiased estimation of neuronal numbers in the human nucleus coeruleus during aging. **Neurobiology of aging** 18:393-399.

Ortega-Ramírez A, Vega R, Soto E (2017) Acid-sensing ion channels as potential therapeutic targets in neurodegeneration and neuroinflammation. **Mediators of inflammation**.

Oyamada Y, Ballantyne D, Mückenhoff K, Scheid P (1998) Respiration -modulated membrane potential and chemosensitivity of locus coeruleus neurones in the in vitro brainstem-spinal cord of the neonatal rat. **The Journal of physiology** 513:381-398.

Patrone LGA, Biancardi V, Marques DA, Bícago KC, Gargaglioni LH (2018) Brainstem catecholaminergic neurones and breathing control during postnatal development in male and female rats. **The Journal of Physiology** 596:3299-3325.

Pendergast JS, Tuesta LM, Bethea JR (2008) Oestrogen receptor β contributes to the transient sex difference in tyrosine hydroxylase expression in the mouse locus coeruleus. **Journal of neuroendocrinology** 20:1155-1164.

Perna G, Bertani A, Politi E, Colombo G, Bellodi L (1997) Asthma and panic attacks. **Biological psychiatry** 42:625-630.

Pineda J, Aghajanian GK (1997) Carbon dioxide regulates the tonic activity of locus coeruleus neurons by modulating a proton-and polyamine-sensitive inward rectifier potassium current. **Neuroscience** 77:723-743.

Pinos H, Collado P, Rodríguez-Zafra M, Rodríguez C, Segovia S, Guillamón A (2001). The development of sex differences in the locus coeruleus of the rat. **Brain research bulletin** 56:73-78.

Price MP, Gong H, Parsons MG, Kundert JR, Reznikov LR, Bernardinelli L, Welsh M (2014) Localization and behaviors in null mice suggest that ASIC1 and ASIC2 modulate responses to aversive stimuli. **Genes, Brain and Behavior** 13:179-194.

Putnam RW, Filosa JA, Ritucci, NA (2004) Cellular mechanisms involved in CO₂ and acid signaling in chemosensitive neurons. **American Journal of Physiology-Cell Physiology** 287:C1493-C1526.

Raisman G, Field PM (1971) Sexual dimorphism in the preoptic area of the rat. **Science** 173:731-733.

Rook ML, Musgaard M, MacLean DM (2021) Coupling structure with function in acid- sensing ion channels: challenges in pursuit of proton sensors. **The Journal of**

physiology 599:417-430.

Serova OV, Gantsova EA, Deyev IE, Petrenko AG (2020) The value of pH sensors in maintaining homeostasis of the nervous system. **Russian Journal of Bioorganic Chemistry** 46:506-519.

Sheikh JI, Leskin GA, Klein DF (2002) Gender differences in panic disorder: findings from the National Comorbidity Survey. **American Journal of Psychiatry** 159:55-58.

Smoller JW, Tsuang MT (1998) Panic and phobic anxiety: defining phenotypes for genetic studies. **American Journal of Psychiatry** 155:1152-1162.

Smith JC, Ellenberger HH, Ballanyi K, Richter DW, Feldman JL (1991) Pre-Bötzinger complex: a brainstem region that may generate respiratory rhythm in mammals. **Science** 254:726-729.

Spiacci A, Vilela-Costa HH, Sant'Ana AB, Fernandes GG, Frias AT, da Silva GSF, Zangrossi Jr H (2018) Panic-like escape response elicited in mice by exposure to CO₂, but not hypoxia. **Progress in Neuro-Psychopharmacology and Biological Psychiatry** 81:178-186.

Sutherland SP, Benson CJ, Adelman JP, McCleskey EW (2001) Acid-sensing ion channel 3 matches the acid-gated current in cardiac ischemia-sensing neurons. **Proceedings of the National Academy of Sciences** 98:711-716.

Taughher RJ, Dlouhy BJ, Kreple CJ, Ghobbeh A, Conlon MM, Wang Y, Wemmie JA (2020) The amygdala differentially regulates defensive behaviors evoked by CO₂. **Behavioural brain research** 377:112236.

Thanky NR, Son JH, Herbison AE (2002) Sex differences in the regulation of tyrosine hydroxylase gene transcription by estrogen in the locus coeruleus of TH9-LacZ transgenic mice. **Molecular brain research** 104:220-226.

Viemari JC, Bevegut M, Burnet H, Coulon P, Pequignot JM, Tiveron MC, Hilaire G (2004) Phox2a gene, A6 neurons, and noradrenaline are essential for development of normal respiratory rhythm in mice. **Journal of Neuroscience** 24:928-937.

Vollmer LL, Ghosal S, McGuire JL, Ahlbrand RL, Li KY, Santin JM, Sah R (2016) Microglial acid sensing regulates carbon dioxide-evoked fear. **Biological psychiatry** 80:541-551.

Vollmer LL, Strawn JR, Sah R (2015) Acid-base dysregulation and chemosensory mechanisms in panic disorder: a translational update. **Translational psychiatry**, 5:e572-e572.

Xiong ZG, Zhu XM, Chu XP, Minami M, Hey J, Wei WL, Simon RP (2004) Neuroprotection in ischemia: blocking calcium-permeable acid-sensing ion channels. **Cell** 118:687-698.

Waldmann R, Champigny G, Bassilana F, Heurteaux C, Lazdunski M (1997) A proton-gated cation channel involved in acid-sensing. **Nature** 386:173-177.

Wang Q, Wang Q, Song XL, Jiang Q, Wu YJ, Li Y, Xu TL (2018) Fear extinction requires ASIC1a-dependent regulation of hippocampal-prefrontal correlates. **Science Advances** 4:eaau3075.

Wang Y, Sun Y, Zuo L, Wang Y, Huang Y (2019) ASIC1a promotes high glucose and PDGF-induced hepatic stellate cell activation by inducing autophagy through CaMKK β /ERK signaling pathway. **Toxicology Letters** 300:1-9.

Weinshenker D (2018) Long road to ruin: noradrenergic dysfunction in neurodegenerative disease. **Trends in neurosciences** 41:211-223.

Wemmie JA, Askwith CC, Lamani E, Cassell MD, Freeman JH, Welsh MJ (2003) Acid-sensing ion channel 1 is localized in brain regions with high synaptic density and contributes to fear conditioning. **Journal of Neuroscience** 23:5496-5502.

Wenninger JM, Olson Jr EB, Cotter CJ, Thomas CF, Behan M (2009) Hypoxic and hypercapnic ventilatory responses in aging male vs. aging female rats. **Journal of Applied Physiology** 106:1522-1528.

Yermolaieva O, Leonard AS, Schnizler MK, Abboud FM, Welsh MJ (2004) Extracellular acidosis increases neuronal cell calcium by activating acid-sensing ion channel 1a. **Proceedings of the National Academy of Sciences** 101:6752-6757.

Yonkers KA, Zlotnick C, Allsworth J, Warshaw M, Shea T, Keller MB (1998) Is the course of panic disorder the same in women and men?. **American Journal of Psychiatry** 155:596-602.

Yu Z, Wu YJ, Wang YZ, Liu DS, Song XL, Jiang Q, Xu TL (2018) The acid-sensing ion channel ASIC1a mediates striatal synapse remodeling and procedural motor learning. **Science Signaling** 11:eaar4481.

Zha XM, Wemmie JA, Green SH, Welsh MJ (2006) Acid-sensing ion channel 1a is a postsynaptic proton receptor that affects the density of dendritic spines. **Proceedings of the National Academy of Sciences** 103:16556-16561.

Ziemann AE, Allen JE, Dahdaleh NS, Drebot II, Coryell MW, Wunsch AM, Wemmie JA (2009) The amygdala is a chemosensor that detects carbon dioxide and acidosis to elicit fear behavior. **Cell** 139:1012-102.

CAPÍTULO 2- SEXUAL DIFFERENCES REGARDING THE PARTICIPATION OF AN IONIC CHANNEL ASIC1a WITHIN THE *LOCUS COERULEUS* IN BEHAVIORAL RESPONSES INDUCED BY CO₂

Abstract: Panic disorder (PD) is a condition marked by intense anxiety, in which the patient quickly develops respiratory symptoms such as dyspnea, hyperventilation and a feeling of suffocation. Although the mechanisms of this disorder remain unknown, recent studies suggest a possible connection between respiratory challenges and PD, since episodes of hypoxia or hypercapnia can trigger anxiety-related disorders. The locus coeruleus (LC) is a chemosensitive region capable of generating emotional and physical responses during stress episodes. This region has noradrenergic neurons that participate in the ventilatory response to CO₂. The acid-sensitive ion channel type 1a (ASIC1a) has been suggested to participate in the panicogenic response induced by CO₂. Despite this, it is not yet fully known the involvement of this channel in the LC in the ventilatory response and behavioral changes induced by hypercapnia. We studied the effects of injection of an ASIC1a antagonist [Psalmostoxin -1 (Pstx-1 - 50ng/0.1μL)] into the LC of C57BL/6 male and female mice on respiratory, metabolic and behavioral responses to exposure to 20% CO₂. We observed that Pstx-1 did not change ventilation and metabolism under normocapnic and hypercapnic conditions in both male and female mice. As to CO₂-behavioral responses, intra-LC Pstx-1 injection decreased the number of jumps in males, but there was no statistically significant difference in the behavioral parameters of the females. Therefore, our data suggest that ASIC1a channels in LC do not participate in respiratory control under room air and hypercapnia, but are involved in CO₂-induced panic behavior, demonstrating a specific sex-dependent response.

Keywords: Locus coeruleus, sexual differences, ventilation, behavior, ASIC1a

Introduction

Panic disorder (PD) is characterized by panic attacks, with behavioral changes and respiratory symptoms such as shortness of breath, hyperventilation and feeling of suffocation (Grant et al., 2006). According to the DSM-V (American Psychiatry Association, 2013), the recurrent PD are categorized as spontaneous (unexpected) or signaled (expected) and are defined as a period of intense distress or discomfort characterized by typical symptoms such as palpitations, tremors, sweating, asphyxia, chest tightness, dyspnea, nausea and feeling of death.

Some studies suggest a link between changes in chemosensitivity to CO₂/pH and PD, since episodes of hypoxia or hypercapnia can trigger panic attacks (Kessler et al., 2005; Esquivel et al., 2010). In fact, brain regions that are sensitive to H⁺ and CO₂ are related to both ventilation control and defensive behaviors, including those related to panic (Esquivel et al., 2010).

Among these structures, the Locus coeruleus (LC) plays an important role in breathing control and anxiety related behaviors (Gargaglioni et al., 2010; Borodovitsyna et al., 2018). Imbalance in the modulation of the LC/noradrenergic system is implicated in numerous psychiatric disorders, such as attention deficit hyperactivity disorder, neurodegenerative diseases, post-traumatic stress disorder, depression and anxiety disorders, including PD (Bangasser and Valentino, 2014; Fortress et al., 2015; Isingrini et al., 2016; Weinshenker, 2018). Clinical studies have shown that altered noradrenergic regulation is present in PD patients and during panic attacks (Charney et al., 1990; Bremner et al., 1996; Balaban, 2002; Dierssen et al., 2002).

The mechanisms by which LC cells detect changes caused by CO₂ are not yet fully understood (Putnam et al., 2004). Among the possibilities, acid-sensitive ion channels (ASIC1a) stand out. ASIC1a are particularly abundant in the amygdala and other structures of the fear circuit, suggesting a contribution to behavioral disorders such as anxiety and depression (Taughner et al., 2017; Zhu et al., 2022). ASIC1a channels respond to decreasing pH, which has been identified as relevant to CO₂-induced panicogenic responses (Wemmie et al., 2003; Coryell et al., 2009). Currents originating from the ASIC1a channel can be inhibited by selective

modulators, such as Psalmotoxin-1 (Pstx-1) (Escoubas et al., 2003; Dawson et al., 2012). Pstx-1 specifically inhibits ASIC1a, without inhibiting any other ASIC channel, binding to these channels to change their conformation, causing their affinity to H⁺ to change (Escoubas et al., 2000; Chen et al., 2006).

It is well known that the sex of the individual can affect breath control and susceptibility to some respiratory diseases (Gargaglioni et al., 2019; Holley et al., 2012; Jiang et al., 2017). Women are two to three times more likely to develop PD than men during the life (Kessler et al., 2005; Sheikh et al., 2002; Yonkers et al., 1998). However, the role of ASIC1a channels located in LC in CO₂-induced hyperventilation and panic remains unknown, as well as their differences regarding sex. Therefore, we sought to test the hypothesis that LC is an important site of ASIC1A action in CO₂-induced panic behaviors and breathing control in male and female mice.

Methods

Animals

The experiments were performed with male and female mice C57BL/6, weighting around 20 to 25 g (8 weeks old). All the animals were housed in a temperature-controlled room, maintained at 25 ± 1°C with a 12 h light-dark cycle (lights on at 6:30 a.m.), with water and food provided *ad libitum*.

Experiments were performed between 8:00 a.m. and 1:00 p.m. at the Department of Animal Morphology and Physiology of the Faculty of Agricultural and Veterinary Sciences – FCAV/UNESP, campus of Jaboticabal. All experimental protocols were approved by the local ethics committee (CEUA - protocol n°3451/22), being in norm with the National Council for the Control of Animal Experimentation (CONCEA-MTC, Brazil).

Surgical procedures

The animals were anesthetized with isoflurane (5% for induction and 1.5% for maintenance). After anesthetic induction, the animals were fixed in a stereotactic apparatus (Kopf Instruments, Kent, England) and a stainless-steel guide cannula was implanted (10 mm in length and 0.7 mm in external diameter) aimed at the LC.

The coordinates (Paxinos and Franklin, 2003) used were antero posterior (AP): -5.2 mm from the bregma; lateral (LL): ± 0.8 mm, and dorso ventral (DV): -3.9 mm from the skull for males and AP: -5.2 mm; LL: ± 0.9 mm and DV: -4.0 mm for females. The cannula was fixed to the skull by using dental acrylic and sealed with a stainless-steel wire to prevent occlusion and infection.

At the end of surgery, the animals were placed in boxes with water and food "*ad libitum*" and kept in a room with controlled temperature, humidity, and luminosity. Antibiotic injection (Enrofloxacin, Baytril; 2.5 mg/kg s.c.) and analgesic (Flunixin Meglumine, Banamine; 10 mg/kg s.c.) were administered in a single dose after surgery.

Determination of body temperature

T_c was obtained by implanting a temperature sensor (Biomark, USA) in the abdominal cavity on the same day that the stereotactic surgery was performed.

Drug

Psalmotoxin-1 (Pstx-1) (Sigma-Aldrich) solubilized in sterile saline 0.9% and was applied in the LC in a concentration of 50 ng/0.1 μ L. The dose chosen was based on studies carried out in the laboratory.

Induction of panic attacks

The mice were placed in a round acrylic chamber of 5 liters and exposed to a respiratory challenge composed of a gas mixture of 20% CO₂, 21% of O₂ and N₂ to balance during the time of 15 minutes. Exposure to respiratory challenge was used to induce behavioral and ventilatory panic responses in animals.

Pulmonary ventilation assessment

The pulmonary ventilation (V_E) of mice were acquired by whole body plethysmography – closed system. For V_E measurements, the chamber was fully sealed for approximately 2 min. The animal was placed in a 5L experimental chamber for a habituation period of 1 hour. Air inlet and outlet flows were 1,800 L/min measured by a flowmeter. The pressure oscillation due to temperature difference

from inhaled and exhaled air were monitored by a differential pressure transducer (TSD 160A, Biopac Systems, Santa Barbara, CA, USA) and fed into a pre-amplifier (DA 100C, Biopac Systems), passed through an analog-to-digital converter and digitized on computer equipped with data acquisition software (MP100ACE, Biopac Systems). A volume calibration was performed by injecting 1 mL of air into the chamber using a graduated syringe. Tidal volume (V_T) was calculated using Drorbaugh and Fenn's formula (1955):

$$V_T = V_K \times (P_T/P_K) \times T_B \times (P_B - P_C) / T_B \times (P_B - P_C) - T_A \times (P_B - P_R)$$

where P_T : pressure deflection associated with each V_T , P_K : pressure deflection associated with the injection of the calibration volume (V_K), T_A : air temperature in the animal chamber, P_B : barometric pressure, P_C : water vapor pressure in the animal chamber, T_B : body temperature (in Kelvin), and P_R : water vapor pressure at T_C .

The V_E was calculated as the product of respiratory frequency (f_R) and V_T and normalized to the animal's body weight. V_E and V_T were presented under ambient barometric pressure conditions at T_B and saturated with water vapor (BTPS). P_C and P_R were calculated indirectly using an appropriate table (Dejours, 1981). The LabChart software (PowerLab System, ADInstruments®/LabChart Software, version 7.3, Sydney, Australia) was used for data analysis.

Determination of oxygen consumption

The indirect calorimetry method by measuring O_2 consumption (VO_2) with flow through push mode configuration was used for metabolic rate inference. The chamber air flow (1.5 L/min) was maintained through a compressor pump that directed the air to the chamber entrance when the animals were exposed to room air and by gas mixture cylinders coupled to a flowmeter when exposed to hypercapnia. The expired gas was dried over a small column of Drierite (W.A. Hammond Drierite Co. Ltd, Xenia, OH, USA) before goes through the gas analyzer. The air was continuously sampled allowing for the determination of VO_2 by a data acquisition program (Power-Lab System, ADInstruments® / Chart Software, version

7.3, Sydney, Australia). The CO₂ was not analysed or scrubbed, then aiming at a better metabolic rate measurement the VO₂ was calculated using the following equation (Koteja, 1996):

$$\dot{V}O_2 = [FRe (\dot{F}iO_2 - \dot{F}eO_2)] / [1 - \dot{F}iO_2 (1 - RQ)]$$

where FRe: end flow rate of air through the chamber, $\dot{F}iO_2$: inlet O₂ fraction, $\dot{F}eO_2$: end O₂ fraction, and RQ: respiratory quotient (considered to be 0.85). The $\dot{V}O_2$ as divided for the body mass and values were presented as STPD (standard conditions of temperature, pressure and dry air).

Behavioral measurements

To analyze the response of the panic attack behavioral, the experiments were recorded by a camera (Logitech, USA), placed to clearly see the animal and its movements. The escape response was analyzed, being considered the number of jumps. The freezing behavioral (total lack of movement except breathing) and the rearing (standing on the hind legs with or without contact on the anterior legs on the side of the chamber) was evaluated by a trained observer.

Central microinjection

For intra-LC microinjection, a needle (10 mm long, 0.3 mm outer diameter) was introduced through the guide cannula, using a 5 µL microsyringe (Hamilton, USA). The volume of injections was 0.1 µL for 1 minute using a microinjection machine (model 310, Stoelting Co., IL, EUA). The displacement of an air bubble inside the polyethylene catheter connecting to the syringe was used to monitor the microinjection.

Experimental protocol

All the animals were exposed to the experimental chamber under the condition of normocapnia (ambient air 21% O₂ and 0,03% CO₂) for a habituation period of approximately 1 hour before the experiment began (Figure 1).

Seven days after stereotactic surgery, the animals underwent the experimental procedure. At first, two control ventilatory and metabolic

measurements were performed in ambient air after 10 and 20 minutes of habituation. Escape responses were recorded during all experiment. Subsequently, 0.1 μ L of vehicle of saline or Pstx-1 (50ng) was microinjected into the mice LC. After 15 min of the microinjection, respiratory and metabolic measurements were performed under normocapnic conditions. Then, the respiratory challenge was presented (20% CO₂) for 15 minutes and after this period respiratory and metabolic measurements were performed again. Behavioral panic responses were recorded throughout the experiment. However, behavioral data were be presented only during the period of hypercapnia since there was no behavioral change during the period of normocapnia.

Confirmation of the microinjection site

For the determination of the microinjection site the animals were deeply anesthetized and 0.1 μ L of Evans blue was injected through the guide cannula. The animals were perfused transcardially with phosphate buffered saline (PBS) 0.1 M, followed by paraformaldehyde 4%. The brain was extracted and maintained in paraformaldehyde 4% for 24 hours. Coronal sections of the brain at 35 μ m were made using a cryostat and deposited in slides, for later visualization under a microscope.

Data analyses

The data were evaluated in terms of meeting the assumptions of homogeneity of variances and normality of the errors. Then, they were submitted to analysis of variance by the GLM procedure (General Linear Models) and, in case of significant difference, the means were compared by the Tukey test with probability level of 5% and confidence interval of 95%. For all ventilatory and metabolic analyses, the repeated measures two-way ANOVA test was performed with the following factors: treatment (Pstx-1 or saline) and gas condition (normocapnia and hypercapnia). For behavioral analysis, the T Student test was used to analyze the number of jumps, rearing, running and freezing.

Results

Figure 2 represents the representative section of the brainstem of an animal, demonstrating the confirmation of the microinjection site in the LC. It is possible to

observe the presence of Evans blue dye restricted to LC. Only animals with blue color in this region were incorporated to the data analysis.

Effect of intra-LC microinjection of Pstx-1 on ventilation and metabolism under normocapnia and hypercapnia in males and females

In Figure 3, the values of V_E , V_T and f_R of males (Figure 3A, 3B and 3C) and females (Figure 3D, 3E and 3F) treated with Pstx-1 or saline in normoxia before and post-injection and hypercapnia condition are represented. The two-way ANOVA revealed significant effects of gas condition in V_E [males: $F(2,35)= 74.227$, $p<0.001$; females: $F(2,35)= 43.107$, $p<0.001$], V_T [males: $F(2,35)= 140.923$, $p<0.001$; females: $F(2,35)= 39.440$, $p<0.001$] and f_R [males: $F(2,35)= 4.943$, $p=0.032$; females: $F(2,35)= 16.934$, $p<0.001$]. As can be seen in the Figure 3C, in males the treatment with Pstx-1 tend to decreased the f_R in the animals exposed to CO_2 [$F(1,35)= 6.478$; $p=0.052$]. In females, treatment with Pstx-1 did not change any of the analyzed parameters. For the other parameters, there is no significant statistics difference between saline and Pstx-1 groups.

In Figure 4, the values of VO_2 , V_E/VO_2 and Tb of males (Figure 4A, 4B and 4C) and females (Figure 4D, 4E and 4F) treated with Pstx-1 or saline in normoxia before and post-injection and hypercapnia condition are represented. For males, The two-way ANOVA revealed significant effects of treatment [$F(1,35)= 15.551$, $p=0.011$] and gas condition [$F(2,35)= 104.126$, $p<0.001$] in Tb , but not in VO_2 and V_E/VO_2 . For females, there is significant effect of gas condition in V_E/VO_2 [$F(2,35)= 5.800$, $p=0.021$] and Tb [$F(2,35)= 167.175$, $p<0.001$], but not in VO_2 . There is no significant statistics difference between saline and Pstx-1 females.

Effect of intra-LC microinjection of Pstx-1 on panic-related responses under normocapnia and hypercapnia in males and females

Figure 5 shows the effect of intra-LC administration of Pstx-1 on jumping, freezing, running, and rearing behaviors expressed by male (Figure 5A, 5B, 5C and 5D) and female (Figure 5E, 5F, 5G and 5H) mice exposed to 20% CO_2 . For males, the

Student T test showed that there was a significant difference between Pstx- 1 and saline group in the number of jumps [$t(10)$ 3.412, $p=0.006$] and rearing [$t(12)$ 2.491, $p=0.028$], but not in freezing [$t(10)$ 0.264, $p=0.797$]. As can be seen in the Figure 5C, the treatment with Pstx-1 tend to decreased the running [$t(10)$ 1.990, $p=0.074$] in the animals exposed to CO₂.

For females, there was no significant difference between Pstx-1 and saline group in the number of jumps [$t(10)$ 1.542, $p=0.154$], freezing [$t(10)$ 0.629, $p=0.543$], running [$t(10)$ 0.859, $p=0.410$] and rearing [$t(10)$ 0.868, $p=0.405$].

Discussion

In the present study, we examined a potential role for ASIC1a channels in the LC in the hypercapnic ventilatory, metabolic and behavioral response of male and female C57BL/6 mice. Our results indicate that this channel in the LC do indeed play a role in the hypercapnic behavioral response in males since their blockade with Pstx- 1 resulted in a significant reduction in the number of jumps and rearing. However, the behavioral data found in females demonstrated no different response to respiratory challenge. Regarding ventilation data in males and females, ASIC1a blockade in the LC did not change ventilation, metabolic rate and respiratory equivalent (V_E/VO_2) during room air and hypercapnic conditions, suggesting that this channel in the LC does not participate in breathing and metabolic control. In relation to Tb, there was a significant attenuation in the drop of Tb under CO₂ exposure only in male animals treated with Pstx-1, indicating that ASIC1a located in LC region plays a role in the hypothermic response induced by hypercapnia.

Under normocapnic conditions, Pstx-1 microinjection into the LC did not alter the ventilatory or metabolic parameters in females suggesting that ASIC1a in this nucleus does not play a tonic respiratory role. In males, a decrease in f_R was observed, however, this reduction was not enough to change ventilation. Our findings support data from previous studies from our laboratory showing that removal of LC noradrenergic neurons does not affect cardiorespiratory control during resting conditions (Biancardi et al., 2008, Carvalho et al., 2010).

Regarding the ventilatory response to 20% CO₂, the ventilatory parameters

increased in relation to exposure to normocapnia in all groups independent of sex. However, ASIC1a blockade did not change the hypercapnic ventilatory response in males and females, indicating that this channel located in LC does not participate in the hyperventilatory response induced by CO₂. In relation to the ventilatory response to 20% CO₂, the ventilatory parameters exhibited an increase across all groups, irrespective of sex. However, it is noteworthy that blocking the ASIC1a channel did not alter the hypercapnic ventilatory response in both males and females. Indicating that this acid channel, situated in the LC, does not play a role in the ventilatory response induced by high level of CO₂. A previous study has demonstrated that Pstx-1 injection (10 nM/0.1 µl) in the lateral hypothalamus abolished the increase of phrenic nerve activity induced by acidification of this region, which suggests that ASIC1a is responsible for acid sensing in the hypothalamus of Sprague-Dawley male rats (Song et al., 2012). The same pattern of response was observed in the ventrolateral medulla in rats using a higher dose (100 nM/0.1 µl; Song et al., 2016). Therefore, ASIC1a channels seem to be important to the increase in respiration in some CO₂/pH chemosensitive areas such as hypothalamus and medulla, but not in LC. As previously mentioned, LC acts as a central chemoreceptor in mammals, (Elam et al., 1981; Filosa et al., 2002; Gargaglioni et al., 2010; Imber et al., 2014; Pineda e Aghajanian, 1997) and it is known to send projections to several areas of the central nervous system, including those responsible for respiratory control and for activating behavioral responses related to panic, such as the amygdala (McCall, et al., 2017; Giustino et al., 2020).

In the present study, the observed behavioral pattern exhibited by the animals during exposure to 20% CO₂ aligns with existing literature, illustrating an augmentation in the frequency of jumps and other responses typical of fight-or-flight behaviors in the presence of elevated CO₂ concentrations (Spiacci et al., 2018, Taugher et al., 2019). Indeed, studies conducted in mice show that inhalation of 20% CO₂ can trigger behavioral responses such as freezing and avoidance (D'Amato et al., 2011; Leibold et al., 2016; Taugher et al., 2014; Vollmer et al., 2016; Ziemann et al., 2009) suggesting that the animals are facing an intense aversive situation, which resembles the triggering of a panic attack (Spiacci et al., 2018). Although the amygdala may be more important in the conditioning process that leads to avoidance behavior, a previous study has demonstrated that three patients with

focal bilateral amygdala lesions (due to Urbach-Wiethe disease) experienced panic attacks following a CO₂ challenge suggesting that other regions besides amygdala is involved in triggering panic attacks (Goossens et al., 2014). One of the regions suggested by the authors was LC. In fact, we have demonstrated that noradrenaline originating from the LC has an important role in the characterization of jumping and freezing behaviors in both male and female mice, since mutant males lacking or with reduced LC noradrenaline (NA) synthesis had a blunted jumping response and an increased freezing duration compared to wild type animals (Ripamonte et al., in revision).

In the study conducted by Ziemann et. (2009) it was demonstrated that inhaled CO₂ evoked fear behavior in male mice and elimination or inhibition of ASIC1a impaired the activity of the animals. Interestingly, the localized expression of ASIC1a in the amygdala rescued the fear deficit induced by CO₂ in ASIC1a^{-/-} animals (Ziemann et., 2009). In addition, the nucleus of the stria terminalis (BNST) is considered a key structure in fear-related behaviors evoked by CO₂ (Taughner et al., 2014). BNST mediates CO₂-evoked freezing and conditioned avoidance and the data found in the study also suggest that CO₂ acidifies BNST and that this region can directly detect acidosis through ASIC1a. The interruption of these channels specifically in the BNST reduced the freezing evoked by CO₂, while the expression of ASIC1a mediated by virus vector in the BNST of mice ASIC1a^{-/-} and ASIC1a^{+/+} increased the freezing evoked by CO₂.

The results observed in the study suggest that the inhibition of ASIC1a channels present in the LC has sex-specific effects on the behavioral response of mice exposed to high concentrations of CO₂. We observed that Pstx-1 only reduced the CO₂ behavioral responses in males. Our data add to the scientific literature the important participation of ASIC1a located in the LC are also involved in the expression of flight behavior. However, for females we still find lags in the literature, and this may hinder possible correlations, such as the estrous/menstrual cycle being related to PD.

Interestingly, a study from our laboratory demonstrated that exposure to 20% CO₂ activates LC-NA neurons of male mice while female counterparts do not, highlighting a notable disparity in CO₂ response between the sexes (Ripamonte et al., in revision). Additionally, the release of noradrenaline from the LC is important

for the tonic excitatory control of ventilation in male mice, but not in females. One could argue that the difference in panic-related behavior may be due to β -estradiol. As demonstrated by Zhou et al. (2019), treatment with β -estradiol suppressed the expression of ASIC1a due to increased protein degradation. It has also been shown that the expression level of ASIC1a protein is significantly lower in the cerebral cortex of female mice, which is correlated with a lower degree of brain damage and protection by ASIC1a inhibitor (Zhou et al., 2019). These differences can, however, be attenuated by ovariectomy. Therefore, these results indicated that β -estradiol can protect neurons against acidosis-mediated neurotoxicity and ischemic brain injury, reducing ASIC1a protein expression and channel function in females.

In conclusion, our data suggest that ASIC1a channels located in the LC do not participate in respiratory control under room air and hypercapnia, but are involved in CO₂-induced panic behavior in males, demonstrating a specific sex-dependent response. The results of the present study extend to previous studies on the mechanisms of PD and provide evidence regarding behavioral responses between males and females to a CO₂ panic challenge.

References

- AMERICAN PSYCHIATRIC ASSOCIATION. DSM 5. 2013. Washington, DC: American Psychiatric Association. 2013.
- Balaban CD (2002) Neural substrates linking balance control and anxiety. **Physiology & behavior** 77:469-475.
- Bangasser DA, Valentino RJ (2014) Sex differences in stress-related psychiatric disorders: neurobiological perspectives. **Frontiers in neuroendocrinology** 35:303-319.
- Biancardi V, Bicego KC, Almeida MC, Gargaglioni LH (2008) Locus coeruleus noradrenergic neurons and CO₂ drive to breathing. **Pflügers Archiv-European Journal of Physiology** 455:1119-1128.
- Bremner JD, Krystal JH, Southwick SM, Charney DS (1996) Noradrenergic mechanisms in stress and anxiety: I. Preclinical studies. **Synapse** 23:28-38.
- Borodovitsyna O, Flamini MD, Chandler DJ (2018) Acute stress persistently alters locus coeruleus function and anxiety-like behavior in adolescent rats. **Neuroscience** 373:7-19.

Carvalho D, Bicego KC, de Castro OW, da Silva GS, Garcia-Cairasco, N, Gargaglioni LH (2010) Role of neurokinin-1 expressing neurons in the locus coeruleus on ventilatory and cardiovascular responses to hypercapnia. **Respiratory physiology & neurobiology** 172:24-31.

Charney DS, Woods SW, Nagy LM, Southwick SM, Krystal JH, Heninger GR (1990) Noradrenergic function in panic disorder. **The Journal of clinical psychiatry** 51:5-11.

Chen X, Kalbacher H, Grunder S (2006) Interaction of acid-sensing ion channel (ASIC)1 with the tarantula toxin psalmotoxin 1 is state dependent. **The Journal of general physiology** 127:267-276.

Coryell MW, Wunsch AM, Haenfler JM, Allen JE, Schnizler M, Ziemann AE, Wemmie JA (2009) Acid-sensing ion channel-1a in the amygdala, a novel therapeutic target in depression-related behavior. **Journal of Neuroscience** 29:5381-5388.

D'Amato FR, Zanettini C, Lampis V, Coccurello R, Pascucci T, Ventura R, Battaglia M (2011) Unstable maternal environment, separation anxiety, and heightened CO₂ sensitivity induced by gene-by-environment interplay. **PLoS One** 6:e18637.

Dawson RJ, Benz J, Stohler P, Tetaz T, Joseph C, Huber S, Ruf A (2012) Structure of the acid-sensing ion channel 1 in complex with the gating modifier Psalmotoxin 1. **Nature communications** 3:936.

Dierssen M, Fotaki V, de Lagrán MM, Gratacos M, Arbones M, Fillat C, Estivill X (2002) Neurobehavioral development of two mouse lines commonly used in transgenic studies. **Pharmacology Biochemistry and Behavior** 73:19-25.

Drorbaugh JE, Fenn WO (1955) A barometric method for measuring ventilation in newborn infants. **Pediatrics** 16:81-87.

Elam M, Yao T, Thore P, Svensson TH (1981) Hypercapnia and hypoxia: chemoreceptor-mediated control of locus coeruleus neurons and splanchnic, sympathetic nerves. **Brain research** 222:373-381.

Escoubas P, De Weille JR, Lecoq A, Diochot S, Waldmann R, Champigny G, Lazdunski M (2000) Isolation of a tarantula toxin specific for a class of proton-gated Na⁺ channels. **Journal of Biological Chemistry** 275:25116-25121.

Escoubas P, Bernard C, Lambeau G, Lazdunski M, Darbon H (2003) Recombinant production and solution structure of PcTx1, the specific peptide inhibitor of ASIC1a proton-gated cation channels. **Protein Science** 12:1332-1343.

Esquivel G, Schruers KR, Maddock RJ, Colasanti A, Griez EJ (2010). Acids in the brain: a factor in panic?. **Journal of psychopharmacology** 24:639-647.

Filosa JA, Dean JB, Putnam RW (2002) Role of intracellular and extracellular pH in

the chemosensitive response of rat locus coeruleus neurones. **The Journal of physiology** 54:493-509.

Fortress AM, Hamlett ED, Vazey EM, Aston-Jones G, Cass WA, Boger HA, GranholmACE (2015) Designer receptors enhance memory in a mouse model of Down syndrome. **Journal of Neuroscience** 35:1343-1353.

Gargaglioni LH, Hartzler LK, Putnam RW (2010) The locus coeruleus and central chemosensitivity. **Respiratory physiology & neurobiology** 173:264-273.

Gargaglioni LH, Marques DA, Patrone LGA (2019) Sex differences in breathing. **Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology** 238:110543.

Giustino TF, Ramanathan KR, Totty MS, Miles OW, Maren S (2020) Locus coeruleus norepinephrine drives stress-induced increases in basolateral amygdala firing and impairs extinction learning. **Journal of Neuroscience** 40:907-916.

Goossens L, Leibold N, Peeters R, Esquivel G, Knuts I, Backes W, Schruers K (2014) Brainstem response to hypercapnia: a symptom provocation study into the pathophysiology of panic disorder. **Journal of psychopharmacology** 28:449-456.

Grant BF, Hasin DS, Stinson FS, Dawson DA, Goldstein RB, Smith S, Saha TD (2006) The epidemiology of DSM-IV panic disorder and agoraphobia in the United States: results from the National Epidemiologic Survey on Alcohol and Related Conditions. **The Journal of clinical psychiatry** 67: 6517.

Holley HS, Behan M, Wenninger, JM (2012) Age and sex differences in the ventilatory response to hypoxia and hypercapnia in awake neonatal, pre-pubertal and young adults. **Respiratory physiology & neurobiology** 180:79-87.

Imber AN, Santin JM, Graham CD, Putnam RW (2014) A HCO₃⁻-dependent mechanism involving soluble adenylyl cyclase for the activation of Ca²⁺ currents in locus coeruleus neurons. **Biochimica et Biophysica Acta (BBA)-Molecular Basis of Disease** 1842:2569-2578.

Isingrini E, Perret L, Rainer Q, Amilhon B, Guma E, Tanti A, Giros B (2016) Resilience to chronic stress is mediated by noradrenergic regulation of dopamine neurons. *Nature neuroscience*, 19(4), 560-563.

Jiang C, Cui N, Zhong W, Johnson CM, Wu Y (2017) Breathing abnormalities in animal models of Rett syndrome a female neurogenetic disorder. **Respiratory physiology & neurobiology** 245:45-52.

Kessler RC, Chiu WT, Demler O, Walters EE (2005) Prevalence, severity, and comorbidity of 12-month DSM-IV disorders in the National Comorbidity Survey Replication. **Archives of general psychiatry** 62:617-627.

Koteja P (1996) Measuring Energy Metabolism with Open -Flow Respirometric

Systems: Which Design to Choose? **Funct. Ecol** 10:675.

Leibold NK, Van Den Hove DLA, Viechtbauer W, Buchanan GF, Goossens L, Lange I, Schruers KRJ (2016) CO₂ exposure as translational cross-species experimental model for panic. **Translational psychiatry** 6:e885-e885.

McCall JG, Siuda ER, Bhatti DL, Lawson LA, McElligott ZA, Stuber GD, Bruchas MR (2017) Locus coeruleus to basolateral amygdala noradrenergic projections promote anxiety-like behavior. **Elife** 6:e18247.

Paxinos G, Franklin KB (2003) Paxinos and Franklin's the mouse brain in stereotaxic coordinates. Academic press.

Pineda J, Aghajanian GK (1997) Carbon dioxide regulates the tonic activity of locus coeruleus neurons by modulating a proton- and polyamine-sensitive inward rectifier potassium current. **Neuroscience** 77:723-743.

Putnam RW, Filosa JA, Ritucci, NA (2004) Cellular mechanisms involved in CO₂ and acid signaling in chemosensitive neurons. **American Journal of Physiology-Cell Physiology** 287:C1493-C1526.

Song N, Zhang G, Geng W, Liu Z, Jin W, Li L, Shen L (2012) Acid sensing ion channel 1 in lateral hypothalamus contributes to breathing control. **PloS one** 7:e39982.

Sheikh JI, Leskin GA, Klein DF (2002) Gender differences in panic disorder: findings from the National Comorbidity Survey. **American Journal of Psychiatry** 159:55-58.

Song N, Guan R, Jiang Q, Hassanzadeh CJ, Chu Y, Zhao X, Shen L (2016) Acid - sensing ion channels are expressed in the ventrolateral medulla and contribute to central chemoreception. **Scientific reports** 6:38777.

Spiacchi A, Vilela-Costa HH, Sant'Ana AB, Fernandes GG, Frias AT, da Silva GSF, Zangrossi H (2018) Panic-like escape response elicited in mice by exposure to CO₂, but not hypoxia. **Progress in Neuro-Psychopharmacology and Biological Psychiatry** 81:178-186.

Ripamonte GC, Fonseca EM, Frias AT, Patrone LGA, Vilela-Costa HH, Costa Silva KS, Szawka RE, Bícigo KC, Zangrossi Jr H, Plummer NW, Jensen P, Gargaglioni LH. Locus coeruleus noradrenaline depletion and its differential impact on CO₂-induced panic and hyperventilation in male and female mice. *In revision*.

Taughner RJ, Lu Y, Fan R, Ghobbeh A, Kreple CJ, Faraci FM, Wemmie JA (2017) ASIC1A in neurons is critical for fear-related behaviors. **Genes, Brain and Behavior** 16:745-755.

Taughner RJ, Lu Y, Wang Y, Kreple CJ, Ghobbeh A, Fan R, Wemmie JA (2014) The bed nucleus of the stria terminalis is critical for anxiety-related behavior evoked by CO₂ and acidosis. **Journal of Neuroscience** 34:10247-10255.

Taugher RJ, Dlouhy BJ, Kreple CJ, Ghobbeh A, Conlon MM, Wang Y, Wemmie JA (2019) The amygdala differentially regulates defensive behaviors evoked by CO₂. **Behavioural brain research** 377:112236.

Vollmer LL, Strawn JR, Sah R (2015) Acid–base dysregulation and chemosensory mechanisms in panic disorder: a translational update. **Translational psychiatry**, 5:e572-e572.

Weinshenker D (2018) Long road to ruin: noradrenergic dysfunction in neurodegenerative disease. **Trends in neurosciences** 41:211-223.

Wemmie JA, Askwith CC, Lamani E, Cassell MD, Freeman JH, Welsh MJ (2003) Acid-sensing ion channel 1 is localized in brain regions with high synaptic density and contributes to fear conditioning. **Journal of Neuroscience** 23:5496-5502.

Yonkers KA, Zlotnick C, Allsworth J, Warshaw M, Shea T, Keller MB (1998) Is the course of panic disorder the same in women and men?. **American Journal of Psychiatry** 155:596-602.

Zhou R, Leng T, Yang T, Chen F, Hu W, Xiong ZG (2019) β -Estradiol protects against acidosis-mediated and ischemic neuronal injury by promoting ASIC1a (acid-sensing ion channel 1a) protein degradation. **Stroke** 50:2902-2911.

Zhu Y, Hu X, Wang L, Zhang J, Pan X, Li Y, Huang Y (2022) Recent advances in acid- sensitive ion channels in central nervous system diseases. **Current Pharmaceutical Design** 28:1406-1411.

Ziemann AE, Allen JE, Dahdaleh NS, Drebot II, Coryell MW, Wunsch AM, Wemmie JA (2009) The amygdala is a chemosensor that detects carbon dioxide and acidosis to elicit fear behavior. **Cell** 139:1012-102.

Figure Legends

Fig 1. Illustrative figure of the experimental protocol.

Fig 2. Photomicrographs of cross-sections of the pons of mice showing the site of microinjection of Pstx-1 or saline in the LC with the presence of Evans blue.

Fig 3. Effect of intra-LC microinjection of Pstx-1 or saline on ventilation (VE) [A, D], tidal volume (VT) [B, E] and respiratory frequency (fR) [C, F], during resting conditions in normocapnia or hypercapnia (20% of CO₂) in male [A, B, C] and female [D, E, F] mice (n=6 animals/group).

Fig 4. Effect of intra-LC microinjection of Pstx-1 or saline on ventilation oxygen consumption (VO₂) [A, D], respiratory equivalent (VE/VO₂) [B, E] and body temperature (Tb) [C, F] during resting conditions in normocapnia or hypercapnia (20% of CO₂) in male [A, B, C] and female [D, E, F] mice (n=6 animals/group).

Fig 5. Effects of intra-LC microinjection with Pstx-1 or saline on the number of jumps [A, E], % freezing time (s) [B, F], running frequency [C, G] and rearing frequency [D, H] of male [A, B, C, D] and female [E, F, G, H] mice submitted to normocapnia and 20% CO₂ (n=6 animals/group). Pstx-1: psalmotoxin-1. *p<0.05 compared to the control group.

Figures

Fig. 1




Figure 1

Fig 2.



Fig 3.






Fig 4.


A




D




B



E



C



F




Fig 5.

