

Bats (Mammalia: Chiroptera) from Guarulhos, state of São Paulo, Brazil

Maria Ester Chaves^{1*}, Wilson Uieda², Cristiane Espinosa Bolochio³, Cláudia Almeida Igayara de Souza³, David de Almeida Braga³, Carlos Henrique Ferreira³, Carlos Leandro Firmo¹, Rebeca G. G. Calaça Mariano¹, Kelly Cardinale Silva de Oliveira¹, Eduardo Guilherme dos Santos¹ and Fábio Moreira da Costa^{3,4}

1 Universidade Guarulhos, Campus Centro, Laboratório de Biociências, Grupo de Zoologia e Saúde Pública. CEP 07023-070. Guarulhos, SP, Brazil.

2 Universidade Estadual Paulista, Instituto de Biociências, Departamento de Zoologia. CEP 18618-970. Botucatu, SP, Brazil.

3 Zoológico de Guarulhos, Grupo de Fauna. Secretaria do Meio Ambiente de Guarulhos (SEMA). CEP 07081-120. Guarulhos, SP, Brazil.

4 Divisão Técnica para Conservação da Biodiversidade e Educação Ambiental (SEMA). CEP 07114-010. Guarulhos, SP, Brazil.

* Corresponding author. E-mail: mechaves@ig.com.br

ABSTRACT: The present study provides a record of the bat fauna captured in Guarulhos, state of São Paulo, Brazil, between February and November 2011. The capture sessions were carried out using mist nets set on trails, around and above streams and lakes, near built areas and around livestock. After a total effort of 26,208 m².h, we recorded 17 bat species belonging to Phyllostomidae (13 species), Vespertilionidae (3) and Molossidae (1). *Sturnira lilium* (73 captures), *Artibeus fimbriatus* (69), and *A. lituratus* (59) were the most abundant species, corresponding to 57.8% of all captures. Some information about shelters was also commented. This paper is the first to report *Molossops neglectus* for Guarulhos.

INTRODUCTION

During the last decades, Guarulhos had a population growth rate higher than the average of its region and higher even than that of the city of São Paulo (state capital). As a result of the urban expansion, the native vegetation was removed, giving way to commercial, industrial, and service activities (Oliveira and Andrade 2008). In spite of this, several forest fragments remained in Guarulhos, most of which located in the north and northeastern regions of the municipality (33% of the fragments). Thus, these areas also comprise the greatest biodiversity found in Guarulhos (Andrade *et al.* 2008).

Although many faunal inventories have been carried out in southeastern Brazil, in Guarulhos they are still scarce. The latest initiative to counteract this scarcity was the project “Guarulhos has Biodiversity”, which started officially in 2007 (Diário Oficial do Município de Guarulhos N° 042/2007). Our study on the bat fauna of Guarulhos is part of this project.

MATERIAL AND METHODS

Study area

Guarulhos is located in the transition between tropical and temperate zones; it has a mesothermal humid climate, with two or three dry months in the winter. Annual mean temperature ranges from 15° to 24°C and rainfall ranges from 1,250 to 1,500 mm. Serra do Itaberaba is the highest point located at northeastern Guarulhos, with 1,422 m of altitude, and preserves most of the municipality vegetation, described as Ombrophylous Dense Forest by Andrade *et al.* (2008).

Sampling sites

The present study was carried out in 10 different sites (Figure 1) in Serra do Itaberaba, Guarulhos, state of São

Paulo, around the State Park of Itaberaba, created in 2010. Most of the sampling sites were private properties with animal-raising areas surrounded by fragments of Atlantic Forest. The sites were the following: (A) a private property situated on a hill, along a stream with riparian vegetation, and some domestic animals (chickens, pigs and cows) (23°20' S, 46°23' W); (B) a fishing farm with artificial lakes for fishing activity, fish ponds, streams with riparian vegetation, a restaurant, and the owner's house (23°20' S, 46°24' W); (C) a farm with forest fragments, a stream, ponds, animal-raising such as chickens, cattle, sheep, goats, ostrich, and dogs (23°20' S, 46°24' W); (D) a forest park (Horto Florestal de Guarulhos) with fragments of Atlantic Forest, greenhouses, and seedlings (23°22' S, 46°23' W); (E) an area with a historic mansion called “Candinhas House” surrounded by forest fragments and streams with riparian vegetation (23°23' S, 46°28' W); (F) a private property with a hennery, an artificial lake, a stream with riparian vegetation, and the owner's residence (23°20' S, 46°24' W); (G) a private property that was formerly used as a recreational area, with open spaces and trails through the Atlantic Forest and a tunnel of groundwater that flows into an artificial lake (23°21' S, 46°24' W); (H) a crops farm with a hennery, forest fragments, swamps, and abandoned buildings (23°20' S, 46°24' W); (I) an area with a hennery, a corral, some goats and turkeys, an abandoned house, and other uninhabited buildings surrounded by forest (23°18' S, 46°22' W); and (J) a green area with trails, artificial lakes, and forest fragments owned by a beverage company (23°18' S, 46°23' W).

Capture Sessions

The bat sampling was conducted during 18 nights, from February to November 2011, using six to nine mist nets each night. The mist nets size ranged from 7x3 m

to 12x3 m. They were opened during the first six hours after sunset, 0.2 m to 3.0 m high, in trails at the border and inside forest fragments, around and above streams and lakes, and near building areas, depending on the site options. At places with livestock, some nets were set up around the animals in order to capture vampire bats.

Captured bats were identified, measured, and released, except for one individual of each species by sampling night. These specimens were deposited as voucher in the Vertebrate Collection of Department of Zoology, Institute of Biosciences, UNESP, Botucatu campus (Appendix 1). Individuals were identified according to Vizotto and Taddei (1973), Gregorin and Taddei (2002), LaVal (1973), and other references (Taddei et al. 1998; Gregorin et al. 2004; Bernardi et al. 2007; Araújo and Langguth 2010). The capture and collection of all bats reported here were conducted under the permanent license of Wilson Uieda (IBAMA nº 23151-1).

RESULTS AND DISCUSSION

We carried out 108 sampling hours, comprising a total effort of 26,208 m².h (Straube and Bianconi 2002). A total of 17 bat species of three families (Phyllostomidae, Vespertilionidae and Molossidae) were recorded in 348 captures (Table 1). Phyllostomidae was the richest group with 13 species, which corroborates the majority of the bat fauna surveys of the Atlantic Forest (Reis et al. 2000; Falcão et al. 2003; Bianconi et al. 2004; Bordignon 2006; Dias and Peracchi 2008) that used mist nets as the main sampling methodology. Mist net is an useful method for sampling phyllostomids and other bats that forage in the understory (Kalko 1998; Straube and Bianconi 2002).

Most captured species belong to Stenodermatinae (Phyllostomidae). This group has 12 genera and 33 species

with occurrence in Brazil (Zortéa 2007). Of these, six (*Artibeus lituratus*, *A. fimbriatus*, *Pygoderma bilabiatum*, *Platyrrhinus lineatus*, *Sturnira lilium*, and *Carollia perspicillata*) were recorded by us in Serra do Itaberaba.

Similar to other inventories in secondary forests (Falcão et al. 2003; Passos et al. 2003; Zanon and Reis 2007), our study showed that *S. lilium* (73 captures), *A. fimbriatus* (69), and *A. lituratus* (59) were the most abundant species, corresponding to 57.8% of all individuals captured. The former species is common in open and secondary growth areas and feeds on a variety of fruits, especially pioneer plants (Bredt et al. 2012), as well as on pollen and insects (Gardner 1977). According to Fleming (1988), *S. lilium* could be classified as an understory frugivore. The two other bat species feed mainly on Moraceae (especially *Ficus* spp.) and Urticaceae (Mikich 2002; Passos et al. 2003; Passos and Gracioli 2004). The families Urticaceae and Moraceae have several pioneer species, such as *Cecropia pachystachya* and *Ficus dendrocida* (Lorenzi 1992; 1998).

According to Zortéa (2003), the three species of glossophagine bats captured in Guarulhos (Table 1) are the most common species of this subfamily and, in general, few species of this group have been found in sympatry (Passos et al. 2003; Nogueira et al. 2007). The largest number of *A. caudifer* (N= 47) in our study is a result of their netting while emerging from the shelter. A group of 12 *A. caudifer* was captured while emerging from the Candinhas House, where these individuals were harbored. Two *G. soricina* also emerged from that shelter. This nectar feeding bat has been usually found in Brazilian urban areas (Bredt and Uieda 1996; Silva et al. 1996; De Knegt et al. 2005; Uieda and Chaves 2005; Bredt et al. 2012), where they harbor in attics and basements. On the other hand, *A. caudifer* is more adapted to medium and large forest fragments (Reis et al. 2003), as is the case of regions surrounding the Serra do Itaberaba.

The phyllostomines *Mimon bennettii* and *Micronycteris megalotis* were very scarce in our samples (Table 1), and they were recorded only during the dry season (May, July and August). Just one individual of the former species was netted in Guarulhos (site D). Two individuals of *M. megalotis* were captured in two different Atlantic Forest fragments (sites C and G). These phyllostomines seem to be negatively affected by the process of forest fragmentation and are usually recorded in low numbers in fragmented landscapes (Reis et al. 2000; 2006; Dias et al. 2002).

Of the three species of vampire bat guild, two (*D. rotundus* and *D. ecaudata*) were captured in Serra do Itaberaba. The former, also known as the common vampire bat, feeds on the blood of cattle, horses, pigs, and even humans (Gardner 1977; Schneider et al. 2001; Gomes and Uieda 2004), whereas the second feeds on avian blood (Gardner 1977; Uieda et al. 1992). *D. rotundus* was netted on trails, at the outside of their shelters (a groundwater tunnel and an abandoned house), and around corrals. The hairy-legged vampire bat *D. ecaudata* was sampled in four of the sites (C, F, G and I) and was always netted near their food source (chickens) or their day roost. In one occasion, we observed two individuals of *D. ecaudata* while hanging on the chicken's tail and feeding on their cloacal region. The chickens were resting on the hennery perches at 0.5 m high (site I). This feeding behavior was previously cited




FIGURE 1. Map of the northeastern region of the municipality of Guarulhos, state of São Paulo, showing the Serra do Itaberaba and the location of the 10 sites (A to J) sampled for bats (see text for details).

by Greenhall (1988), Uieda (1992), and Uieda *et al.* (1992).

In one of the properties sampled (site G), there was a groundwater tunnel underneath a trail. This tunnel was at least 100 m long, had four openings, was full of water at several points, and its walls consisted of overlapping rocks. Seven species (*D. rotundus*, *C. perspicillata*, *G. soricina*, *A. caudifer*, *S. lilium*, *A. lituratus*, and *A. fimbriatus*) were captured in the net set in front of the tunnel's entrance. We did not check if these species were roosting inside the tunnel, but in the literature only *D. rotundus* and *C. perspicillata* were frequently observed harboring in this type of day roost (Marques 1985; Gomes and Uieda 2004; Oliveira *et al.* 2009).

An abandoned house (site I) was also used as shelter by small groups of *D. rotundus* (three individuals), *D. ecaudata* (three), *C. perspicillata* (seven), and *A. caudifer* (nine). The vampire bats harbored in only one of the three rooms of the house, where their guano was accumulated on the floor. *Carollia perspicillata* and *A. caudifer* were roosting in the opposite room. All four species have been observed cohabiting caves of the valley of the Rio Ribeira, SP (Trajano 1985), and the first three were observed in cohabitation in caves in the Federal District, Central-Western Brazil (Bredt *et al.* 1999). Marques (1985) had recorded *C. perspicillata* and *A. caudifer* cohabiting manholes in Manaus, AM, northern Brazil.

Vespertilionidae was represented by three species in our samples. *Myotis nigricans* was more abundant than *M. albescens* and was netted during its low flights along trails, above ponds, and built areas. *Myotis albescens*, in turn, was only captured during low flights above ponds (artificial lakes in areas B and J). *Myotis nigricans* is frequently listed in bat inventories from different regions (e.g. Peracchi and Albuquerque 1993; Teixeira and Peracchi 1996; Dias *et al.*

2002; Uieda and Chaves 2005; Reis *et al.* 2006; Ortêncio-Filho and Reis 2009), although in a low number. On the other hand, *M. albescens* is scarcely recorded in bat surveys and little is known about its biology. Only one juvenile bat of *Histiotus velatus* was found fallen on the floor of an old and uninhabited house (site I). According to Bianconi and Pedro (2007), *H. velatus* can use different kinds of shelter such as attics, roofs, crevices in rocks, caves, and abandoned mines.

Molossid bats usually harbor in house roofs, attics, and building expansion joints (Bredt and Uieda 1996, Chaves and Sodr e 2007, Quintela *et al.* 2011). Although some mist nets were set around houses and trees of an orchard, only one individual (not pregnant adult female) of *Molossops neglectus* was netted in the orchard. Except by the slightly smaller length of the forearm (34.6 mm) and the greatest length of the skull (15.1 mm), measurements of this specimen, including breadth of the braincase (8.3 mm), length of the maxillary toothrow (5.9 mm), and length of mandible (11.4 mm) are within the range described for the species (Gregorin and Taddei 2002; Gregorin *et al.* 2004; Bernardi *et al.* 2007). Diagnostic qualitative characters described by Gregorin *et al.* (2004) and Barquez *et al.* (1999; 2011), such as dark brown dorsal hairs with distinctly white bases, ears and membranes almost black, and separated and triangular ears were also observed in our specimen. *M. neglectus* was only recently recorded in Rio Grande do Sul (Bernardi *et al.* 2007), significantly expanding its known geographic distribution. The present study is the first to report this rare bat for Guarulhos, indicating that new species of bats can still be found at this municipality. Future works using different sampling methods should also improve our knowledge about the bat fauna of Guarulhos.

TABLE 1. Bat species captured in the municipality of Guarulhos, state of S o Paulo, in 2011.

TAXON	NUMBER OF CAPTURES											TOTAL (%)	
	A	B	C	D	E	F	G	H	I	J			
Family Phyllostomidae													
<i>Mimon bennettii</i> (Gray, 1838)				1									1 (0.3)
<i>Micronycteris megalotis</i> (Gray, 1842)			1				1						2 (0.6)
<i>Artibeus fimbriatus</i> Gray, 1838	4	3	5	7	8	21	16	2	1	2			69 (19.8)
<i>Artibeus lituratus</i> (Olfers, 1818)	5		4	9	7	3	26	1	4				59 (17.0)
<i>Pygoderma bilabiatum</i> (Wagner, 1843)	1			1					1				3 (0.9)
<i>Platyrrhinus lineatus</i> (�. Geoffroy, 1810)					1	1							2 (0.6)
<i>Sturnira lilium</i> (�. Geoffroy, 1810)	11	2	3	6	7	7	24	5	8				73 (21.0)
<i>Carollia perspicillata</i> (Linnaeus, 1758)	3			1	7	3	7	2	9				32 (9.2)
<i>Glossophaga soricina</i> (Pallas, 1766)	1	1		1	4		9						16 (4.6)
<i>Anoura caudifer</i> (�. Geoffroy, 1818)			2		25	1	4		15				47 (13.5)
<i>Anoura geoffroyi</i> Gray, 1838	1					2	1						4 (1.1)
<i>Desmodus rotundus</i> (�. Geoffroy, 1810)	2						3		9				14 (4.0)
<i>Diphylla ecaudata</i> Spix, 1823			2			2	1		7				12 (3.4)
Family Vespertilionidae													
<i>Myotis albescens</i> (�. Geoffroy, 1806)		1								4			5 (1.4)
<i>Myotis nigricans</i> (Schinz, 1821)		1	1		1	1		1	1	1			7 (2.0)
<i>Histiotus velatus</i> (I. Geoffroy, 1824)									1				1 (0.3)
Family Molossidae													
<i>Molossops neglectus</i> Williams and Genoways, 1980					1								1 (0.3)
Total Species/area	8	5	7	7	9	9	10	5	10	3			348 (100)

ACKNOWLEDGMENTS: We are grateful to Universidade Guarulhos for providing financial support (Project of Professor Research, Round I, 2010-2011) and to Gabriela Uieda for improving the English version of the manuscript. We thank Alexandre Kise from Secretaria do Meio Ambiente (SEMA) of Guarulhos for logistical support, all owners for permitting the bat samplings in their properties and João Carlos Campos, Jorge Nascimento, Valéria de Moraes and Samuel G.C. Mariano for helping in the field work. We thank Isaac Passos de Lima, Marcelo R. Nogueira and one anonymous referee, for reviews and comments on the manuscript. We also thank Dr. Antônio Roberto Saad and William de Queiroz, from Geoprocessing Laboratory of Universidade Guarulhos, for making the map of study area.

LITERATURE CITED

- Andrade, M.R.M., A.M.S. Oliveira, W. Queiroz, S.E. Sato, E.J. Barros, G. Bagattini and A.A. Aleixo. 2008. Aspectos fisiográficos da paisagem guarulhense; p. 25-37 In Omar, E.E.H. (org.). *Guarulhos tem história: questões sobre história natural, social e cultural*. São Paulo: Ananda Gráfica e Editora. 200p.
- Araújo, P. and A. Langguth. 2010. Caracteres distintivos das quatro espécies de grandes *Artibeus* (Phyllostomidae) de Paraíba e Pernambuco, Brasil. *Chiroptera Neotropical* 16(2): 715-722.
- Barquez, R.M., M.A. Mares and J.K. Braun. 1999. *The Bats of Argentina. Special Publications of the Museum of Texas Tech University* 42: 1-275.
- Barquez, R.M., A.V. Franzoy and M.M. Díaz. 2011. Mammalia, Chiroptera, Molossidae *Molossops neglectus* Williams and Genoways, 1980: Range extension and first record for Corrientes Province, Argentina. *Check List* 7(6): 889-890.
- Bernardi, I.P., A. Pulchério-Leite, J.M.D. Miranda and F.C. Passos. 2007. Ampliação da distribuição de *Molossops neglectus* Williams & Genoways (Chiroptera, Molossidae) para o Sul da América do Sul. *Revista Brasileira de Zoologia* 24(2): 505-507.
- Bianconi, G.V. and W.A. Pedro. 2007. Família Vespertilionidae; p.167-195 In N.R. Reis, A.L. Peracchi, W.A. Pedro and I.P. Lima (ed.). *Morcegos do Brasil*. Londrina: Nélio R. dos Reis.
- Bianconi, G.V., S.B. Mikich and W.A. Pedro. 2004. Diversidade de morcegos (Mammalia, Chiroptera) em remanescentes florestais do município de Fênix, noroeste do Paraná, Brasil. *Revista Brasileira de Zoologia* 21(4): 943-954.
- Bordignon, M.O. 2006. Diversidade de morcegos (Mammalia, Chiroptera) do Complexo Aporé-Sucuriú, Mato Grosso do Sul, Brasil. *Revista Brasileira de Zoologia* 23(4): 1002-1009.
- Bredt, A. and W. Uieda. 1996. Bats from urban and rural environments of the Distrito Federal, Mid-western Brazil. *Chiroptera Neotropical* 2(2): 54-57.
- Bredt, A., W. Uieda and E.D. Magalhães. 1999. Morcegos cavernícolas da região do Distrito Federal, centro-oeste do Brasil (Mammalia, Chiroptera). *Revista Brasileira de Zoologia* 16(3): 731-770.
- Bredt, A., W. Uieda and W. A. Pedro. 2012. *Plantas e morcegos: na recuperação de áreas degradadas e na paisagem urbana*. Brasília: Rede de Sementes do Cerrado. 273p.
- Chaves, M.E. and M.M. Sodrê. 2007. Manejo de morcegos em áreas urbanas; p. 139-144 In A.S. Pinto, M.M. Rossi and E. Salmeron (org.). *Manejo de Pragas Urbanas*. Piracicaba: CP 2.
- De Knecht, L.V., J.A. Silva, E.C. Moreira and G.L. Sales. 2005. Morcegos capturados no município de Belo Horizonte, 1999-2003. *Arquivo Brasileiro de Medicina Veterinária e Zootecnia* 57(5): 576-583.
- Diário Oficial do Município de Guarulhos No. 042/2007. *Portaria nº018/2007-SM*. Electronic Database accessible at <http://novo.guarulhos.sp.gov.br/>. Captured on 11 November 2010.
- Dias, D., A.L. Peracchi and S.S.P. Silva. 2002. Quirópteros do Parque Estadual da Pedra Branca, Rio de Janeiro, Brasil (Mammalia, Chiroptera). *Revista Brasileira de Zoologia* 19(Supl.2): 113-140.
- Dias, D. and A.L. Peracchi. 2008. Quirópteros da Reserva Biológica do Tinguá, estado do Rio de Janeiro, sudeste do Brasil (Mammalia: Chiroptera). *Revista Brasileira de Zoologia* 25(2): 333-369.
- Falcão, F.C., V.F. Rebêlo and S.A. Talamoni. 2003. Structure of a bat assemblage (Mammalia, Chiroptera) in Serra do Caraça Reserve, South-east Brazil. *Revista Brasileira de Zoologia* 20(2): 347-350.
- Fleming, T. H. 1988. *The short-tailed fruit bat: a study in plant-animal interactions*. Chicago, Illinois: University of Chicago Press. 365p.
- Gardner, A.L. 1977. Feeding Habits; p. 293-350 In R. J. Baker, J.K. Jones Jr. and D.C. Carter (ed.) *Biology of bats of the New World Family Phyllostomatidae, part II. Special Publications of Museum Texas Technical University* 13: 1-364.
- Gomes, M.N. and W. Uieda. 2004. Abrigos diurnos, composição de colônias, dimorfismo sexual e reprodução do morcego hematófago *Desmodus rotundus* (E. Geoffroy) (Chiroptera, Phyllostomidae) no Estado de São Paulo, Brasil. *Revista Brasileira de Zoologia* 21(3): 629-638.
- Greenhall, A.M. 1988. Feeding behavior. p.111-131 In A.M. Greenhall and W. Schmidt (eds). *Natural History of Vampire Bats*. Boca Raton, CRC Press, 246p.
- Gregorin, R., B.K. Lim, W.A. Pedro, F.C. Passos and V.A. Taddei. 2004. Distributional extension of *Molossops neglectus* (Chiroptera, Molossidae) into southeastern Brazil. *Mammalia* 68(2-3): 233-237.
- Gregorin, R. and V.A. Taddei. 2002. Chave artificial para a identificação de Molossídeos brasileiros (Mammalia, Chiroptera). *Mastozoologia Neotropical* 9(1):13-32.
- Kalko, E. K. V. 1998. Organization and diversity of tropical bat communities through space and time. *Zoology* 101: 281-297.
- LaVal, R.K. 1973. A revision of the neotropical bats of the genus *Myotis*. *Science Bulletin Natural History Museum Los Angeles County* 15: 1-53.
- Lorenzi, H. 1992. *Árvores Brasileiras: manual de identificação e cultivo de plantas arbóreas nativas do Brasil*. São Paulo: Editora Plantarum. 352p.
- Lorenzi, H. 1998. *Árvores Brasileiras: manual de identificação e cultivo de plantas arbóreas nativas do Brasil*. Volume II. São Paulo: Editora Plantarum. 352p.
- Marques, S.A. 1985. Espécies associadas e algumas características físicas influenciando na presença de *Carollia perspicillata* em bueiros na região de Manaus, AM (Mammalia, Chiroptera: Phyllostomidae). *Acta Amazônica* 15(1-2): 243-248.
- Mikich, S.B. 2002. A dieta dos morcegos frugívoros (Mammalia, Chiroptera, Phyllostomidae) de um pequeno remanescente de Floresta Estacional Semidecidual do Sul do Brasil. *Revista Brasileira de Zoologia* 19(1): 239-249.
- Nogueira, M.R., D. Dias and A.L. Peracchi. 2007. Subfamília Glossophaginae; p.45-59 In N.R. Reis, A.L. Peracchi, W.A. Pedro and I.P. Lima (eds.). *Morcegos do Brasil*. Londrina: Nélio R. dos Reis, 253p.
- Oliveira, A.M.S. and M.R.M. Andrade. 2008. Expansão urbana e problemas geoambientais do uso do solo em Guarulhos; p. 47-55 In Omar, E.E.H. (org.). *Guarulhos tem história: questões sobre história natural, social e cultural*. São Paulo: Ananda Gráfica e Editora. 200p.
- Oliveria, P.R., D.A.R. Silva, J.H. Rocha, S.M.A. Melo, N.G. Bombonato and F.O. Carneiro e Silva. 2009. Levantamento, cadastramento e estimativa populacional das habitações de morcegos hematófagos, antes e após atividades de controle, no município de Araguari, MG. *Arquivos do Instituto Biológico*, 76(4): 553-560.
- Ortêncio-Filho, H. and N.R. Reis. 2009. Species richness and abundance of bats in fragments of the seasonal semidecidual forest, Upper Paraná River, southern Brazil. *Brazilian Journal of Biology* 69(Suppl. 2): 727-734.
- Passos, F.C., W.R. Silva, W.A. Pedro and M. R. Bonin. 2003. Frugivoria em morcegos (Mammalia, Chiroptera) no Parque Estadual Intervalles, sudeste do Brasil. *Revista Brasileira de Zoologia* 20(3): 511-517.
- Passos, F.C. and G. Gracioli. 2004. Observações da dieta de *Artibeus lituratus* (Olfers) (Chiroptera, Phyllostomidae) em duas áreas do sul do Brasil. *Revista Brasileira de Zoologia* 21(3): 487-489.
- Peracchi, A.L. and S.T. Albuquerque. 1993. Quirópteros do município de Linhares, Estado do Espírito Santo, Brasil (Mammalia, Chiroptera). *Revista Brasileira de Biologia* 53(4): 575-581.
- Quintela, F.M., C. Ibarra, S.V. Oliveira, I.G. Medvedovsky, F. Correa, D. Gianuca, A. Gava and S.M. Pacheco. 2011. Mammalia, Chiroptera, Rio Grande, state of Rio Grande do Sul, Brazil. *Check List* 7(4): 443-447.
- Reis, N.R., M.L.S. Barbieri, I.P. Lima and A.L. Peracchi. 2003. O que é melhor para manter a riqueza de espécies de morcegos (Mammalia, Chiroptera): um fragmento florestal grande ou vários fragmentos de pequeno tamanho? *Revista Brasileira de Zoologia* 20(2): 225-230.
- Reis, N.R., A.L. Peracchi, I.P. Lima and W.A. Pedro. 2006. Riqueza de espécies de morcegos (Mammalia, Chiroptera) em dois diferentes habitats, na região centro-sul do Paraná, sul do Brasil. *Revista Brasileira de Zoologia* 23(3): 813-816.
- Reis, N.R., A.L. Peracchi, M.L. Sekiama and I.P. Lima. 2000. Diversidade de morcegos (Chiroptera, Mammalia) em fragmentos florestais no Estado do Paraná, Brasil. *Revista Brasileira de Zoologia* 17(3): 697-704.
- Schneider, M.A., J. Aron, C. Santos-Burgoa, W. Uieda and S. Ruiz-Velazco. 2001. Common vampire bat attacks on humans in a village of the Amazon region of Brazil. *Cadernos de Saúde Pública* 17(6): 1531-1536.
- Silva, M.M.S., N.M.S. Harmani, E.F.B. Gonçalves and W. Uieda. 1996. Bats from the Metropolitan region of São Paulo, Southeastern Brazil. *Chiroptera Neotropical* 2(1): 39-41.
- Straube, F.C. and G.V. Bianconi. 2002. Sobre a grandeza e a unidade utilizada para estimar esforço de captura com utilização de redes-de-neblina. *Chiroptera Neotropical* 8:150-152.
- Taddei, V.A., C.A. Nobile and E. Morielle-Versute. 1998. Distribuição geográfica e análise morfométrica comparativa em *Artibeus obscurus* (Schinz, 1821) e *Artibeus fimbriatus* Gray, 1838 (Mammalia, Chiroptera, Phyllostomidae). *Ensaios e Ciência* 2(2): 71-127.
- Trajano, E. 1985. Ecologia de populações de morcegos cavernícolas em uma região cárstica do Sudeste do Brasil. *Revista Brasileira de Zoologia* 2(5): 255-320.
- Teixeira, S. C. and A.L. Peracchi. 1996. Morcegos do Parque Estadual da

- Serra da Tiririca, Rio de Janeiro, Brasil (Mammalia, Chiroptera). *Revista Brasileira de Zoologia* 13(1): 61-68.
- Uieda, W. 1992. Período de atividade alimentar e tipos de presa dos morcegos hematófagos (Phyllostomidae) no Sudeste do Brasil. *Revista Brasileira de Biologia* 52(4): 563-573.
- Uieda, W. and M.E. Chaves. 2005. Bats from Botucatu region, State of São Paulo, Southeastern Brazil. *Chiroptera Neotropical* 11(1-2): 224-226.
- Uieda, W., S. Buck and I. Sazima. 1992. Feeding behavior of the vampire bats *Diaemus youngi* and *Diphylla ecaudata* on smaller birds in captivity. *Ciência e Cultura* 44(6): 410-412.
- Vizotto, L.D. and V.A. Taddei. 1973. Chave para determinação de quirópteros brasileiros. *Boletim de Ciências, Faculdade de Filosofia, Ciências e Letras* 1:1-72.
- Zanon, C.M.V and N.R. Reis. 2007. Bats (Mammalia, Chiroptera) in the Ponta Grossa region, Campos Gerais, Paraná, Brazil. *Revista Brasileira de Zoologia* 24(2): 327-332.
- Zortéa, M. 2003. Reproductive patterns and feeding habits of three nectarivorous bats (Phyllostomidae: Glossophaginae) from the Brazilian Cerrado. *Brazilian Journal of Biology* 63(1): 159-168.
- Zortéa, M. 2007. Subfamília Stenodermatinae; p.107-128 In N.R. Reis, A.L. Peracchi, W.A. Pedro and I.P. Lima (ed.). *Morcegos do Brasil*. Londrina: Nélío R. dos Reis.

RECEIVED: April 2012
ACCEPTED: August 2012
PUBLISHED ONLINE: November 2012
EDITORIAL RESPONSIBILITY: Marcelo R. Nogueira

APPENDIX 1. Voucher list.

Mimon bennettii: (1475); *Micronycteris megalotis*: (1488); *Artibeus fimbriatus*: (1471, 1476, 1480, 1487); *Artibeus lituratus*: (1477, 1489, 1490); *Pygoderma bilabiatum*: (1467); *Platyrrhinus lineatus*: (1494); *Sturnira lilium*: (1465, 1469, 1473, 1479); *Carollia perspicillata*: (1464, 1474, 1481, 1493); *Glossophaga soricina*: (1468, 1486, 1497); *Anoura caudifer*: (1478, 1484, 1498); *Anoura geoffroyi*: (1483, 1496); *Desmodus rotundus*: (1466, 1485); *Diphylla ecaudata*: (1482, 1491); *Myotis albescens*: (1472); *Myotis nigricans*: (1470, 1495); *Molossops neglectus*: (1492).