

# Towards a Realization of the Condensed-Matter–Gravity Correspondence in String Theory via Consistent Abelian Truncation of the Aharony-Bergman-Jafferis-Maldacena Model

Asadig Mohammed,<sup>1,\*</sup> Jeff Murugan,<sup>1,†</sup> and Horatiu Nastase<sup>2,‡</sup>

<sup>1</sup>*The Laboratory for Quantum Gravity and Strings, Department of Mathematics and Applied Mathematics, University of Cape Town, Private Bag, Rondebosch 7700, South Africa*

<sup>2</sup>*Instituto de Física Teórica, UNESP-Universidade Estadual Paulista, Rua Dr. Bento T. Ferraz 271, Bloco II, São Paulo 01140-070, São Paulo, Brazil*

(Received 12 June 2012; revised manuscript received 21 August 2012; published 31 October 2012)

We present an embedding of the three-dimensional relativistic Landau-Ginzburg model for condensed matter systems in an  $\mathcal{N} = 6$ ,  $U(N) \times U(N)$  Chern-Simons-matter theory [the Aharony-Bergman-Jafferis-Maldacena model] by consistently truncating the latter to an Abelian effective field theory encoding the collective dynamics of  $\mathcal{O}(N)$  of the  $\mathcal{O}(N^2)$  modes. In fact, depending on the vacuum expectation value on one of the Aharony-Bergman-Jafferis-Maldacena scalars, a mass deformation parameter  $\mu$  and the Chern-Simons level number  $k$ , our Abelianization prescription allows us to interpolate between the Abelian Higgs model with its usual multivortex solutions and a  $\phi^4$  theory. We sketch a simple condensed matter model that reproduces all the salient features of the Abelianization. In this context, the Abelianization can be interpreted as giving a dimensional reduction from four dimensions.

DOI: [10.1103/PhysRevLett.109.181601](https://doi.org/10.1103/PhysRevLett.109.181601)

PACS numbers: 11.25.Tq, 11.15.Yc, 74.40.Kb

**Introduction.**—The gauge-gravity duality, as manifest in the AdS/CFT correspondence [1] has evolved from its humble origins in string theory into one of the most powerful tools in the arsenal of physicists studying nonperturbative and strong-coupling phenomena in quantum field theories today. The original and, arguably, most studied of these *applied string theory* phenomena was the physics of QCD. In particular, a lot of recent work has focused on the quark-gluon plasma observed in heavy ion colliders such as the Relativistic Heavy Ion Collider (RHIC) and the ALICE experiment at CERN (see for example, Ref. [2] for a recent review). More recently though, the ideas of holography have found a new, lower-dimensional hunting ground in condensed matter physics. The  $pp$ -wave or Berenstein-Maldacena-Nastase limit of AdS/CFT selects operators with large charge [3] in  $\mathcal{N} = 4$  super Yang-Mills theory. The physics of such large operators is, in a very concrete sense, isomorphic to the physics of certain spin chains [4]; a realization that has led, not only to an enormous development in our understanding of integrability in string theory but also served as a forerunner to many of the developments in what has now become known as the anti-de Sitter condensed matter theory (*AdS/CMT*) correspondence.

In these cases however, the duality was heuristically motivated by a relation between some system of branes and a gravitational background, in some decoupling limit. Later, it was realized that if physics in AdS is always holographic, we can consider simple theories in AdS, that should be related to some strongly coupled conformal field theory on the boundary. Thus naturally AdS/CFT came to be applied to condensed matter systems, where one encounters strongly coupled conformal field theories that cannot be dealt with in other ways (see for instance the reviews

Refs. [5,6] for an introduction and relevant references). In all these cases, however, the argument is mostly one of universality, that a variety of (large  $N$ ) theories have some small set of (Abelian) operators dual to some finite and small number of fields in AdS, usually a  $U(1)$  gauge field, a scalar and maybe some spinors, representing a (sometimes consistent) truncation of some AdS/CFT pair. In other words, either one truncates the number of operators of the system, in which case it is not entirely clear (a) why one should focus on a subset, or how one understands from the point of view of a condensed matter system the focus on the few operators of the large  $N$  system, or (b) one thinks of an Abelian condensed matter *analog* of the large  $N$  theory, in which case it is not clear why we can use just a gravity dual, as opposed to a full string theory. In either case, we find the argument less than persuasive.

In this Letter, we will take some steps towards a better understanding of AdS/CMT, by proposing a modification of the above set-up. We consider instead a consistent truncation of the three-dimensional Aharony-Bergman-Jafferis-Maldacena (ABJM) theory (which has a known gravity dual), a truncation that corresponds to the collective dynamics of  $\mathcal{O}(N)$  fields out of the  $\mathcal{O}(N^2)$  of ABJM theory, and gives an effective theory that is easily identified as the relativistic Landau-Ginzburg model. We also sketch a simple CMT model that has the same qualitative features as the ABJM Abelianization, allowing us to understand better in what sense we can use the ABJM theory for condensed matter systems. Here we present only the main ideas, leaving the technical details to a longer paper [7].

As a point of clarity, we note that the idea of a consistent truncation in string theory is not a new one, having been featured before in two primary contexts. On the gravity side

of the AdS/CFT correspondence, when one is interested in a classical limit only, a consistent truncation means that we can safely drop the *nonzero modes*, as these will only appear in quantum loops. In supergravity compactifications, however, a consistent truncation for a dimensional reduction means that we can drop all the nonzero (KK) modes from the low energy quantum theory as well, provided that the coupling to the nonzero modes can be made arbitrarily small, or that the masses of the nonzero modes are much larger than the mass parameters of the low energy theory. Hence quantum loops of these nonzero modes may be ignored. We will argue below that it is this latter case that arises here. To understand the collective dynamics that is crucial to our argument, consider a large number,  $N$ , of branes in some background. A classical solution that is obtained by turning

on fields in all the  $N$  branes will curve the background space nontrivially corresponding, via AdS/CFT, to some finite deformation of the dual theory. On the other hand, just turning on fields on a single brane will produce a negligible deformation that will not deform the background space nor the dual. In our case it is the former situation that arises so that in this sense, the collective dynamics of  $\mathcal{O}(N)$  fields really is different from the dynamics of a single field, and it is this that allows for a dual gravitational interpretation.

*ABJM model and its massive deformation.*—The ABJM model [8] is an  $\mathcal{N} = 6$  supersymmetric  $U(N) \times U(N)$  Chern-Simons gauge theory at level  $(k, -k)$ , with bifundamental scalars  $C^I$  and fermions  $\psi_I$ ,  $I = 1, \dots, 4$  in the fundamental of the  $SU(4)_R$  symmetry group and gauge fields for the two groups  $A_\mu$  and  $\hat{A}_\mu$ . Its action is given by

$$S = \int d^3x \left[ \frac{k}{4\pi} \epsilon^{\mu\nu\lambda} \text{Tr} \left( A_\mu \partial_\nu A_\lambda + \frac{2i}{3} A_\mu A_\nu A_\lambda - \hat{A}_\mu \partial_\nu \hat{A}_\lambda - \frac{2i}{3} \hat{A}_\mu \hat{A}_\nu \hat{A}_\lambda \right) - \text{Tr} D_\mu C_I^\dagger D^\mu C^I - i \text{Tr} \psi^{I\dagger} \gamma^\mu D_\mu \psi_I \right. \\ \left. + \frac{4\pi^2}{3k^2} \text{Tr} (C^I C_I^\dagger C^J C_J^\dagger C^K C_K^\dagger + C_I^\dagger C^I C_J^\dagger C^J C_K^\dagger C^K + 4C^I C_J^\dagger C^K C_I^\dagger C^J C_K^\dagger - 6C^I C_J^\dagger C^J C_I^\dagger C^K C_K^\dagger) \right. \\ \left. + \frac{2\pi i}{k} \text{Tr} (C_I^\dagger C^I \psi^{J\dagger} \psi_J - \psi^{J\dagger} C^I C_I^\dagger \psi_J - 2C_I^\dagger C^J \psi^{I\dagger} \psi_J + 2\psi^{J\dagger} C^I C_J^\dagger \psi_J + \epsilon^{IJKL} C_I^\dagger \psi_J C_K^\dagger \psi_L \right. \\ \left. - \epsilon_{IJKL} C^I \psi^{J\dagger} C^K \psi^{L\dagger} \right), \quad (1)$$

where the gauge covariant derivative is

$$D_\mu C^I = \partial_\mu C^I + iA_\mu C^I - iC_I \hat{A}_\mu. \quad (2)$$

The action has an  $SU(4) \times U(1)$   $R$  symmetry associated with the  $\mathcal{N} = 6$  supersymmetries. It admits a maximally supersymmetric (i.e., preserving all  $\mathcal{N} = 6$ ) massive deformation with mass parameter  $\mu$  [9,10], which breaks the  $R$  symmetry down to  $SU(2) \times SU(2) \times U(1)_A \times U(1)_B \times \mathbb{Z}_2$  by splitting the scalars as

$$C^I = (Q^\alpha, R^\alpha); \quad \alpha = 1, 2. \quad (3)$$

The mass deformation changes the potential to

$$V = \text{Tr}(|M^\alpha|^2 + |N^\alpha|^2), \quad (4)$$

where

$$M^\alpha = \mu Q^\alpha + \frac{2\pi}{k} (2Q^{[\alpha} Q_\beta^\dagger Q^{\beta]} + R^\beta R_\beta^\dagger Q^\alpha - Q^\alpha R_\beta^\dagger R^\beta \\ + 2Q^\beta R_\beta^\dagger R^\alpha - 2R^\alpha R_\beta^\dagger Q^\beta), \\ N^\alpha = -\mu R^\alpha + \frac{2\pi}{k} (2R^{[\alpha} R_\beta^\dagger R^{\beta]} + Q^\beta Q_\beta^\dagger R^\alpha - R^\alpha Q_\beta^\dagger Q^\beta \\ + 2R^\beta Q_\beta^\dagger Q^\alpha - 2Q^\alpha Q_\beta^\dagger R^\beta). \quad (5)$$

This mass-deformed Aharony-Bergman-Jafferis-Maldacena theory has ground states of the *fuzzy sphere* type given by [9,10]

$$R^\alpha = cG^\alpha; \quad Q^\alpha = 0 \quad \text{and} \quad Q_\alpha^\dagger = cG^\alpha; \quad R^\alpha = 0, \quad (6)$$

where  $c \equiv \sqrt{\frac{\mu k}{2\pi}}$  and the matrices  $G^\alpha$ ,  $\alpha = 1, 2$ , bifundamental under  $U(N) \times U(N)$ , satisfy (with no summation on repeated indices)

$$G^\alpha = G^\alpha G_\beta^\dagger G^\beta - G^\beta G_\beta^\dagger G^\alpha. \quad (7)$$

This ground state corresponds to a fuzzy two-sphere [11,12]. An explicit solution for  $G^\alpha$  is given by

$$(G^1)_{m,n} = \sqrt{m-1} \delta_{m,n}, \quad (G^2)_{m,n} = \sqrt{(N-m)} \delta_{m+1,n} \\ (G_1^\dagger)_{m,n} = \sqrt{m-1} \delta_{m,n}, \quad (G_2^\dagger)_{m,n} = \sqrt{(N-n)} \delta_{n+1,m}. \quad (8)$$

We will now use these so-called Gomis-Rodriguez-Gomez-van Raamsdonk-Verlinde matrices to posit an ansatz that effectively Abelianizes the ABJM model while retaining the large  $N$  limit.

*Consistent Abelian truncation and condensed matter model.*—Consider then the following ansatz for the Chern-Simons fields and the scalar matter in the supermultiplet:

$$\begin{aligned}
A_\mu &= a_\mu^{(2)} G^1 G_1^\dagger + a_\mu^{(1)} G^2 G_2^\dagger \\
\hat{A}_\mu &= a_\mu^{(2)} G_1^\dagger G^1 + a_\mu^{(1)} G_2^\dagger G^2 \\
Q^\alpha &= \phi_\alpha G^\alpha \\
R^\alpha &= \chi_\alpha G^\alpha,
\end{aligned} \tag{9}$$

where, again, there is no summation over the repeated  $\alpha$ ;  $a_\mu^{(1)}$  and  $a_\mu^{(2)}$  are real-valued vector fields and  $\phi_\alpha, \chi_\alpha$  are complex-valued scalar fields. This provides a consistent truncation of the ABJM action to

$$\begin{aligned}
S = & -\frac{N(N-1)}{2} \int d^3x \left[ \frac{k}{4\pi} \epsilon^{\mu\nu\lambda} (a_\mu^{(2)} f_{\nu\lambda}^{(1)} + a_\mu^{(1)} f_{\nu\lambda}^{(2)}) \right. \\
& \left. + |D_\mu \phi_i|^2 + |D_\mu \chi_i|^2 + U(|\phi_i|, |\chi_i|) \right], \tag{10}
\end{aligned}$$

where  $U \equiv 2V/N(N-1)$  is a rescaling of the potential

$$\begin{aligned}
V = & \frac{2\pi^2}{k^2} N(N-1) [ (|\phi_1|^2 + |\chi_1|^2)(|\chi_2|^2 - |\phi_2|^2 - c^2)^2 \\
& + (|\phi_2|^2 + |\chi_2|^2)(|\chi_1|^2 - |\phi_1|^2 - c^2)^2 \\
& + 4|\phi_1|^2 |\phi_2|^2 (|\chi_1|^2 + |\chi_2|^2) \\
& + 4|\chi_1|^2 |\chi_2|^2 (|\phi_1|^2 + |\phi_2|^2) ], \tag{11}
\end{aligned}$$

and where the, now Abelian, gauge covariant derivatives are  $D_\mu \phi_i = (\partial_\mu - ia_\mu^{(i)})\phi_i$  and  $D_\mu \chi_i = (\partial_\mu - ia_\mu^{(i)})\chi_i$ . Different choices of scalars turned on lead to different consistent truncations we collect below:

$\chi_2 = \phi_2 = 0$ : This leads to a model with two massive complex scalars with no self-interactions. This is essentially trivial and will not merit further attention.

$\chi_1 = \phi_2 = 0$ : After a minor relabeling of  $\chi_2 \rightarrow \phi_2$  this choice produces

$$\begin{aligned}
S = & -\frac{N(N-1)}{2} \int d^3x \left[ \frac{k}{4\pi} \epsilon^{\mu\nu\lambda} (a_\mu^{(2)} f_{\nu\lambda}^{(1)} + a_\mu^{(1)} f_{\nu\lambda}^{(2)}) \right. \\
& \left. + |D_\mu \phi_i|^2 + U(|\phi_i|) \right], \\
V = & \frac{2\pi^2 N(N-1)}{k^2} \left[ |\phi_1|^2 (|\phi_2|^2 - c^2)^2 + |\phi_2|^2 (|\phi_1|^2 + c^2)^2 \right] \tag{12}
\end{aligned}$$

a model which has vortex solutions with  $\phi_1 = |\phi_1| e^{iN_1\alpha}$  and  $\phi_2 = |\phi_2| e^{iN_2\alpha}$ , where  $\alpha$  is the polar angle on the plane, and  $|\phi_{1,2}|$  go to zero at  $r = 0$  and  $r = \infty$ .

$\phi_1 = \phi_2 = 0$ : If we also set  $\chi_1 = b = \text{const}$ , and solve for the (now auxiliary) gauge field  $a_\mu^{(1)}$  we obtain the action

$$S = -\frac{N(N-1)}{2} \int d^3x \left[ \frac{k^2}{8\pi^2 |b|^2} (f_{\mu\nu}^{(2)})^2 + |D_\mu \chi_2|^2 + V \right], \tag{13}$$

where the potential is

$$V = \frac{2\pi^2}{k^2} N(N-1) \{ |b|^2 |\chi_2|^4 + |\chi_2|^2 [(|b|^2 - c^2)^2 - 2c^2 |b|^2] + c^4 |b|^2 \}. \tag{14}$$

This is the most interesting case. Indeed, we see that for  $||b|^2 - c^2| > \sqrt{2}|c||b|$ , we obtain a regular  $\phi^4$  phase, whereas for  $||b|^2 - c^2| < \sqrt{2}|c||b|$  we obtain the Abelian-Higgs phase; i.e., this is a relativistic Landau-Ginzburg theory, with  $(|b|^2 - c^2)^2 \sim g$  and  $2c|b|^2 \sim g_c$ . However, in order to have a consistent truncation, in the above action we need to also satisfy the *equation of motion* for the constant  $|b|^2$ . This is indeed the case for Bogomolnyi-Prasad-Sommerfeld (BPS) solutions of the Abelian-Higgs action.

When canonically normalizing all the fields, the quartic coupling for the canonical scalar  $\tilde{\chi}_2$  from (14) becomes  $g^2$ , with  $g = 2\pi|b|/(Nk)$ , and the coefficient of the middle term in (14) becomes  $N^2 g^2/2$ . Then, for  $|b| \sim c$ , with  $k \sim 1$  and  $N$  large,  $g^2 \sim \mu/(N^2 k) \sim \mu/N^2 \ll \mu$ , and generically the mass of  $|\phi|$  is  $\sim \mu$ . But we can tune the system to be near zero mass, so that (for  $m^2 \neq 0$ ) we have  $|m^2| \ll \mu^2$ . Generic modes of the ABJM model (the *nonzero modes* dropped in our consistent truncation) have mass  $\mu$ , as easily checked in (1) and (5). Therefore we can drop the nonzero modes in the reduced low energy theory even at the quantum level, as advertised in the introduction, and consistently truncate to (13). Note also that substituting the reduction ansatz into the ABJM action, we find that the (sextic) potential gives a term with bilinear coupling to the nonzero modes  $\delta\phi$  of the type  $(\tilde{\chi}_2)^4 (\delta\phi)^2 / (k^2 N^2) \propto 1/N^2$ , the 2-fermion-2-scalar term gives a bilinear coupling  $(\tilde{\chi}_2)^2 \delta\bar{\psi} \delta\psi / (kN) \propto 1/N$ , and mass deformation quartic in the scalars gives a bilinear coupling  $(\tilde{\chi}_2)^2 (\delta\phi)^2 \mu / (kN) \propto \mu/N$ , which is  $\ll \mu$ , though still  $\gg \mu/N^2 \sim g^2$ .

Now also note that for  $|b| = c$ , the potential (14) has the vacuum  $|\phi| = |b| = c$ , which is nothing but the fuzzy sphere vacuum of the massive ABJM model; therefore, classical solutions of the reduced theory (13) are some type of deformations of the fuzzy sphere. Other examples of such classical solutions will be given elsewhere [7]. These solutions, giving a collective dynamics of  $\mathcal{O}(N)$  modes, then correspond to finite deformations of the gravity dual, unlike any solutions obtained by turning on a single mode. Therefore we retain the good features of the large  $N$  behavior (classical gravity dual) with this Abelian reduction, as advertised in the introduction.

At this point, one might ask whether the modes in the Landau-Ginzburg action we consider are the only light ones and if not, do they couple to any others? For the extra modes in (10) that we dropped, the answer is simple. All canonically normalized fields couple to the Landau-Ginzburg (LG) mode with coupling  $\sim \mu/(N^2 k) \sim g^2$ . Moreover, all  $\phi_i, \chi_i$  have an explicit mass term of the order  $N^2 c^4 / k^2 \sim \mu^2$  and there is also a contribution from the vacuum expectation value (VEV)  $\chi_1 = b$  of

$1/k^2[-4c^2b^2|\tilde{\chi}_2|^2 + b^4(|\tilde{\chi}_2|^2 + |\tilde{\phi}_2|^2)]$ . Consequently in the region of parameter space where our LG mode  $\tilde{\chi}_2$  can be tuned to be light all the other modes *stay heavy*. For generic modes outside the action (10) the answer is a bit more difficult. As we have already argued, since generic mass terms are of the order  $m^2 \sim \mu^2 > 0$ , the only thing remaining to check is if they can be (almost) cancelled by the terms coming from the Higgs VEV  $\chi_1 = b$ . We can obtain the total mass term by keeping only two  $C^I$ 's in the potential (4), and replacing the rest by  $C^I = (R^1 = bG^1, R^2 = Q^1 = Q^2 = 0)$ . Setting this to zero produces a very long equation for the trace of products of two  $C^I$  matrices and up to four  $G^1$  matrices being zero, which we will not reproduce here. One solution is given by our LG light mode,  $C^I = (R^2 = \chi_2 G^2, R^1 = Q^1 = Q^2 = 0)$ , and amounts to an identity between the  $G^1$  and  $G^2$  matrices (together with the condition  $|b|^2 = c^2(2 \pm \sqrt{3})$  for a massless LG field). The question is whether the solution is unique. While we don't know of a general *mathematical* proof of uniqueness, *physically* it is clear that there can't be another solution. Indeed, this solution is related to the existence of the maximally supersymmetric fuzzy sphere vacuum characterized by  $G^1, G^2$ ; once we have  $G^1$  turned on, there is an instability towards turning on  $G^2$  also. Consequently, the mass of  $\chi_2$  can become negative, passing through zero. Another solution would amount to another instability towards a different vacuum with the same  $G^1$  turned on. As there are no vacua connected in this way to the maximally supersymmetric one, this is impossible. Finally, we note that there can be other light modes in other regions of parameter space, but since all we need here is that at  $N$  large,  $k \sim 1$  and the only VEV turned on is  $bG^1$ , there are no other light modes.

The question at this stage is how relevant are any of our effective field theories in the condensed matter context? Following [13], we now outline an argument to suggest that an appropriate answer is very. Beginning with the Hubbard model for spinless bosons, with a ground state where each site in the lattice is populated by an equal number of bosons, one can construct a discretized field  $\phi_i \sim \alpha_i a_i + \beta_i h_i^\dagger$ , where  $a_i^\dagger$  creates a *particle* above the ground state and  $h_i^\dagger$  creates a *hole*. One then obtains the relativistic Landau-Ginzburg action

$$S = \int d^3x (-|\partial_t \phi|^2 + v^2 |\vec{\nabla} \phi|^2 + (g - g_c)|\phi|^2 + u|\phi|^4), \quad (15)$$

in the continuum limit. For  $g < g_c$  we have an Abelian-Higgs system, i.e., a superconducting phase, while for  $g > g_c$  we have an insulator phase. At  $g = g_c$  (and temperature  $T = 0$ ) the model describes a conformal field theory. The systems described by the above model also have a quantum critical phase which opens up at nonzero temperature for a  $T$ -dependent window around  $g = g_c$ . This quantum critical phase is strongly coupled and

difficult to analyze using usual condensed matter methods and hence a good candidate for a holographic description.

Moreover, the Hubbard model is a drastic simplification of a real condensed matter system. The model was used to describe the quantum critical phase of (bosonic)  $^{87}\text{Rb}$  cold atoms on an optical lattice, but the description is believed to hold more generally for the quantum critical phase. For instance, high  $T_c$  superconductors have a *strange metal* phase that is believed to be of the same quantum critical phase type. In fact, a simple model for a solid with free electrons describes the qualitative features of the ABJM Abelianization. One can consider an electron at site  $i$  in the model coupled to an electron at site  $j$  to form a spinless boson  $\phi_{ij} = \bar{\psi}_i \psi_j$ . In two spatial dimensions there are  $\mathcal{O}(N^2)$  neighbours of maximum distance  $N$  away. It is not unreasonable to consider that the length between the sites has a maximum value  $N \geq |\vec{i} - \vec{j}|$ . We can write this field as  $\phi_{i'}^{ab}$ , where  $i'$  is at midpoint between  $i$  and  $j$ , and  $a, b$  correspond to  $1, 2, \dots, N$  in spatial directions  $x, y$ . If the normalized wave functions for  $\phi_{i'}^{ab}$  give probabilities for the existence of the pair  $|\phi_{i'}^{ab}|^2$ , and assuming rotational invariance so only rotationally invariant modes  $\psi(a)$ , thought of as eigenvalues of the matrices  $\phi^{ab}$ , are nonzero, we can consider a decaying solution  $|\psi(a)|^2 \propto N - a$ . This corresponds to the ABJM matrix  $G^2$ , with  $(G^2 G_2^\dagger)_{mn} = (N - m)\delta_{mn}$ . Indeed, in the ABJM model we have the field  $\chi_2(G^2)_{mn}$ , corresponding to  $b_{i'} \sim \sum_a \psi(a) \phi_{i'}^{aa}$ . The average distance in between sites is then

$$\langle a \rangle = \frac{\int |\psi(a)|^2 a (2\pi a da)}{\int |\psi(a)|^2 (2\pi a da)} = \frac{N}{2}, \quad (16)$$

consistent with the fact that there is a large distance between sites that couple, as is known to be the case.

Note that while the above fields are the only ones that are turned on, the system has, in principle, several more possibilities. For example, we can form more than one matrix scalar field, like the 4  $C^I$ 's of ABJM, by having more electrons at each site that can couple to form spinless bosons, as well as matrix fermions, by having two electrons at site  $i$  couple among themselves and with an electron at site  $j$ . We can also construct two Chern-Simons (topological) gauge fields by a generalization of the Abelian Chern-Simons case (see e.g., Ref. [14]) as follows: Consider two fermions at sites  $i$  and  $i'$  coupling to form  $\phi_{i'}^{aa'}$  at their midpoint  $i'$  and two fermions at sites  $j$  and  $j''$  coupling to form  $\phi_{j'}^{bb'}$  at their midpoint  $j'$ . Then the field

$$e\vec{a}(\vec{r}_{i'}) = \vec{\nabla}_{i'} \sum_{j' \neq i'} \alpha(\vec{r}_i - \vec{r}_j), \quad (17)$$

where  $\alpha(\vec{r}_i - \vec{r}_j)$  is the angle made by  $\vec{r}_i - \vec{r}_j$  with a fixed axis, corresponds to a Chern-Simons gauge field. The indices on the gauge field are the planar indices for the only variable above,  $\vec{r}_{i'} - \vec{r}_{j'}$  (changing  $\vec{r}_{i'}$  by itself just gives a harmless overall translation), as well as the discrete

choice for  $\vec{r}_{ii'}$  to belong to  $i'$  or  $j'$ , giving two gauge fields  $A$  and  $\hat{A}$ . The scalars  $\phi_{ij}^{ab}$  act as bifundamental with respect to them. It is clear then that, qualitatively at least, the model outlined above describes all the fields of ABJM, as well as the Abelianization.

In some sense, the three-dimensional Landau-Ginzburg model makes more sense as a dimensional reduction from four dimensions; the same is true of our Abelianization picture. The matrix  $G^1$ , one of the two matrices  $G^1, G^2$  that describe the fuzzy two-sphere, is multiplied by the constant  $|b|$ , so in a sense we have a *fuzzy circle* (limit of a fuzzy two-sphere), becoming classical at large  $N$ . The physical radius of a fuzzy sphere construction was argued in the literature on brane polarizations to be (see e.g., Ref. [11])

$$R_{\text{ph}}^2 = \frac{2}{N} \text{Tr}[X^I X_I^\dagger] = \frac{2}{N} \text{Tr}[C^I C_I^\dagger] 4\pi^2 l_p^3, \quad (18)$$

where  $l_p^3 = l_s^2 R_{11}$ . Assuming the same formula holds for the *fuzzy circle* case, and that like in the pure fuzzy sphere case, the 11th direction has radius  $R_{11} = R_{\text{ph}}/k$ , we obtain

$$R_{\text{ph}} = (N-1) l_s^2 \frac{4\pi^2 |b|^2}{k}. \quad (19)$$

The pure (massless) ABJM model corresponds to the IR limit of  $M2$  branes on  $\mathbb{R}^{2,1} \times \mathbb{C}^4/\mathbb{Z}_k$  and has as a gravity dual type IIA string theory on  $\text{AdS}_4 \times \mathbb{CP}^3$ . In the massive case, the spacetime for  $M2$ -brane propagation is more complicated [15], and the gravity dual even more so [15,16], so we will not reproduce the formulas here.

To summarize, in this Letter we have presented a consistent truncation of the ABJM model to a collective model of  $\mathcal{O}(N)$  modes out of the  $\mathcal{O}(N^2)$ , reducing to an Abelian Landau-Ginzburg model. We have also seen that we can map this process to a simple condensed matter model that reproduces the same general features. This provides a concrete step towards a well-defined AdS/CMT model, where there is a large  $N$  theory for the condensed matter

system, with a gravity dual, and yet the relevant physics is encoded in a simple Abelian model.

We thank the anonymous referees for useful comments on an earlier version of this article that led to several clarifying remarks. The work of H.N. is supported in part by CNPq Grant No. 301219/2010-9. J.M. acknowledges support from an NRF Incentive Funding for Rated Researchers Grant.

\*asadig@gmail.com

†jeff@nassp.uct.ac.za

‡nastase@ift.unesp.br

- [1] J.M. Maldacena, *Adv. Theor. Math. Phys.* **2**, 231 (1998).
- [2] S. S. Gubser, *Nucl. Phys.* **A830**, 657c (2009).
- [3] D. Berenstein, J. Maldacena, and H. Nastase, *J. High Energy Phys.* **04** (2002) 013.
- [4] J. A. Minahan and K. Zarembo, *J. High Energy Phys.* **03** (2003) 013.
- [5] S. A. Hartnoll, *Classical Quantum Gravity* **26**, 224002 (2009).
- [6] C. P. Herzog, *J. Phys. A* **42**, 343001 (2009).
- [7] A. Mohammed, J. Murugan, and H. Nastase (to be published).
- [8] O. Aharony, O. Bergman, D. L. Jafferis, and J. Maldacena, *J. High Energy Phys.* **10** (2008) 091.
- [9] J. Gomis, D. Rodriguez-Gomez, M. Van Raamsdonk, and H. Verlinde, *J. High Energy Phys.* **09** (2008) 113.
- [10] S. Terashima, *J. High Energy Phys.* **08** (2008) 080.
- [11] H. Nastase, C. Papageorgakis, and S. Ramgoolam, *J. High Energy Phys.* **05** (2009) 123.
- [12] H. Nastase and C. Papageorgakis, *SIGMA* **6**, 058 (2010).
- [13] S. Sachdev, *Annu. Rev. Condens. Matter Phys.* **3**, 9 (2012).
- [14] S. H. Simon, in *Composite Fermions*, edited by O. Heinonen (World Scientific, Singapore, 1998), Chap. 3; [arXiv: cond-mat/9812186](https://arxiv.org/abs/cond-mat/9812186).
- [15] A. Mohammed, J. Murugan, and H. Nastase, *Phys. Rev. D* **82**, 086004 (2010).
- [16] R. Auzzi and S. P. Kumar, *J. High Energy Phys.* **10** (2009) 071.